

4-1-2007

## Application of white noise calculus in evaluating the path integral in relativistic quantum mechanics

C C Bernido

J B Bornales

M V Carpio-Bernido

Follow this and additional works at: <https://repository.lsu.edu/cosa>



Part of the [Analysis Commons](#), and the [Other Mathematics Commons](#)

---

### Recommended Citation

Bernido, C C; Bornales, J B; and Carpio-Bernido, M V (2007) "Application of white noise calculus in evaluating the path integral in relativistic quantum mechanics," *Communications on Stochastic Analysis*: Vol. 1: No. 1, Article 12.

DOI: 10.31390/cosa.1.1.12

Available at: <https://repository.lsu.edu/cosa/vol1/iss1/12>

## APPLICATION OF WHITE NOISE CALCULUS IN EVALUATING THE PATH INTEGRAL IN RELATIVISTIC QUANTUM MECHANICS

C. C. BERNIDO, J. B. BORNALES, AND M. V. CARPIO-BERNIDO

ABSTRACT. The Hida-Streit method of evaluating the Feynman path integral is applied to relativistic quantum mechanical systems. A charged particle in a uniform magnetic field is taken as an example where the Green function for the Dirac particle is obtained.

### 1. Introduction

Since Feynman introduced his path integral formulation of quantum mechanics [11], various approaches aimed at providing a mathematically rigorous meaning to the path integral have been given. Notable among these would be the prodistributions of DeWitt-Morette [10], the oscillatory integrals of Albeverio and Høegh-Krohn [1] which explores the Fresnel integrals, and the white noise analysis approach of Hida and Streit [27]. In this paper, we shall consider the Hida-Streit approach which, through the years, has solved various classes of potentials [2, 3, 5, 6, 7, 8, 13, 18, 19, 21, 27] in nonrelativistic quantum mechanics. In white noise analysis, the velocity of Brownian motion  $B$ , i.e.,  $\omega(\tau) = dB/d\tau$ ,  $\tau \in \mathbb{R}$ , which are independent random variables at each point in time, form the coordinates of an infinite dimensional space [14]. We note that white noise analysis as applied here is also referred to as Hida calculus [14, 16, 20, 23], and differs from the use of white noise in Parisi-Wu stochastic quantization [25]. The calculus allows the generalization of concepts in finite dimensions to the infinite dimensional case, including differential operators and integral transforms such as Fourier and Fourier-Mehler transforms [16]. This makes it particularly suitable for the evaluation of the Feynman path integral. Here we extend the applicability of the Hida-Streit white noise path integral to solve systems in relativistic quantum mechanics where the Green function for the Dirac equation is obtained.

In Section 2 we briefly discuss a general procedure for obtaining the Dirac Green function using the path integral method. We then take as an example in Section 3 the case of a relativistic charged particle in a uniform magnetic field. In Section 4, the paths of this particle are parametrized in terms of the white noise variable  $\omega$  and the effective potential becomes similar in form to that of Lévy's stochastic area. As in the nonrelativistic case, the path integral can be evaluated with the

---

2000 *Mathematics Subject Classification.* Primary 60H40, 81S40; Secondary 81Q05.

*Key words and phrases.* white noise calculus, Feynman path integral, relativistic quantum mechanics.

help of the  $T$ -transform in white noise analysis from which we obtain the Dirac Green function. Concluding remarks are given in Section 5 where we cite other relativistic problems that may be solved using this approach.

## 2. Quantum Relativistic Systems

The Hida-Streit white noise path integral can be used to solve the Dirac equation for a spin  $\frac{1}{2}$  particle of mass  $m$  given by

$$\left(m - \widehat{M}\right) G(\mathbf{r}'', \mathbf{r}') = \delta(\mathbf{r}'' - \mathbf{r}'), \quad (2.1)$$

where we have defined an operator  $\widehat{M}$  in terms of the Dirac matrices  $\alpha$  and  $\beta$  of the form,

$$\widehat{M} = -\beta\boldsymbol{\alpha} \cdot (\mathbf{p} - e\mathbf{A}) - \beta V + \beta E. \quad (2.2)$$

Here  $V$  and  $\mathbf{A}$  are the scalar and vector potentials, respectively. By expressing the Green function  $G(\mathbf{r}'', \mathbf{r}')$  in Eq. (2.1) as,

$$G(\mathbf{r}'', \mathbf{r}') = \left(m + \widehat{M}\right) g(\mathbf{r}'', \mathbf{r}'). \quad (2.3)$$

we obtain an iterated Dirac equation,

$$\left(m^2 - \widehat{M}^2\right) g(\mathbf{r}'', \mathbf{r}') = \delta(\mathbf{r}'' - \mathbf{r}'). \quad (2.4)$$

The solution of Eq. (2.1) are also solutions of Eq. (2.4), but not conversely.

The Green functions,  $G(\mathbf{r}'', \mathbf{r}') = \langle \mathbf{r}'' | G | \mathbf{r}' \rangle$  and  $g(\mathbf{r}'', \mathbf{r}') = \langle \mathbf{r}'' | g | \mathbf{r}' \rangle$ , are matrix elements of the operators,  $G = (m - \widehat{M})^{-1}$ , and  $g = (m^2 - \widehat{M}^2)^{-1}$ , respectively. We can also write, for example operator  $g$ , in integral form using the relation (see, e.g., [11], Eq. 5-17),

$$\lim_{\epsilon \rightarrow 0} \left[ -(\theta + i\epsilon)^{-1} \right] = i \int_0^\infty \exp(i\theta\Lambda) d\Lambda. \quad (2.5)$$

Defining

$$H = \left(m^2 - \widehat{M}^2\right) / 2m, \quad (2.6)$$

we can express the operator  $g = (1/2m)(1/H)$  as (see also [9, 26])

$$g = (i/2m) \int_0^\infty \exp(-iH\Lambda) d\Lambda. \quad (2.7)$$

Taking the matrix element of Eq. (2.7),

$$g(\mathbf{r}'', \mathbf{r}') = (i/2m) \int_0^\infty \langle \mathbf{r}'' | \exp(-iH\Lambda) | \mathbf{r}' \rangle d\Lambda, \quad (2.8)$$

we observe that the integrand in Eq. (2.8) is analogous in form to a quantum propagator evolving in  $\Lambda$ -time with an effective Hamiltonian  $H$ . Following Feynman's prescription for handling quantum propagators [11] we can express the integrand  $\langle \mathbf{r}'' | \exp(-iH\Lambda) | \mathbf{r}' \rangle$  as a path integral. After evaluating the propagator  $\langle \mathbf{r}'' | \exp(-iH\Lambda) | \mathbf{r}' \rangle$ , we can integrate out  $\Lambda$  as in Eq. (2.8) to get  $g(\mathbf{r}'', \mathbf{r}')$ ,

and from Eq. (2.3), the Green function  $G(\mathbf{r}'', \mathbf{r}')$  for the Dirac equation can then be subsequently calculated. This procedure, therefore, enables us to arrive at a solution for the Dirac equation via the path integral method. We evaluate as an example a charged particle in a uniform magnetic field where white noise analytical methods are used.

### 3. Particle in a Uniform Magnetic Field

Let us consider an electron of charge  $e$  ( $e < 0$ ) and mass  $m$  subjected to a uniform magnetic field  $\mathbb{B}$  along the  $z$ -axis and represented by the vector potential  $\mathbf{A} = (1/2)(-\mathcal{B}y, \mathcal{B}x, 0)$ . Here we represent the magnetic field by  $\mathbb{B} = \mathcal{B}\hat{k}$  in order to distinguish it from our notation for the Brownian motion,  $B$ . With  $\widehat{M} = -\beta\alpha \cdot (\mathbf{p} - e\mathbf{A}) + \beta E$ , the operator  $H$  in Eq. (2.6) has the form,

$$H = (1/2m) \left[ (p_x + e\mathcal{B}y/2)^2 + (p_y - e\mathcal{B}x/2)^2 + p_z^2 \right] - (k_0^2/2m) - (e\sigma \cdot \mathbb{B}/2m), \quad (3.1)$$

where  $k_0^2 = E^2 - m^2$ . The spin part, with the magnetic field  $\mathbb{B}$  along the  $z$ -axis, is just  $-e\sigma \cdot \mathbb{B}/2m = -e\mathcal{B}\sigma_z/2m$ . We can then use Eq. (3.1) as the effective Hamiltonian in  $\langle \mathbf{r}'' | \exp(-iH\Lambda) | \mathbf{r}' \rangle$  of Eq. (2.8). Defining the operator  $S_z = (1/2)\sigma_z$ , then  $S_z$  acting upon an eigenspinor has the eigenvalues,  $s_i = \pm 1/2$ . Noting that  $|\mathbf{r}\rangle = |xyz\rangle |\chi\rangle$ , with  $|\chi\rangle$  a spinor, we can use  $\sum |s_i\rangle \langle s_i| = 1$ , to write the integrand in Eq. (2.8) as,

$$\begin{aligned} \langle \mathbf{r}'' | \exp(-iH\Lambda) | \mathbf{r}' \rangle &\equiv \sum_{s_i} \langle \chi | s_i \rangle \langle x''y''z'' | \exp(-iH_s\Lambda) | x'y'z' \rangle \langle s_i | \chi \rangle \\ &= \sum_{s_i} \eta_{s_i} \eta_{s_i}^+ \langle x''y''z'' | \exp(-iH_s\Lambda) | x'y'z' \rangle \end{aligned} \quad (3.2)$$

where  $\eta_{s_i} = \langle \chi | s_i \rangle$ . The coordinate part,  $\langle x''y''z'' | \exp(-iH_s\Lambda) | x'y'z' \rangle$ , has an effective Hamiltonian of the form (setting  $\gamma = e\mathcal{B}/2$ ),

$$H_s = (1/2m) \left[ (p_x + e\mathcal{B}y/2)^2 + (p_y - e\mathcal{B}x/2)^2 + p_z^2 \right] - (k_0^2/2m) - (2\gamma s_i/m). \quad (3.3)$$

We then evaluate the coordinate part as the path integral

$$\langle x''y''z'' | \exp(-iH_s\Lambda) | x'y'z' \rangle = \int \exp(iS) D[xyz] \quad (3.4)$$

for a system evolving with a time-like parameter  $\Lambda$ , and an action corresponding to Eq. (3.3) given by

$$S = \int_0^\Lambda \left[ \frac{1}{2}m(\dot{x}^2 + \dot{y}^2 + \dot{z}^2) + \gamma(xy - yx) + (k_0^2/2m) + (2\gamma s_i/m) \right] d\lambda. \quad (3.5)$$

The path integral, Eq. (3.4), with an action given by Eq. (3.5) can be written as,

$$\langle x''y''z'' | \exp(-iH_s\Lambda) | x'y'z' \rangle = e^{i[(k_0^2/2m) + (2\gamma s_i/m)]\Lambda} K(x'', y''; x', y') K(z'', z') \quad (3.6)$$

where the propagator along the  $z$ -coordinate is similar to the free particle case, i.e.,

$$\begin{aligned} K(z'', z') &= \int \exp \left[ (im/2) \dot{z}^2 \right] D[z] \\ &= (1/2\pi) \int \exp \left\{ [ik_z (z'' - z') - (ik_z^2/2m)] \Lambda \right\} dk_z. \end{aligned} \quad (3.7)$$

On the other hand, the propagator along the  $(x, y)$  axes is,

$$K(x''y''; x'y') = \int \exp \left\{ i \int_0^\Lambda \left[ \frac{1}{2} m (\dot{x}^2 + \dot{y}^2) + \gamma (xy - y\dot{x}) \right] d\lambda \right\} D[xy]. \quad (3.8)$$

This equation is similar to the propagator for a nonrelativistic charged particle in a uniform magnetic field [6, 11] which can be evaluated using white noise calculus.

#### 4. Evaluation in terms of White Noise

Let us now evaluate Eq. (3.8),  $K(x'', y''; x', y')$ , by parametrizing the paths in terms of Brownian motion variables as (setting,  $m = \hbar = c = 1$ ),

$$\begin{aligned} x(\Lambda) &= x' + B_x(\Lambda) \\ &= x' + \int_0^\Lambda \omega_x(\lambda) d\lambda, \end{aligned} \quad (4.1)$$

where  $\omega_x(\lambda) = dB_x(\lambda)/d\lambda$ , is the corresponding white noise variable. Similarly,

$$\begin{aligned} y(\Lambda) &= y' + B_y(\Lambda) \\ &= y' + \int_0^\Lambda \omega_y(\lambda) d\lambda, \end{aligned} \quad (4.2)$$

with  $\omega_y(\lambda) = dB_y(\lambda)/d\lambda$ . Here  $x'$  and  $y'$  are the fixed initial points. With this, the velocity of the particle, for instance along the  $x$ -direction, becomes,  $(dx/d\lambda) = \omega_x$ , and the kinetic part acquires the form,

$$\exp \left[ \frac{i}{2} \int_0^\Lambda (\dot{x}^2) d\lambda \right] = \exp \left[ \frac{i}{2} \int_0^\Lambda \omega_x(\tau)^2 d\lambda \right]. \quad (4.3)$$

The integration over all paths ( $\lim_{N \rightarrow \infty} \prod d[x_j]$  or  $d^\infty x$ ), on the other hand, leads to an integration over the Gaussian white noise measure of the form,  $d\mu(\omega_x) = N_\omega \exp \left( -\frac{1}{2} \int \omega_x(\tau)^2 d\tau \right) d^\infty \omega_x$ . The exponential factor  $\exp \left( -\frac{1}{2} \int \omega_x(\tau)^2 d\tau \right)$  in  $d\mu(\omega_x)$ , which is responsible for the Gaussian fall-off, spoils the direct correspondence between  $d^\infty x$  of the Feynman path integral and  $d^\infty \omega_x$  of the white noise measure. A better comparison can, therefore, be attained by multiplying  $d\mu(\omega_x)$  by a factor  $\exp \left( \frac{1}{2} \int \omega_x(\tau)^2 d\tau \right)$  to obtain the correspondence,  $d^\infty x \rightarrow d^\infty \omega$ . Together with Eq. (4.3), the translation of the integrand for the Feynman integral into the language of white noise analysis leads us to consider the Gauss kernel (for

the  $x$ -degree of freedom),

$$I_0 = \mathcal{N} \exp \left[ \left( \frac{i+1}{2} \right) \int_0^\Lambda \omega_x(\lambda)^2 d\lambda \right], \quad (4.4)$$

where  $\mathcal{N}$  is a normalization.

The kinetic part of Eq. (3.8) leads to a white noise functional which is a two-dimensional version of Eq. (4.4). We have,

$$I_{xy} = N \exp \left\{ \left( \frac{i+1}{2} \right) \int_0^\Lambda [\omega_x(\lambda)^2 + \omega_y(\lambda)^2] d\lambda \right\}. \quad (4.5)$$

On the other hand, the interaction part in Eq. (3.8) can be written as,

$$\begin{aligned} \exp \left[ i\gamma \int_0^\Lambda (xy - yx) d\lambda \right] &= \exp \left\{ i\gamma \int_0^\Lambda [x' + B_x(\lambda)] \omega_y(\lambda) d\lambda \right. \\ &\quad \left. - i\gamma \int_0^\Lambda [y' + B_y(\lambda)] \omega_x(\lambda) d\lambda \right\} \\ &= \exp \left\{ i\gamma \int_0^\Lambda [x' \omega_y(\lambda) - y' \omega_x(\lambda)] d\lambda \right\} \\ &\quad \times \exp(2i\gamma S_T), \end{aligned} \quad (4.6)$$

where  $S_T$  is Lévy's stochastic area given by

$$S_T = \frac{1}{2} \int_0^\Lambda [B_x(\lambda) dB_y(\lambda) - B_y(\lambda) dB_x(\lambda)]. \quad (4.7)$$

This may be handled by noting that the two-dimensional Brownian motion,  $B_x(\lambda)$  and  $B_y(\lambda)$ , can be realized on the probability space of a one-dimensional white noise [6, 15, 22], where

$$B_x(\lambda) = \int_0^\lambda \omega(\tau) d\tau; \quad dB_x(\lambda) = \omega(\lambda) d\lambda. \quad (4.8)$$

$$B_y(\lambda) = \int_{-\lambda}^0 \omega(\tau) d\tau; \quad dB_y(\lambda) = \omega(-\lambda) d\lambda. \quad (4.9)$$

Eqs. (4.8) and (4.9) enable us to write Eq. (4.7) as

$$\begin{aligned} S_T &= \int_{\mathbf{R}^2} \omega(\tau_1) F_S(\tau_1, \tau_2) \omega(\tau_2) d\tau_1 d\tau_2, \\ &= \langle \omega, F_S(\tau_1, \tau_2) \omega \rangle \end{aligned} \quad (4.10)$$

where

$$\begin{aligned} F_S(\tau_1, \tau_2) &= \frac{1}{4} [\chi_{[-\Lambda, 0]}(\tau_1) \chi_{[0, -\tau_1]}(\tau_2) + \chi_{[-\Lambda, 0]}(\tau_2) \chi_{[0, -\tau_2]}(\tau_1)] \\ &\quad - \frac{1}{4} [\chi_{[0, \Lambda]}(\tau_1) \chi_{[-\tau_1, 0]}(\tau_2) + \chi_{[0, \Lambda]}(\tau_2) \chi_{[-\tau_2, 0]}(\tau_1)]. \end{aligned} \quad (4.11)$$

The  $\chi_{[\alpha, \beta]}(\tau_j)$ ,  $j = 1, 2$ , with  $[\alpha, \beta]$  as the limits of integration, denotes the integration over  $\tau_j$  in Eq. (4.10).

On the other hand, the terms in the exponent of the kinetic part, Eq. (4.5), can be written as,

$$\begin{aligned} \left(\frac{i+1}{2}\right) \int_0^\Lambda [\omega_x(\tau)^2 + \omega_y(\tau)^2] d\tau &= \left(\frac{i+1}{2}\right) \int_0^\Lambda [\omega(\tau)^2 + \omega(-\tau)^2] d\tau \\ &= -\frac{1}{2} \int_{-\Lambda}^\Lambda \omega(\tau) K \omega(\tau) d\tau, \end{aligned} \quad (4.12)$$

where  $K = -(i+1)$ . Furthermore, to fix the endpoints  $x''$  and  $y''$ , we use the Donsker delta function  $\delta(x(\Lambda) - x'')$  and  $\delta(y(\Lambda) - y'')$ . where  $x(\Lambda)$  and  $y(\Lambda)$  are given by Eqs. (4.1) and (4.2). With Eqs. (4.6) to (4.12), we can now write Eq. (3.8) as,

$$\begin{aligned} K(x'', y''; x', y') &= \int I_{xy} \exp \left[ i\gamma \left( x' \int_{-\Lambda}^0 \omega(\lambda) d\lambda - y' \int_0^\Lambda \omega(\lambda) d\lambda \right) \right] \\ &\quad \times \exp \left( -\frac{1}{2} \langle \omega, L \omega \rangle \right) \\ &\quad \times \delta \left( x' - x'' + \int_0^\Lambda \omega(\lambda) d\lambda \right) \\ &\quad \times \delta \left( y' - y'' + \int_{-\Lambda}^0 \omega(\lambda) d\lambda \right) d\mu(\omega), \end{aligned} \quad (4.13)$$

where  $L = -4i\gamma F_S(\tau_1, \tau_2)$  and  $F_S(\tau_1, \tau_2)$  is given by Eq. (4.11). To evaluate the integration over the white noise measure, we write the delta functions in terms of their Fourier representation so that Eq. (4.13) becomes,

$$\begin{aligned} K(x'', y''; x', y') &= \frac{1}{(2\pi)^2} \int_{\mathbf{R}^2} d^2 \mathbf{p} \exp [i\mathbf{p} \cdot (\mathbf{x}' - \mathbf{x}'')] \\ &\quad \times \int N \exp \left[ i(p_x - \gamma y') \int_0^\Lambda \omega(\lambda) d\lambda \right] \end{aligned}$$

$$\begin{aligned}
& \times \exp \left[ i (p_y + \gamma x') \int_{-\Lambda}^0 \omega(\lambda) d\lambda \right] \\
& \times \exp \left( -\frac{1}{2} \langle \omega, K \omega \rangle \right) \exp \left( -\frac{1}{2} \langle \omega, L \omega \rangle \right) d\mu(\omega) \\
= & \frac{1}{(2\pi)^2} \int_{\mathbf{R}^2} d^2 \mathbf{p} \exp [i \mathbf{p} \cdot (\mathbf{x}' - \mathbf{x}'')] \int N \exp (i \langle \omega, \xi \rangle) \\
& \times \exp \left( -\frac{1}{2} \langle \omega, K \omega \rangle \right) \exp \left( -\frac{1}{2} \langle \omega, L \omega \rangle \right) d\mu(\omega),
\end{aligned} \tag{4.14}$$

where  $\mathbf{x} = (x, y)$ ,  $\mathbf{p} = (p_x, p_y)$ , and,  $\xi = (p_x - \gamma y') \chi_{[0, \Lambda]} + (p_y + \gamma x') \chi_{[-\Lambda, 0]}$ . We then note that the integration over  $d\mu(\omega)$  is just the  $T$ -transform of the white noise functional,  $\Phi = N \exp \left( -\frac{1}{2} \langle \omega, K \omega \rangle \right) \exp \left( -\frac{1}{2} \langle \omega, L \omega \rangle \right)$ , i.e.,

$$\begin{aligned}
(T\Phi)(\xi) &= \int N \exp (i \langle \omega, \xi \rangle) \exp \left( -\frac{1}{2} \langle \omega, K \omega \rangle \right) \exp \left( -\frac{1}{2} \langle \omega, L \omega \rangle \right) d\mu(\omega) \\
&= \left[ \det \left( 1 + L(1 + K)^{-1} \right) \right]^{-\frac{1}{2}} \exp \left[ -\frac{1}{2} \langle \xi, (1 + K + L)^{-1} \xi \rangle \right],
\end{aligned} \tag{4.15}$$

where we used Eq. (33) of ref. [6]. With  $K = -(i + 1)$ , we also have,

$$\begin{aligned}
\left[ \det \left( 1 + L(1 + K)^{-1} \right) \right]^{-\frac{1}{2}} &= [\det (1 + iL)]^{-\frac{1}{2}} \\
&= [\cos (\gamma \Lambda)]^{-1},
\end{aligned} \tag{4.16}$$

and [6],

$$\begin{aligned}
\langle \xi, (1 + K + L)^{-1} \xi \rangle &= (i/\gamma) \tan (\gamma \Lambda) [(p_x^2 + p_y^2) \\
&\quad + \gamma^2 (x'^2 + y'^2) + 2\gamma (p_y x' - p_x y')].
\end{aligned} \tag{4.17}$$

With Eqs. (4.15), (4.16) and (4.17), Eq. (4.14) becomes,

$$\begin{aligned}
K(x'', y''; x', y') &= \frac{\cos (\gamma \Lambda)^{-1}}{(2\pi)^2} \exp \left[ -\frac{i}{2} \gamma \tan (\gamma \Lambda) (x'^2 + y'^2) \right] \\
&\quad \times \int_{-\infty}^{+\infty} \exp \left\{ -i \tan (\gamma \Lambda) (2\gamma)^{-1} p_x^2 \right\} \\
&\quad \times \exp \{ i [(x' - x'') + \tan (\gamma \Lambda) y'] p_x \} dp_x \\
&\quad \times \int_{-\infty}^{+\infty} \exp \left\{ -i \tan (\gamma \Lambda) (2\gamma)^{-1} p_y^2 \right\} \\
&\quad \times \exp \{ i [(y' - y'') - \tan (\gamma \Lambda) x'] p_y \} dp_y.
\end{aligned} \tag{4.18}$$



The Gaussian integration in Eq. (4.18) over  $dp_x$  and  $dp_y$  can be performed and we obtain,

$$K(x'', y''; x', y') = \frac{\gamma}{2\pi i \sin(\gamma\Lambda)} \exp \left\{ \frac{i}{2} \gamma \cot(\gamma\Lambda) \left[ (x' - x'')^2 + (y' - y'')^2 \right] \right\} \\ \times \exp \{ i\gamma (x' y'' - x'' y') \}. \quad (4.19)$$

In terms of the Hermite polynomials  $H_n$ , Eq.(4.19) becomes,

$$K(x'', y''; x', y') = (\gamma/\pi) \exp \{ i\gamma (x' y'' - x'' y') \} \\ \times \exp \left\{ (-\gamma/2) \left[ (x'' - x')^2 + (y'' - y')^2 \right] \right\} \\ \times \sum_{n=0}^{\infty} \sum_{q=0}^n \frac{(-1)^{-n} 2^{-2n}}{q! (n-q)!} \\ \times H_{2q}(\sqrt{\gamma} (x'' - x')) H_{2(n-q)}(\sqrt{\gamma} (y'' - y)) \\ \times \exp \left[ -i \left( n + \frac{1}{2} \right) 2\gamma\Lambda \right]. \quad (4.20)$$

Using Eqs. (3.7) and (4.20), we can now write Eq. (3.6) as,

$$\langle x'' y'' z'' | \exp(-iH_s\Lambda) | x' y' z' \rangle \\ = (\gamma/2\pi^2) \exp \{ i\gamma (x' y'' - x'' y') \} \\ \times \exp \left\{ (-\gamma/2) \left[ (x'' - x')^2 + (y'' - y')^2 \right] \right\} \\ \times \int dk_z \exp [ik_z (z'' - z')] \sum_{n=0}^{\infty} \sum_{q=0}^n \frac{(-1)^{-n} 2^{-2n}}{q! (n-q)!} \\ \times H_{2q}(\sqrt{\gamma} (x'' - x')) H_{2(n-q)}(\sqrt{\gamma} (y'' - y')) \\ \times \exp [-i (k_z^2 - b^2) \Lambda/2m], \quad (4.21)$$

where  $b^2/2m = (k_0^2/2m) + [s_i - (n + \frac{1}{2})] (2\gamma/m)$  (putting back the mass  $m$ ). From Eqs. (3.2) and (4.21), the Green function for the iterated Dirac equation (2.8), appears as,

$$g(\mathbf{r}'', \mathbf{r}') = \left[ i\gamma/m (2\pi)^2 \int dk_z \sum_{s_i} \eta_{s_i} \eta_{s_i}^+ \exp [ik_z (z'' - z')] \right] \\ \times \exp \{ i\gamma (x' y'' - x'' y') \} \exp \{ (-\gamma/2) \\ \times \left[ (x'' - x')^2 + (y'' - y')^2 \right] \} \sum_{n=0}^{\infty} \sum_{q=0}^n \left[ \frac{(-1)^{-n} 2^{-2n}}{q! (n-q)!} \right] \\ \times H_{2q}(\sqrt{\gamma} (x'' - x')) H_{2(n-q)}(\sqrt{\gamma} (y'' - y')) \\ \times \int_0^{\infty} \exp [-i (k_z^2 - b^2) \Lambda/2m] d\Lambda. \quad (4.22)$$

The integration over  $\Lambda$  in Eq. (4.22) makes use of Eq. (2.5) which yields the result [12],

$$\begin{aligned}
g(\mathbf{r}'', \mathbf{r}') &= \left[ i\gamma/m (2\pi)^2 \right] \int dk_z \sum_{s_i} \eta_{s_i} \eta_{s_i}^+ \exp [ik_z (z'' - z')] \\
&\times \exp \{i\gamma (x' y'' - x'' y')\} \exp \{(-\gamma/2) \\
&\times [(x'' - x')^2 + (y'' - y')^2]\} \sum_{n=0}^{\infty} \sum_{q=0}^n \left[ \frac{(-1)^{-n} 2^{-2n}}{q! (n-q)!} \right] \\
&\times H_{2q}(\sqrt{\gamma} (x'' - x')) H_{2(n-q)}(\sqrt{\gamma} (y'' - y')) \\
&\times \left\{ (1/2m) (E^2 - m^2 - k_z^2) + \left[ s_i - \left( n + \frac{1}{2} \right) \right] (2\gamma/m) + i\varepsilon \right\}^{-1},
\end{aligned} \tag{4.23}$$

where  $\varepsilon \rightarrow 0$ . The discrete energy spectrum [17] can be obtained from the poles of the Green function and is given by,  $E^2 = m^2 + k_z^2 + [(2n+1) - 2s_i] e\mathcal{B}$ .

The Green function  $G(\mathbf{r}'', \mathbf{r}')$  for the first-order Dirac equation can be obtained with the help of Eqs. (2.3) and (4.23). We can rewrite  $\widehat{M}$  in Eq. (2.2) with  $V = 0$  as

$$\begin{aligned}
\widehat{M} &= -\beta\alpha_x (-i\partial/\partial x'' + e\mathcal{B}y''/2) - \beta\alpha_y (-i\partial/\partial y'' - e\mathcal{B}x''/2) \\
&+ (i\beta\alpha_z \partial/\partial z'') + \beta E
\end{aligned}$$

and using the recursion relations of the Hermite polynomials

$$\frac{dH_n(x)}{dx} = 2nH_{n-1}(x),$$

we obtain,

$$\begin{aligned}
G(\mathbf{r}'', \mathbf{r}') &= \left\{ m - \left( \frac{e\mathcal{B}}{2} \right) \beta\alpha_+ [(y'' - y') + i(x'' - x')] + \beta E - k_z \beta\alpha_z \right\} \\
&\times g(\mathbf{r}'', \mathbf{r}') + i\beta\alpha_x \bar{g} \sqrt{\frac{e\mathcal{B}}{2}} 4q \\
&\times H_{2q-1} \left( \sqrt{\frac{e\mathcal{B}}{2}} (x'' - x') \right) H_{2(n-q)} \left( \sqrt{\frac{e\mathcal{B}}{2}} (y'' - y') \right) \\
&+ i\beta\alpha_y \bar{g} \sqrt{\frac{e\mathcal{B}}{2}} 4(n-q) \\
&\times H_{2(n-q)-1} \left( \sqrt{\frac{e\mathcal{B}}{2}} (y'' - y') \right) H_{2q} \left( \sqrt{\frac{e\mathcal{B}}{2}} (x'' - x') \right),
\end{aligned}$$

where  $\alpha_+ = \alpha_y + i\alpha_x$ , and

$$\bar{g} = \left[ ie\mathcal{B}/2m (2\pi)^2 \right] \int dk_z \sum_{s_i} \eta_{s_i} \eta_{s_i}^+ \exp [ik_z (z'' - z')]$$

$$\begin{aligned} & \times \exp \left\{ (ie\mathcal{B}/2) (x'y'' - x''y') - (e\mathcal{B}/4) \left[ (x'' - x')^2 + (y'' - y')^2 \right] \right\} \\ & \times \sum_{n=0}^{\infty} \sum_{q=0}^n \left[ (-1)^{-n} 2^{-2n} / q! (n - q)! \right] \\ & \times \left\{ (1/2m) (E^2 - m^2 - k_z^2) + \left[ s_i - \left( n + \frac{1}{2} \right) \right] (e\mathcal{B}/m) + i\varepsilon \right\}^{-1}, \end{aligned}$$

as  $\varepsilon \rightarrow 0$ . We note here that an early attempt to provide a path integral treatment of a Dirac particle in a uniform magnetic field can be found in reference [24].

## 5. Conclusion

In this paper, we demonstrated how white noise calculus may be applied to solve relativistic quantum problems. By iterating the Dirac equation and expressing the corresponding Green function in terms of a path integral, the form of the effective Lagrangian enables one to take advantage of results available in the non-relativistic case. This similarity with the nonrelativistic problem is, in fact, the feature which allows us to extend the applicability of the Hida-Streit white noise path integral to solve quantum relativistic systems. Using the procedure presented in this paper, other relativistic systems can likewise be handled. Examples would be a Dirac particle on a circle, as well as a relativistic particle in an Aharonov-Bohm (AB) potential, and combinations thereof, such as a Dirac particle in the presence of a uniform magnetic field plus the AB potential [4]. We also note that, since we made use of the iterated Dirac equation, the procedure discussed in this paper can likewise be applied to treat spin 0 Klein-Gordon particles.

## References

- [1] Albeverio, S., and Høegh-Krohn, R.: *Mathematical Theory of Feynman Path Integrals*. Lecture Notes in Mathematics Vol. **523**. Springer-Verlag, Berlin, 1976.
- [2] Asai, N., Kubo, I., Kuo, H.-H.: Feynman integrals associated with Albeverio-Hoegh-Krohn and Laplace transform potentials; in: *Stochastic in Finite and Infinite Dimensions, in honor of G. Kallianpur*, eds. T. Hida et al. (Birkhaeuser, Boston, 2000) pp. 29 - 48.
- [3] Bernido, C. C., Carpio-Bernido, M. V.: Path integrals for boundaries and topological constraints: a white noise functional approach; *Jour. Math. Phys.* **43** (2002) 1728-1736.
- [4] Bornales, J.: Ph.D. Thesis (University of the Philippines, 2005).
- [5] Cunha, M., Drumond, C., Leukert, P., Silva, J. L., and Westerkamp, W.: The Feynman integrand for the perturbed harmonic oscillator as a Hida distribution; *Ann. Phys.* **4** (1995) 53 - 67.
- [6] de Falco, D., and Khandekar, D. C.: Applications of white noise calculus to the computation of Feynman integrals; *Stoch. Processes and Appl.* **29** (1988) 257-266.
- [7] de Faria, M., Potthoff, J. and Streit, L.: The Feynman integrand as a Hida distribution; *Jour. Math. Phys.* **32** (1991) 2123-2127.
- [8] de Faria M., Oliveira, M. J., and Streit, L.: Feynman integrals for non-smooth and rapidly growing potentials; *Jour. Math. Phys.* **46** (2005) Art. No.063505.
- [9] DeWitt, B. S.: *Dynamical Theory of Groups and Fields*. Gordon and Breach, New York, 1965.
- [10] DeWitt-Morette, C.: Feynman path integrals: I. Linear and affine techniques, II. The Feynman-Green function; *Comm. Math. Phys.* **37** (1974) 63-81.
- [11] Feynman, R. P., and Hibbs, A. R.: *Quantum Mechanics and Path Integrals*. McGraw-Hill, New York, 1965.

- [12] Gradshteyn, I. S., and Ryzhik, I. M.: *Table of Integrals, Series, and Products*, 5th ed. Academic Press, San Diego, 1994.
- [13] Grothaus, M., Khandekar, D. C., Silva, J. L. and Streit, L.: The Feynman integral for time-dependent anharmonic oscillators; *Jour. Math. Phys.* **38** (1997) 3278-3299.
- [14] Hida, T.: *Analysis of Brownian Functionals*, *Carleton Mathematical Lecture Notes* **13**. Princeton University book, 1975
- [15] Hida, T.: *Brownian Motion*. Springer-Verlag, New York, 1980.
- [16] Hida, T., Kuo, H. H., Potthoff, J., and Streit, L.: *White Noise. An Infinite Dimensional Calculus*; Kluwer, Dordrecht, 1993.
- [17] Itzykson, C., and Zuber, J. B.: *Quantum Field Theory*. McGraw-Hill, New York, 1985.
- [18] Khandekar, D. C., Streit, L.: Constructing the Feynman integrand; *Ann.Phys.(Leipzig)* **1** (1992) 46-55.
- [19] Kuna, T., Streit, L., Westerkamp, W.: Feynman integrals for a class of exponentially growing potentials; *Jour. Math. Phys.* **39** (1998) 4476-4491.
- [20] Kuo, H. H.: *White Noise Distribution Theory*. CRC, Boca Raton, FL, 1996.
- [21] Lascheck, A., Leukert, P., Streit, L., and Westerkamp, W.: Quantum mechanical propagators in terms of Hida distributions; *Rep. Math. Phys.* **33** (1993) 221-232.
- [22] Lévy, P.: *Le Mouvement Brownien*. Gauthier-Villars, Paris, 1954.
- [23] Obata, N.: *White Noise Calculus and Fock Space*, Lecture Notes in Mathematics, Vol. 1577. Springer, Berlin, 1994.
- [24] Papadopoulos, G. J., and Devreese, J. T.: Path integral solutions of the Dirac equation; *Phys. Rev. D***13** (1976) 2227.
- [25] Parisi, G. and Wu, Y. S.: Perturbation theory without gauge fixing; *Scientia Sinica* **24** (1981) 483-496.
- [26] Schwinger, J.: On gauge invariance and vacuum polarization; *Phys. Rev.* **82** (1951) 664.
- [27] Streit, L., and Hida, T.: Generalized Brownian functionals and the Feynman integral; *Stoch. Proc. Appl.* **16** (1983) 55-69.

CHRISTOPHER C. BERNIDO: RESEARCH CENTER FOR THEORETICAL PHYSICS, CENTRAL VISAYAN INSTITUTE FOUNDATION, JAGNA, BOHOL 6308, PHILIPPINES  
*E-mail address:* `cbernido@mozcom.com`

JINKY B. BORNALES: PHYSICS DEPARTMENT, MSU-ILIGAN INSTITUTE OF TECHNOLOGY, ILIGAN CITY 9200, PHILIPPINES  
*E-mail address:* `jbornales@yahoo.com`

M. VICTORIA CARPIO-BERNIDO: RESEARCH CENTER FOR THEORETICAL PHYSICS, CENTRAL VISAYAN INSTITUTE FOUNDATION, JAGNA, BOHOL 6308, PHILIPPINES  
*E-mail address:* `cbernido@mozcom.com`