

2-23-2001

## A note on doubles of groups

Nadia Benakli  
*Columbia University*

Oliver T. Dasbach  
*Columbia University*

Yair Glasner  
*Hebrew University of Jerusalem*

Brian Mangum  
*Barnard College*

Follow this and additional works at: [https://repository.lsu.edu/mathematics\\_pubs](https://repository.lsu.edu/mathematics_pubs)

---

### Recommended Citation

Benakli, N., Dasbach, O., Glasner, Y., & Mangum, B. (2001). A note on doubles of groups. *Journal of Pure and Applied Algebra*, 156 (2-3), 147-151. [https://doi.org/10.1016/S0022-4049\(99\)00176-0](https://doi.org/10.1016/S0022-4049(99)00176-0)

This Article is brought to you for free and open access by the Department of Mathematics at LSU Scholarly Repository. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Scholarly Repository. For more information, please contact [ir@lsu.edu](mailto:ir@lsu.edu).



ELSEVIER

Journal of Pure and Applied Algebra 156 (2001) 147–151

---

---

JOURNAL OF  
PURE AND  
APPLIED ALGEBRA

---

---

www.elsevier.com/locate/jpaa

## A note on doubles of groups

Nadia Benakli<sup>a</sup>, Oliver T. Dasbach<sup>a,b,1</sup>, Yair Glasner<sup>c</sup>, Brian Mangum<sup>d,\*</sup>

<sup>a</sup>*Department of Mathematics, Columbia University, New York, NY 10027, USA*

<sup>b</sup>*Mathematisches Institut, Heinrich-Heine-Universität, Abt. M.+M.D., D-40225 Düsseldorf, Germany*

<sup>c</sup>*Institute of Mathematics, The Hebrew University, Jerusalem 91904, Israel*

<sup>d</sup>*Department of Mathematics, Barnard College, Columbia University, New York, NY 10027, USA*

Received 28 February 1999; received in revised form 28 August 1999

Communicated by C.A. Weibel

---

### Abstract

Recently, Rips produced an example of a double of two free groups which has unsolvable generalized word problem. In this paper, we show that Rips's example fits into a large class of doubles of groups, each member of which contains  $F_2 \times F_2$  and therefore has unsolvable generalized word problem and is incoherent. © 2001 Elsevier Science B.V. All rights reserved.

MSC: 20E26; 20E06; 20F10; 20E08; 20F65

---

By a result of Mihailova [7] every group which contains the direct product of two (non-abelian) free groups does not admit a solution to the generalized word problem. Recently, Rips showed [9] — without using Mihailova's result — that the double of two free groups of rank 2 over a certain normal subgroup of index 3 does not have a generalized word problem solution. It was, however, already shown by Gersten [4] that such a group must contain  $F_2 \times F_2$ , and thus Rips's result follows from Gersten's and Mihailova's.

The purpose of this paper is to clarify the situation. We will show that Rips's example, and, in fact, any double of a virtually free finitely generated group over a finite index subgroup, is virtually the direct product of two free groups. Thus, Rips's proof yields a new proof for Mihailova's result. Furthermore, we will show that a more general construction of a double of a group must always contain  $F_2 \times F_2$ .

---

\* Corresponding author.

*E-mail addresses:* benakli@math.columbia.edu (N. Benakli), kasten@math.columbia.edu (O.T. Dasbach), yairgl@math.huji.ac.il (Y. Glasner), mangum@math.columbia.edu (B. Mangum).

<sup>1</sup> Work supported by the Deutsche Forschungsgemeinschaft (DFG).

One might naively think that every finitely presented group which admits a solution to the word problem but which is neither coherent nor does it admit a solution to the generalized word problem must contain  $F_2 \times F_2$ .

A counterexample follows from a construction of Rips [8]. He constructed for every finitely presented group  $G$  a short exact sequence

$$1 \rightarrow K \rightarrow H \rightarrow G \rightarrow 1,$$

so that  $H$  is a finitely presented word hyperbolic group, and  $K$  is — as a group — finitely generated. (In fact, Rips's construction yields for every  $\lambda$  a finitely presented group  $H$  satisfying the small cancellation condition  $C'(\lambda)$ . In particular,  $\lambda$  can be chosen so that  $H$  is word hyperbolic.)

It follows that if either  $G$  is not coherent or  $G$  does not admit a solution to the generalized word problem, then the same is true for  $H$ .

Since  $H$  is word hyperbolic, it has a solution to the word problem but cannot contain  $F_2 \times F_2$  or even  $\mathbb{Z} \times \mathbb{Z}$ .

Rips's construction starts with a group  $G$  that is already known to be incoherent and which does not admit a solution to the generalized word problem. Very recently, however, Wise [11] gave a construction of an incoherent word hyperbolic group without referring to a group which contains  $F_2 \times F_2$ .

**Definition 1.1.** For a group  $G$  one defines the following properties:

1. The group  $G$  is coherent if every finitely generated subgroup is finitely presented.
2. It is locally extended residually finite (LERF) or subgroup separable if for every finitely generated subgroup  $H$  of  $G$  and every  $g \in G$ ,  $g \notin H$  there is a subgroup of finite index in  $G$  that contains  $H$  but does not contain  $g$ . In particular, a LERF group is residually finite.
3. The group  $G$  admits a solution to the generalized word problem or occurrence problem if there is an algorithm that decides for every element  $g \in G$  and every finitely generated subgroup  $H \subset G$  whether  $g$  is in  $H$  or not.

As for a finitely presented group  $G$  residual finiteness implies that the word problem is solvable, it is easy to see that if  $G$  is LERF then the generalized word problem for  $G$  is solvable.

Moreover, it follows more or less directly from the definition that if a group  $G$  has one of these properties then so does every subgroup and every finite extension of  $G$ .

An important obstacle for a group to be coherent, LERF or to admit a solution to the generalized word problem is given by

**Proposition 1.2.** *The group  $F_2 \times F_2$  is neither coherent [5] nor does it admit a solution to the generalized word problem (Mihailova, see [6,7] and compare with [5]) and is thus not LERF. Therefore every group which contains  $F_2 \times F_2$  has also none of these properties.*




Fig. 1. The covering of graphs corresponding to  $K \hookrightarrow L$ .

Let  $G$  be a group and  $H$  a subgroup. By the double of  $G$  over  $H$  we mean the group that we get by the free product of  $G$  with itself amalgamated over  $H$ . Here, we will show that a large family of group doubles contain  $F_2 \times F_2$  as a subgroup.

**Lemma 1.3.** *Let  $G$  be a group and  $H < G$ . Let  $L = G *_H G$  be the double of  $G$  over  $H$ . Let  $\phi_1 : L \rightarrow G$  be the homomorphism obtained by identifying the two factors of  $L$ . Then  $K = \ker(\phi_1)$  is a free group. If  $H$  is of finite index in  $G$  then the rank of  $K$  is  $[G : H] - 1$ .*

**Proof.** The group  $L$  is an amalgam of groups and has a natural action on its Bass–Serre tree  $T$  (which is locally finite iff  $[G : H] < \infty$ ). By definition  $K$  intersects trivially the two copies of  $G$  in  $L$ , and since  $K$  is normal, it intersects trivially all their conjugates, which are all the vertex stabilizers. We conclude that  $K$  acts freely and effectively on  $T$ . (Note that  $L$  does not necessarily act effectively on  $T$ .) Therefore,  $K \cong \pi_1(K \backslash T, \{pt\})$  is a free group (see also [10]). Now,  $\phi_1$  is a surjective homomorphism and therefore the index of  $K$  in  $L$  is  $[L : K] = \#|G|$  (note that  $G$  might be infinite). The graph  $K \backslash T$  is therefore a  $\#|G|$ -fold cover of the edge of groups corresponding to the amalgam  $L$  (in the sense of Bass [1]). The only possibility of such a covering graph is shown in Fig. 1. The rank of  $K$  as a free group is now easily verified as  $K$  is the fundamental group of the covering graph in Fig. 1.  $\square$

The idea for the proof of the following theorem was inspired by the notion of reducible lattices acting on products of trees, defined by Burger et al. [3] and Burger and Mozes [2].

**Theorem 1.4.** *Let  $G$  be a group,  $H < G$  be a subgroup with  $3 \leq [G : H] \leq \infty$ .*

1. *Assume that there exists a subgroup  $N < H$  with the following properties:*
  - (a)  *$N$  contains a non-abelian, free subgroup.*
  - (b)  *$N$  is normal in  $G$ .*

*In particular, this assumption is fulfilled if  $G$  contains a non-abelian free group and  $H$  has finite index in  $G$ . Then the double  $L \stackrel{\text{def}}{=} G *_H G$  contains  $F_2 \times F_2$  as a*

subgroup and is thus neither coherent nor LERF nor does it admit a solution to the generalized word problem.

2. Assume  $G$  is a finitely generated virtually (non-abelian) free group, and  $H$  has finite index in  $G$ . The double  $G *_H G$  is virtually the direct product  $F_{r_1} \times F_{r_2}$  of two non-abelian free groups of finite rank. Moreover,  $F_2 \times F_2$  and  $G *_H G$  are commensurable.

**Proof.** The group  $N$  is normal in both copies of  $G$  in  $L$  and is therefore normal in  $L$ . We consider the following homomorphism of  $L$  into a product:

$$\phi = (\phi_1, \phi_2) : L \rightarrow G \times L/N. \tag{1.1}$$

Here  $\phi_1$  is induced by identifying the two factors of  $L$ , and  $\phi_2$  is just the projection to the quotient. First note that  $\ker(\phi_2) = N$  and that  $N$ , as a subgroup of  $G$ , injects into  $G$  under  $\phi_1$ . Therefore,  $\ker(\phi) = \ker(\phi_1) \cap \ker(\phi_2)$  is trivial. So  $\phi$  is injective and it will be enough to prove that  $\phi(L)$  contains a copy of  $F_2 \times F_2$ . Define  $K_i \stackrel{\text{def}}{=} \phi_i(\ker(\phi_{3-i}))$ ,  $i = 1, 2$ , and look at the embedding  $K_1 \times K_2 \hookrightarrow \phi(L)$ . We will show that each one of the  $K_i$  contains a non-abelian free group and thus conclude the proof of Part 1. The group  $K_1 = N$  contains a non-abelian free group by assumption. In order to deal with the other factor, consider the map  $\bar{\phi}_1 : G/N *_H/N G/N \rightarrow G/N$  (the homomorphism obtained by identifying the two copies of  $G/N$ ). The fact that  $\ker(\phi_1) \cap \ker(\phi_2) = \langle e \rangle$  and the commutativity of the following diagram:

$$\begin{array}{ccc} L \cong G *_H G & \xrightarrow{\phi_2} & L/N \cong G/N *_H/N G/N \\ \phi_1 \downarrow & & \bar{\phi}_1 \downarrow \\ G & \longrightarrow & G/N \end{array}$$

imply that  $K_2 = \ker(\bar{\phi}_1)$ . Now, by Lemma 1.3,  $K_2$  is a free group of rank

$$[G/N : H/N] - 1 = [G : H] - 1.$$

The assumption that this number is at least 2 shows that  $K_2$  is non-abelian.

In the case, where  $G$  is a virtually non-abelian free group,  $G *_H G$  contains  $K_1 \times K_2$  as a subgroup of finite index where  $K_1$  is virtually free and  $K_2$  is free. Any non-abelian free group of finite rank is isomorphic to a subgroup of finite index in  $F_2$ . Therefore,  $G *_H G$  and  $F_2 \times F_2$  are commensurable. This concludes the theorem.  $\square$

**Remark.** In this theorem, if we take  $G$  to be  $F_2$  and take  $H$  to be the normal subgroup in Rips’s example [9], then the double  $G *_H G$  and  $F_2 \times F_2$  are shown to be commensurable. Rips has shown that  $G *_H G$  has no solution to the generalized word problem. Therefore,  $F_2 \times F_2$  has the same property. This gives a new proof of Mihailova’s result.

## Acknowledgements

This paper was written while the second and third author were guests at Columbia University. They would like to thank Columbia and especially Joan Birman and Hyman Bass for their hospitality. The authors would also like to thank Michael Kapovich for bringing to their attention Rips's construction of incoherent word hyperbolic groups mentioned in the introduction.

## References

- [1] H. Bass, Covering theory for graphs of groups, *J. Pure Appl. Algebra* 89 (1–2) (1993) 3–47.
- [2] M. Burger, S. Mozes, Finitely presented simple groups and products of trees, *C. R. Acad. Sci. Paris Sér. I Math.* 324 (7) (1997) 747–752.
- [3] M. Burger, S. Mozes, R.J. Zimmer, Linear representations and arithmeticity for lattices in  $Aut(T_n) \times Aut(T_m)$ , in preparation.
- [4] S.M. Gersten, Coherence in doubled groups, *Comm. Algebra* 9 (18) (1981) 1893–1900.
- [5] F.J. Grunewald, On some groups which cannot be finitely presented, *J. London Math. Soc.* (2) 17 (3) (1978) 427–436.
- [6] C.F. Miller III, On Group-theoretic Decision Problems and Their Classification, *Annals of Mathematics Studies*, vol. 68, Princeton University Press, Princeton, NJ, 1971.
- [7] K.A. Mihailova, The occurrence problem for direct products of groups, *Mat. Sbornik* 70 (1966) 241–251.
- [8] E. Rips, Subgroups of small cancellation groups, *Bull. London Math. Soc.* 14 (1) (1982) 45–47.
- [9] E. Rips, On a double of a free group, *Israel J. Math.* 96 (Part B) (1996) 523–525.
- [10] J.-P. Serre, *Trees*, Springer, Berlin, 1980 (translated from the French by John Stillwell).
- [11] D.T. Wise, Incoherent negatively curved groups, *Proc. Amer. Math. Soc.* 126 (4) (1998) 957–964.