

1-1-2019

A highly contiguous reference genome for northern bobwhite (*Colinus virginianus*)

Jessie F. Salter
Louisiana State University

Oscar Johnson
Louisiana State University

Norman J. Stafford

William F. Herrin

Darren Schilling
Franklinton Veterinary Clinic

See next page for additional authors

Follow this and additional works at: https://repository.lsu.edu/biosci_pubs

Recommended Citation

Salter, J., Johnson, O., Stafford, N., Herrin, W., Schilling, D., Cedotal, C., Brumfield, R., & Faircloth, B. (2019). A highly contiguous reference genome for northern bobwhite (*Colinus virginianus*). *G3: Genes, Genomes, Genetics*, 9 (12), 3929-3932. <https://doi.org/10.1534/g3.119.400609>

This Article is brought to you for free and open access by the Department of Biological Sciences at LSU Scholarly Repository. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Scholarly Repository. For more information, please contact ir@lsu.edu.

Authors

Jessie F. Salter, Oscar Johnson, Norman J. Stafford, William F. Herrin, Darren Schilling, Cody Cedotal, Robb T. Brumfield, and Brant C. Faircloth

A Highly Contiguous Reference Genome for Northern Bobwhite (*Colinus virginianus*)

Jessie F. Salter,* Oscar Johnson,* Norman J. Stafford, III,[†] William F. Herrin, Jr.,[‡] Darren Schilling,[§] Cody Cedotal,** Robb T. Brumfield,* and Brant C. Faircloth*¹

*Museum of Natural Science and Department of Biological Sciences, Louisiana State University, Baton Rouge, Louisiana, [†]39731 LA-10, Franklinton, Louisiana, [‡]15531 Chanove Avenue, Greenwell Springs, Louisiana, [§]Franklinton Veterinary Clinic, 525 11th Avenue, Franklinton, Louisiana, and **Louisiana Department of Wildlife and Fisheries, P.O. Box 98000, Baton Rouge, Louisiana

ORCID IDs: 0000-0002-1764-5708 (O.J.); 0000-0003-2307-0688 (R.T.B.); 0000-0002-1943-0217 (B.C.F.)

ABSTRACT Northern bobwhites (*Colinus virginianus*) are small quails in the New World Quail family (Odontophoridae) and are one of the most phenotypically diverse avian species. Despite extensive research on bobwhite ecology, genomic studies investigating the evolution of phenotypic diversity in this species are lacking. Here, we present a new, highly contiguous assembly for bobwhites using tissue samples from a vouchered, wild, female bird collected in Louisiana. By performing a *de novo* assembly and scaffolding the assembly with Dovetail Chicago and HiC libraries and the HiRise pipeline, we produced an 866.8 Mb assembly including 1,512 scaffolds with a scaffold N50 of 66.8 Mb, a scaffold L90 of 17, and a BUSCO completeness score of 90.8%. This new assembly represents approximately 96% of the non-repetitive and 84% of the entire bobwhite genome size, greatly improves scaffold lengths and contiguity compared to an existing draft bobwhite genome, and provides an important tool for future studies of evolutionary and functional genomics in bobwhites.

KEYWORDS

Colinus virginianus
bobwhites
Dovetail
Chicago
HiC
genome
assembly

INTRODUCTION

Northern bobwhites (*Colinus virginianus*; hereafter bobwhites) are widely distributed quails primarily found in pine woodlands and grasslands of the eastern United States and Mexico (Brennan 1999). Bobwhites hold a significant place in the cultural heritage of both countries due to their status as popular game birds (Bent 1963; Burger *et al.* 1999), and they have also played a significant role in biological research because they are one of the most intensively studied birds in the world (Guthery 1997). Bobwhites are remarkably polytypic: there are 22 subspecies recognized by male plumage (Brennan 1999) – a larger number of subspecies than 99% of all other birds (Dickinson and Remsen 2013). Although bobwhite ecology research has been

extensive, the evolutionary relationships between bobwhite subspecies remain murky (Ellsworth *et al.* 1989; Evans *et al.* 2009; Eo *et al.* 2009; Williford 2013; Williford *et al.* 2014, 2016) and the genetic basis of phenotypic diversity in bobwhites has been largely unstudied (but see Cole *et al.* 1949).

Identifying genotypes associated with specific phenotypes increasingly relies on whole genome sequencing, particularly for investigating the genetic basis of phenotypic differences in non-model organisms (Ellegren 2014). The first draft genome assembly for bobwhites (GCA_000599485.1; hereafter Cv_TX_1.1) was generated from an unvouchered wild female bird from Texas (Halley *et al.* 2014). Cv_TX_1.1 used small and medium insert paired-end (PE) and mate pair (MP) libraries to produce a 1.172 Gb genome assembly with 77x coverage, 50% of the assembly in scaffolds of at least 45.5 Kbp in size (N50), and 90% of the assembly in 25,837 scaffolds (L90, Halley *et al.* 2014). Sequencing of additional PE and MP libraries from the same bird were used to generate a second assembly (GCA_000599465.2; hereafter Cv_TX_2.0), which yielded a 1.5-fold increase in coverage (122x), a 45-fold improvement in N50 (2.042 Mb), and a nearly threefold decrease in L90 (8,990 scaffolds; Oldeschulte *et al.* 2017). Although Cv_TX_2.0 was a marked improvement over Cv_TX_1.1, the scaffolds remained relatively short, which can hinder identification of structural variants (Domyan *et al.* 2014). Recent studies in

Copyright © 2019 Salter *et al.*

doi: <https://doi.org/10.1534/g3.119.400609>

Manuscript received August 6, 2019; accepted for publication October 8, 2019; published Early Online October 14, 2019.

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Supplemental material available at figshare: <https://doi.org/10.25387/g3.9273542>.

¹Corresponding author: Life Sciences Building, Louisiana State University, Baton Rouge, LA 70803. E-mail: brant@faircloth-lab.org

birds and other taxa have demonstrated the importance of structural variants in generating morphological diversity within closely-related taxa (Lamichhaney *et al.* 2016; Tuttle *et al.* 2016; Vijay *et al.* 2016), highlighting the need for highly contiguous genome assemblies in phenotype-genotype studies (Wellenreuther and Bernatchez 2018).

Here, we describe Cv_LA_1.0, a new assembly for bobwhites using DNA extracted from a vouchered, wild female bird collected in Louisiana. To generate this assembly, we scaffolded contigs from small insert libraries with reads from Chicago (Putnam *et al.* 2016) and HiC (Lieberman-Aiden *et al.* 2009) methodologies and the HiRise assembly pipeline (Dovetail Genomics, LLC). The resulting Cv_LA_1.0 assembly is highly contiguous and represents a 32-fold increase in N50 and 528-fold decrease in L90 relative to Cv_TX_2.0 (Oldeschulte *et al.* 2017).

METHODS AND MATERIALS

Specimen collection and DNA extraction

We collected blood, liver, and other tissues for direct storage in liquid nitrogen from a wild, female bird legally harvested at Sandy Hollow Wildlife Management Area (30.827 N, 90.397 W) in Tangipahoa Parish, Louisiana. After tissue collection, we prepared a specimen for the LSU Museum of Natural Science (LSUMNS) Collection of Birds (LSUMZ 197699), and we stored tissue samples from this specimen in the LSUMNS Collection of Genetic Resources (LSUMZ B-91918). We shipped blood and liver to Dovetail Genomics, LLC (Scotts Valley, CA) where Dovetail Staff performed DNA extraction, library preparation, sequencing, and assembly steps. Dovetail staff extracted high molecular weight (HMW) DNA from tissues using the Blood and Cell Culture Midi Kit (Qiagen, GmbH) following the manufacturer's protocol. Mean fragment length of the extracted DNA was 85 kb.

Short-insert library preparation, sequencing, and assembly

Dovetail staff randomly fragmented extracted DNA by sonication using a Bioruptor Pico (Diagenode, Inc.) and 7 cycles of sonication for 15 sec followed by 90 sec of rest. Dovetail staff then prepared a sequencing library by inputting fragmented DNA to the TruSeq DNA PCR-Free Library Preparation Kit (Illumina, Inc.) following the manufacturer's protocol. Resulting libraries were sequenced on an Illumina HiSeq X platform using paired-end (PE) 150 bp sequencing. Resulting data were trimmed for low-quality bases and adapter contamination using Trimmomatic (Bolger *et al.* 2014) and used to assemble scaffolds with Meraculous v2.2.5 (Chapman *et al.* 2011). Before assembly, Dovetail staff used Jellyfish (Marçais and Kingsford 2011) with in-house software similar to GenomeScope (Vurture *et al.* 2017) to profile the short insert reads at a variety of k-mer values (25, 55, 85, 109), estimate genome size, and fit negative binomial models to the data. The resulting profiles suggested a k-mer size of 55 was optimal for assembly, and Dovetail staff assembled contigs using Meraculous with a k-mer size of 55, a minimum k-mer frequency of 12, and the diploid nonredundant haplotigs mode.

Chicago library preparation and sequencing

Following *de novo* assembly with Meraculous, Dovetail staff prepared a single, proprietary "Chicago" library following the methods described in Putnam *et al.* (2016). Briefly, they reconstituted ~500 ng of HMW genomic DNA into chromatin *in vitro* and fixed the reconstituted DNA with formaldehyde. Then, they digested fixed chromatin with DpnII, filled in 5' overhangs with biotinylated nucleotides, and ligated free, blunt ends. After ligation, they reversed crosslinks and purified the DNA from protein. Dovetail staff treated purified DNA to remove

Table 1 Metrics estimated using QCAST, UCSC Browser Utilities, and GNU Coreutils for *Colinus virginianus* genome assembly Cv_LA_1.0 (this manuscript) and comparison to a different assembly of a different individual, Cv_TX_2.0 (GCA_000599465.2; Oldeschulte *et al.* 2017), from the same species

	Cv_LA_1.0	Cv_TX_2.0
Contigs	1,512	42,369
Largest contig (bp)	180,865,729	14,292,544
Total length (bp)	866,266,924	1,254,146,751
N50 (bp)	66,809,948	2,042,136
N75 (bp)	22,391,474	65,386
N90 (bp)	13,127,921	11,797
L50	4	150
L75	10	1,080
L90	17	8,989
GC (%)	41.2	42.7
# N's	11,810,287	119,897,618
# N's per 100 kbp	1,363.4	9,560.1

biotin that was not internal to ligated fragments and sheared the resulting DNA to ~350 bp mean fragment size using a Bioruptor Pico. Dovetail staff then prepared sequencing libraries from the sheared DNA using NEBNext Ultra enzymes (New England Biolabs, Inc.) and Illumina-compatible adapters. They isolated biotin-containing fragments using streptavidin beads before PCR enrichment of each library. Dovetail staff then sequenced amplified libraries on an Illumina HiSeq X platform using PE 150 reads to approximately 90X depth.

Dovetail HiC library preparation and sequencing (multiple libraries)

Dovetail staff also prepared two Dovetail HiC libraries following the procedures outlined in Lieberman-Aiden *et al.* (2009). Briefly, for each library, Dovetail staff used formaldehyde to fix chromatin in place in the nucleus. They extracted and digested fixed chromatin with DpnII, filled in the 5' overhangs with biotinylated nucleotides, and ligated free blunt ends. After ligation, Dovetail staff reversed crosslinks and purified the DNA from protein. They treated the purified DNA to remove biotin that was not internal to ligated fragments and sheared the DNA to ~350 bp mean fragment size using a Bioruptor Pico. Dovetail staff then prepared sequencing libraries using NEBNext Ultra enzymes and Illumina-compatible adapters. They isolated biotin-containing fragments using streptavidin beads before PCR enrichment of each library and sequenced the resulting libraries on an Illumina HiSeq X Platform using PE 150 reads to approximately 60X depth.

Assembly scaffolding with HiRise

To scaffold and improve the bobwhite assembly, Dovetail staff input the *de novo* assembly from Meraculous, along with shotgun reads, Chicago library reads, and Dovetail HiC library reads into HiRise (April 2017 version), a software pipeline designed for this purpose (Putnam *et al.* 2016). Using HiRise, Dovetail staff conducted an iterative analysis. First, they aligned shotgun and Chicago library sequences to the draft contig assembly using a modified SNAP read mapper (<http://snap.cs.berkeley.edu>). Second, they analyzed the separations of Chicago read pairs mapped within draft scaffolds to produce a likelihood model for genomic distance between read pairs, and they used this model to: identify and break putative misjoins, score prospective joins, and make joins above a threshold. Finally, after aligning and scaffolding the draft assembly using the Chicago data, Dovetail staff aligned and scaffolded the Chicago assembly using Dovetail HiC library sequences following the same method. After scaffolding, Dovetail staff

■ **Table 2** Genome completeness estimated using single copy orthologs (BUSCO v3) from *Colinus virginianus* assembly Cv_LA_1.0 (this manuscript) compared to a different assembly, Cv_TX_2.0 (GCA_000599465.2; Oldeschulte *et al.* 2017) from the same species

	Cv_LA_1.0		Cv_TX_2.0	
	Count	Percentage	Count	Percentage
Complete BUSCOs	4,461	90.8%	4,493	91.4%
Complete and single-copy BUSCOs	4,416	89.8%	4,435	90.2%
Complete and duplicated BUSCOs	45	0.9%	58	1.2%
Fragmented BUSCOs	170	3.5%	248	5.0%
Missing BUSCOs	284	5.8%	174	3.5%
Total BUSCO groups searched	4,915	—	4,915	—

used the short-insert sequences to close remaining gaps between contigs where possible.

Assembly polishing, contiguity statistics, and BUSCO analyses

After receiving the assembly from Dovetail, we aligned the short insert data back to the scaffolded assembly using bwa v0.7.17-r1188 (Li and Durbin 2009) and samtools v1.9 (Li *et al.* 2009) and polished the scaffolds using Pilon v1.23 (Walker *et al.* 2014) on a 48-core, 1.5 TB RAM compute node with default parameters. After polishing, we computed contiguity statistics of our scaffolded assembly as well as the Cv_TX_2.0 assembly (Oldeschulte *et al.* 2017) using QUAST v5.0.2 (Mikheenko *et al.* 2018), UCSC Browser Utilities (Kent *et al.* 2002), and GNU Coreutils (<https://www.gnu.org/software/coreutils>), and we performed BUSCO analyses against both genomes using BUSCO v3.1.0 (Waterhouse *et al.* 2018) and the Aves Data Set (aves_odb9).

Data availability

Data from all sequencing runs and the final assembly, Cv_LA_1.0, are available from NCBI BioProject (PRJNA454855). Short-insert, Chicago, and HiC reads are also available from the NCBI SRA (SRP215501), and the assembly is available from NCBI Genome using the accession VONY000000000. The version described in this manuscript is VONY01000000. Outputs from QUAST and BUSCO analyses are available as supplemental files from figshare. Supplemental material available at figshare: <https://doi.org/10.25387/g3.9273542>.

RESULTS AND DISCUSSION

Sequencing of short-insert libraries produced 441.8 million read pairs with an average insert size of 428 bp. Analysis of the k-mer histogram at the optimal value of 55 suggested the genome size was 1.0 Gb, and the estimated Q20 read depth for this genome size was approximately 118x. Meraculous assembly using a k-mer value of 55 produced 23,275 contigs having a total length of 853.1 Mb and an N50 of 113.6 Kb. These contigs were joined by Meraculous into 14,482 scaffolds totaling 854.1 Mb in length with an N50 of 176.8 Kb and a L90 of 5,343 scaffolds. The longest Meraculous scaffold was 1.6 Mb. Meraculous estimated that the assembled contigs comprised 96% of the estimated, non-repetitive genome size and 84% of the entire genome size.

Chicago library sequencing produced 303 million read pairs, and the estimated physical coverage (the number of read pairs with inserts between 1 and 100 Kb) spanning each position in the Meraculous assembly was 382.2. HiRise made 12,824 joins and one break to the Meraculous assembly to produce a Chicago assembly including 1,659 scaffolds totaling 866.68 Mb in length with an N50 of 15.5 Mb and a L90 of 53 scaffolds. The longest Chicago scaffold was 86 Mb.

HiC library sequencing produced 111 million read pairs for Library 1 and 95 million read pairs for Library 2, and the estimated physical coverage (the number of read pairs with inserts between 10 and

10,000 kb) spanning each position in the Chicago assembly was 38,615. HiRise made 147 joins and zero breaks to the Chicago-scaffolded assembly to produce a HiC assembly including 1,512 scaffolds totaling 866.8 Mb in length with an N50 of 66.9 Mb and a L90 of 17 scaffolds. The longest HiC scaffold was 180.8 Mb.

After polishing the HiC assembly, the bobwhite genome assembly Cv_LA_1.0 included 1,512 scaffolds having an N50 of 66.8 Mb and a L90 of 17. Comparison of Cv_LA_1.0 with the Cv_TX_2.0 assembly (Table 1) shows the increase in contiguity of our assembly relative to the assembly produced by Oldeschulte *et al.* (2017). BUSCO analyses of both genomes are similar (Table 2), although we detected slightly fewer BUSCOs (-0.7%) in our Cv_LA_1.0 assembly relative to Cv_TX_2.0, perhaps due to repeat regions that were excluded from the contigs assembled by Meraculous. Future improvements to this assembly will incorporate Pacific Biosciences long-read sequences to help fill gaps that are likely associated with repeat regions that were difficult to assemble using short-reads.

ACKNOWLEDGMENTS

We thank Shaune Hall and other staff members at Dovetail Genomics for working with us, and Donna Dittmann and Steve Cardiff for assistance with specimen preparation and tissue storage. Special thanks to Chick and Lulu for assistance in the field. Funding for this project was provided by Louisiana State University, National Science Foundation grants DEB-1242260 and IOS-1754417 (to BCF) and DEB-1655624 (to BCF and RTB). JFS was supported by the LSU Museum of Natural Science, and portions of this research were conducted with high performance computational resources provided by the Louisiana Optical Network Infrastructure (<http://www.loni.org>). The individual bobwhite used in this study was collected by a private quail hunter with a valid Louisiana hunting license. NJS, WFH, DS, CC, JFS, and OJ performed field work; JFS and OJ prepared specimens; BCF performed analyses; JFS and BCF wrote the paper; BCF and RTB provided funding; and all authors reviewed and approved the manuscript.

LITERATURE CITED

- Bent, A. C., 1963 *Life Histories of North American Gallinaceous Birds*, Dover Publications, Inc., New York.
- Bolger, A. M., M. Lohse, and B. Usadel, 2014 Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30: 2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>
- Brennan, L. A., 1999 Northern bobwhite (*Colinus virginianus*). The birds of North America, number 397. National Academy of Natural Sciences, Philadelphia, Pennsylvania, and The American Ornithologists' Union, Washington, D.C.
- Burger, L. W., D. A. Miller, and R. I. Southwick, 1999 Economic impact of northern bobwhite hunting in the southeastern United States. *Wildl. Soc. Bull.* 27: 1010–1018.

- Chapman, J. A., I. Ho, S. Sunkara, S. Luo, G. P. Schroth *et al.*, 2011 Meraculous: de novo genome assembly with short paired-end reads. *PLoS One* 6: e23501. <https://doi.org/10.1371/journal.pone.0023501>
- Cole, L. J., H. L. Stoddard, and E. V. Komarek, 1949 Red bob-white: A report and correction. *Auk* 66: 28–35. <https://doi.org/10.2307/4080656>
- Dickinson, E. C., and J. V. Remsen, Jr. (Editors), 2013 *The Howard & Moore Complete Checklist of the Birds of the World*, Aves Press, Eastbourne, U.K.
- Domyan, E. T., M. W. Guernsey, Z. Kronenberg, S. Krishnan, R. E. Boissy *et al.*, 2014 Epistatic and combinatorial effects of pigimentary gene mutations in the domestic pigeon. *Curr. Biol.* 24: 459–464. <https://doi.org/10.1016/j.cub.2014.01.020>
- Ellegren, H., 2014 Genome sequencing and population genomics in non-model organisms. *Trends Ecol. Evol.* 29: 51–63. <https://doi.org/10.1016/j.tree.2013.09.008>
- Ellsworth, D. L., J. L. Roseberry, and W. D. Klimstra, 1989 Genetic structure and gene flow in the northern bobwhite. *Auk* 106: 492–495.
- Eo, S. H., J. P. Wares, and J. P. Carroll, 2009 Subspecies and units for conservation and management of the northern bobwhite in the eastern United States. *Conserv. Genet.* 11: 867–875. <https://doi.org/10.1007/s10592-009-9926-9>
- Evans, K. O., M. D. Smith, L. W. Burger, R. J. Chambers, A. E. Houston *et al.*, 2009 Release of pen-reared bobwhites: Potential consequences to the genetic integrity of resident wild populations National Quail Symposium Proceedings 6: 121–133.
- Guthery, F. S., 1997 A philosophy of habitat management for northern bobwhites. *J. Wildl. Manage.* 61: 291–301. <https://doi.org/10.2307/3802584>
- Halley, Y. A., S. E. Dowd, J. E. Decker, P. M. Seabury, E. Bhattarai *et al.*, 2014 A draft de novo genome assembly for the northern bobwhite (*Colinus virginianus*) reveals evidence for a rapid decline in effective population size beginning in the Late Pleistocene. *PLoS One* 9: e90240. <https://doi.org/10.1371/journal.pone.0090240>
- Kent, W. J., C. W. Sugnet, T. S. Furey, K. M. Roskin, T. H. Pringle *et al.*, 2002 The human genome browser at UCSC. *Genome Res.* 12: 996–1006. <https://doi.org/10.1101/gr.229102>
- Lamichhaney, S., G. Fan, F. Widemo, U. Gunnarsson, D. S. Thalmann *et al.*, 2016 Structural genomic changes underlie alternative reproductive strategies in the ruff (*Philomachus pugnax*). *Nat. Genet.* 48: 84–88. <https://doi.org/10.1038/ng.3430>
- Li, H., and R. Durbin, 2009 Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25: 1754–1760. <https://doi.org/10.1093/bioinformatics/btp324>
- Lieberman-Aiden, E., N. L. van Berkum, L. Williams, M. Imakaev, T. Ragoczy *et al.*, 2009 Comprehensive mapping of long-range interactions reveals folding principles of the human genome. *Science* 326: 289–293. <https://doi.org/10.1126/science.1181369>
- Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan *et al.*, 2009 The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25: 2078–2079. <https://doi.org/10.1093/bioinformatics/btp352>
- Marçais, G., and C. Kingsford, 2011 A fast, lock-free approach for efficient parallel counting of occurrences of k-mers. *Bioinformatics* 27: 764–770. <https://doi.org/10.1093/bioinformatics/btr011>
- Mikheenko, A., A. Prjibelski, V. Saveliev, D. Antipov, and A. Gurevich, 2018 Versatile genome assembly evaluation with QUAST-LG. *Bioinformatics* 34: i142–i150. <https://doi.org/10.1093/bioinformatics/bty266>
- Oldeschulte, D. L., Y. A. Halley, M. L. Wilson, E. K. Bhattarai, W. Brashear *et al.*, 2017 Annotated draft genome assemblies for the northern bobwhite (*Colinus virginianus*) and the scaled quail (*Callipepla squamata*) reveal disparate estimates of modern genome diversity and historic effective population size. *G3 (Bethesda)* 7: 3047–3058. <https://doi.org/10.1534/g3.117.043083>
- Putnam, N. H., B. L. O’Connell, J. C. Stites, B. J. Rice, M. Blanchette *et al.*, 2016 Chromosome-scale shotgun assembly using an in vitro method for long-range linkage. *Genome Res.* 26: 342–350. <https://doi.org/10.1101/gr.193474.115>
- Tuttle, E. M., A. O. Bergland, M. L. Korody, M. S. Brewer, D. J. Newhouse *et al.*, 2016 Divergence and functional degradation of a sex chromosome-like supergene. *Curr. Biol.* 26: 344–350. <https://doi.org/10.1016/j.cub.2015.11.069>
- Vijay, N., C. M. Bossu, J. W. Poelstra, M. H. Weissensteiner, A. Suh *et al.*, 2016 Evolution of heterogeneous genome differentiation across multiple contact zones in a crow species complex. *Nat. Commun.* 7: 13195. <https://doi.org/10.1038/ncomms13195>
- Vurture, G. W., F. J. Sedlazeck, M. Nattestad, C. J. Underwood, H. Fang *et al.*, 2017 GenomeScope: fast reference-free genome profiling from short reads. *Bioinformatics* 33: 2202–2204. <https://doi.org/10.1093/bioinformatics/btx153>
- Walker, B. J., T. Abeel, T. Shea, M. Priest, A. Abuouliel *et al.*, 2014 Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS One* 9: e112963. <https://doi.org/10.1371/journal.pone.0112963>
- Waterhouse, R. M., M. Seppey, F. A. Simão, M. Manni, P. Ioannidis *et al.*, 2018 BUSCO applications from quality assessments to gene prediction and phylogenomics. *Mol. Biol. Evol.* 35: 543–548. <https://doi.org/10.1093/molbev/msx319>
- Wellenreuther, M., and L. Bernatchez, 2018 Eco-evolutionary genomics of chromosomal inversions. *Trends Ecol. Evol.* 33: 427–440. <https://doi.org/10.1016/j.tree.2018.04.002>
- Williford, D. L., 2013 Molecular genetics of the northern bobwhite, scaled quail, and Gambel’s quail. Ph.D. Dissertation, Texas A&M University-Kingsville, Kingsville.
- Williford, D., R. W. Deyoung, R. L. Honeycutt, L. A. Brennan, and F. Hernández, 2016 Phylogeography of the bobwhite (*Colinus*) quails. *Wildl. Monogr.* 193: 1–49. <https://doi.org/10.1002/wmon.1017>
- Williford, D., R. W. Deyoung, R. L. Honeycutt, L. A. Brennan, H. F. Wehland *et al.*, 2014 Contemporary genetic structure of the northern bobwhite west of the Mississippi River. *J. Wildl. Manage.* 78: 914–929. <https://doi.org/10.1002/jwmg.733>

Communicating editor: A. Sethuraman