

2-1-2017

## Biogeography of a plant invasion: Genetic variation and plasticity in latitudinal clines for traits related to herbivory

Ganesh P. Bhattarai  
*Louisiana State University*

Laura A. Meyerson  
*University of Rhode Island*

Jack Anderson  
*University of Rhode Island*

David Cummings  
*University of Rhode Island*

Warwick J. Allen  
*Louisiana State University*

*See next page for additional authors*

Follow this and additional works at: [https://repository.lsu.edu/biosci\\_pubs](https://repository.lsu.edu/biosci_pubs)

---

### Recommended Citation

Bhattarai, G., Meyerson, L., Anderson, J., Cummings, D., Allen, W., & Cronin, J. (2017). Biogeography of a plant invasion: Genetic variation and plasticity in latitudinal clines for traits related to herbivory. *Ecological Monographs*, 87 (1), 57-75. <https://doi.org/10.1002/ecm.1233>

This Article is brought to you for free and open access by the Department of Biological Sciences at LSU Scholarly Repository. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Scholarly Repository. For more information, please contact [ir@lsu.edu](mailto:ir@lsu.edu).

---

**Authors**

Ganesh P. Bhattarai, Laura A. Meyerson, Jack Anderson, David Cummings, Warwick J. Allen, and James T. Cronin

1  
2  
3  
4  
5  
6  
7

Received Date : 26-Apr-2016

Revised Date : 25-Aug-2016

Accepted Date : 30-Aug-2016

Article type : Article

Running Head: Latitudinal clines in herbivory

**Biogeography of a plant invasion: genetic variation and plasticity in latitudinal clines for traits related to herbivory**

Ganesh P. Bhattarai<sup>1,3,5</sup>, Laura A. Meyerson<sup>2</sup>, Jack Anderson<sup>2</sup>, David Cummings<sup>2</sup>, Warwick J. Allen<sup>1,4</sup>, and James T. Cronin<sup>1</sup>

<sup>1</sup>*Department of Biological Sciences, Louisiana State University, Baton Rouge, LA 70803  
USA*

<sup>2</sup>*Department of Natural Resource Sciences, University of Rhode Island, 1 Greenhouse Road,  
Kingston, RI 02881 USA*

<sup>3</sup>*Present address: Department of Entomology, Kansas State University, Manhattan, KS  
66506 USA*

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/ecm.1233](https://doi.org/10.1002/ecm.1233)

This article is protected by copyright. All rights reserved

<sup>4</sup>*Present address: The Bio-Protection Research Centre, Lincoln University, PO Box 84,  
Lincoln 7647, New Zealand*

<sup>5</sup>Corresponding Author. E-mail: [bhattaraigp@gmail.com](mailto:bhattaraigp@gmail.com)

Author Manuscript

8 *Abstract:* The juxtaposition of plant-species invasions with latitudinal gradients in herbivore  
9 pressure is an important yet mostly unexplored issue in invasion biology. Latitudinal clines in  
10 defense and palatability to herbivores are expected to exist in native plant species but the  
11 evolution of these clines may lag behind for invasive plant species resulting in non-parallel  
12 latitudinal clines that may impact invasion success. Our study focused on a native and European  
13 invasive lineages of the common reed *Phragmites australis* in North America. Using native and  
14 invasive genotypes of *P. australis* collected across a 17° latitudinal range, we performed  
15 experiments in replicate northern and southern common gardens to investigate whether these two  
16 lineages exhibited different genetically based latitudinal clines in defenses, nutritional condition  
17 and palatability to their herbivores, the aphid *Hyalopterus pruni* and the fall armyworm  
18 *Spodoptera frugiperda*. We also tested whether invasive genotypes are more phenotypically  
19 plastic than native genotypes and whether plasticity varies with latitude. Although invasive  
20 genotypes did not exhibit higher defense levels (leaf toughness, phenolics, % carbon), they were  
21 considerably less palatable to their herbivores than native genotypes. Genetic-based latitudinal  
22 clines were evident for both native and invasive *P. australis* and for all defenses, nutrients and at  
23 least one palatability trait for each herbivore. In 36% of the cases where clines were evident, they  
24 were non-parallel between the two lineages. These data suggest that clines in the invasive  
25 genotypes of *P. australis* evolved within the past ~100 years. Moreover, our study showed that  
26 the occurrence and direction of latitudinal clines in plant traits were commonly dependent on  
27 where the study was conducted (north or south), indicating strong phenotypic plasticity in these  
28 genetic-based clines. Finally, traits for invasive genotypes of *P. australis* were 2.5 times more  
29 plastic than traits for native genotypes. Interestingly, plasticity for native but not invasive  
30 genotypes was strongly dependent on latitude of origin. Such spatial heterogeneity within and  
31 between the native and invasive lineages of *P. australis* with respect to their interactions with  
32 herbivores can generate substantial spatial variability in biotic resistance that can have important  
33 implications for the establishment and spread of invasive genotypes and species.

34  
35 *Keywords:* *biotic resistance, enemy-release hypothesis, Hyalopterus pruni, invasive species,*  
36 *latitudinal gradients, phenotypic plasticity, Phragmites australis, plant-insect interactions, plant*  
37 *defense, Spodoptera frugiperda.*

38

## INTRODUCTION

One of the most well-supported biogeographical patterns in ecology is the increase in primary productivity and species richness as latitude decreases (Rosenzweig 1995, Hillebrand 2004). Over the past several decades, much interest also has focused on latitudinal gradients in consumer-resource interactions (e.g., Coley and Aide 1991, Pennings et al. 2001, 2009, Schemske et al. 2009, Kim 2014, Cronin et al. 2015). For plant-herbivore interactions, herbivore damage is expected to increase toward lower latitudes because of a longer growing season and more benign winter conditions (e.g., Dobzhansky 1950, Coley and Aide 1991, Bolser and Hay 1996, Pennings et al. 2001, 2009, Kozlov et al. 2015, but see Andrew and Hughes 2005, Moles et al. 2011). In response, natural selection should favor increased defenses or reduced palatability in plant species at lower than higher latitudes (Coley and Aide 1991, Schemske et al. 2009).

Although this ecological paradigm is deemed too simplistic because it ignores feedbacks between plant defenses and herbivore abundance, and indirect and multi-trophic interactions (Kim 2014), latitudinal gradients in herbivory and defense are commonly reported. In the meta-analysis by Moles et al. (2011), 37% of the studies showed the expected negative relationship between latitude and herbivory. An additional 21% of the studies exhibited a positive relationship. Fifty-one percent of the studies exhibited a latitudinal gradient (positive and negative) in defense levels. Even though support for this paradigm is mixed, latitudinal gradients in herbivory, defense and other traits related to plant-herbivore interactions are a common occurrence.

The invasion and subsequent spread of a non-native species across a broad geographic range is expected to be followed by evolutionary changes in response to novel environmental and biotic gradients. A number of studies have documented that invasive species have rapidly evolved (< 100 years) in response to an environmental gradient in their introduced range. For example, invasive species have evolved distinct clines in growth and fitness-related traits with latitude (Maron et al. 2004, Novy et al. 2013, Li et al. 2015) that parallel the clines for the same species in their native range. Although the evolution of latitudinal clines in plant defenses or palatability to herbivores is expected to occur with invasive species, no study has ever examined whether or how quickly latitudinal clines have formed in traits related to plant-herbivore interactions.

69           The evolution of latitudinal clines in defenses and/or palatability traits in an invasive  
70 species may have important implications for invasion success (Bezemer et al. 2014, Cronin et al.  
71 2015). For example, the biogeographic paradigm described previously predicts that native plants  
72 should exhibit latitudinal clines in defense and palatability traits (see e.g., Pennings et al. 2001).  
73 However, an invasive species may not have had sufficient time to evolve a gradient that parallels  
74 the gradients for co-occurring native species. These non-parallel gradients in defense or  
75 palatability between native and invasive species, particularly early in the invasion process, may  
76 create large-scale heterogeneity in the relative impact of herbivores on co-occurring native and  
77 invasive plant taxa. As such, in some regions, herbivory may be greater on invasive than native  
78 plants (supporting the biotic resistance hypothesis; Levine et al. 2004, Chun et al. 2010) and in  
79 other regions, the reverse may occur (i.e., biotic susceptibility). In the only example on this  
80 subject, Cronin et al. (2015) found latitudinal gradients in herbivore pressure for native  
81 genotypes of *Phragmites australis* (Poaceae) in the field in North America. Invasive genotypes  
82 that are sympatric with native genotypes exhibited no latitudinal gradients in herbivore pressure.  
83 These non-parallel gradients in herbivory between native and invasive *P. australis* resulted in  
84 greater herbivore pressure on native than invasive genotypes in the south (supporting the local  
85 enemy-release hypothesis; Zheng et al. 2012) but no difference in herbivore pressure in the  
86 north. Importantly, the field study by Cronin et al. (2015) did not allow us to assess whether the  
87 geographic variation in *P. australis*-herbivore traits was genetically based, the result of plastic  
88 responses by the plants to an environmental gradient (e.g., climate), or some combination of the  
89 two.

90           Phenotypic plasticity, variability in the expression of traits in different environments, is  
91 another possible mechanism that can generate spatial heterogeneity in native-invasive plant  
92 responses to herbivores. Phenotypic plasticity is thought to be beneficial to invaders because  
93 plastic “general-purpose genotypes” could have a fitness advantage in founder populations (e.g.,  
94 Richards et al. 2006, Chun 2011, Davidson et al. 2011). Alternatively, plasticity may also evolve  
95 in an invasive species in response to the novel environment after colonization (Richards et al.  
96 2006, Lavergne and Molofsky 2007). In their meta-analysis, Davidson et al. (2011) found strong  
97 support for the prediction that invasive species are more phenotypically plastic than native  
98 species. Moreover, theoretical models suggest that trait plasticity may vary along a climatic  
99 gradient, being greater at range margins than range interiors (Chevin and Lande 2011). A number

100 of studies have demonstrated latitudinal gradients in trait plasticity (e.g., Maron et al. 2004, De  
101 Frenne et al. 2011, Woods et al. 2012). Consequently, we may expect co-occurring native and  
102 invasive species to differ in trait plasticity along a latitudinal gradient which can contribute to  
103 spatial heterogeneity in invasion success. To date, no study has addressed this possibility.

104 Although the common reed, *Phragmites australis*, is native and widespread in North  
105 America, a continent wide invasion by introduced European genotypes of *P. australis* in North  
106 America has been underway for at least 150 years (Saltonstall 2002). We conducted experiments  
107 in replicate common gardens, one in the north (41.49°, University of Rhode Island) and one in  
108 the south (30.35°, Louisiana State University), to assess whether genetic-based latitudinal  
109 gradients exist for different traits associated with the interactions between native and invasive *P.*  
110 *australis* and its herbivores. Because the study was conducted in a common-garden environment  
111 and maternal effects were minimized, any latitudinal patterns found would be genetically based.  
112 Using multiple native and invasive genotypes of *P. australis* collected across a 17° latitudinal  
113 range in North America (Fig. 1), we quantified plant defense levels (leaf toughness, total  
114 phenolics) and nutritional condition (percent carbon, percent nitrogen, CN ratio, water content),  
115 and conducted experiments to assess palatability to two common and widespread generalist  
116 herbivores, the mealy-plum aphid, *Hyalopterus pruni* (Homoptera: Aphididae), and the fall  
117 armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae). We tested the following six  
118 hypotheses. (1) Genetic-based latitudinal clines for plant defenses and palatability to herbivores  
119 are evident for native *P. australis* genotypes. (2) Because European invasive genotypes may not  
120 have had sufficient time to evolve in response to their novel environment, latitudinal clines in  
121 defense and palatability traits will be absent or weak, and thus non-parallel to the gradients for  
122 the native genotypes. Based on differences between replicate common gardens, we also tested  
123 the hypotheses that (3) invasive genotypes are more phenotypically plastic than native  
124 genotypes, (4) plasticity varies with latitude, and (5) latitudinal clines for each lineage differ  
125 between gardens, indicating that these clines are phenotypically plastic. Finally, in light of our  
126 findings with regard to the previous hypotheses, we test the hypothesis that (6) across a broad  
127 latitudinal range in North America, European genotypes of *P. australis* are successful invaders  
128 because they are better defended and less palatable to herbivores than native genotypes. Support  
129 for this latter hypothesis would suggest that native communities have low biotic resistance to  
130 invasion by European *P. australis*.



131

132

## METHODS

### 133 Study system

134 *Phragmites australis* is considered a model organism for the study of plant invasions (for  
135 a detailed discussion, see Meyerson et al. 2016). *Phragmites australis* is a tall and robust  
136 perennial grass of coastal and freshwater marshes with a nearly worldwide distribution (Marks et  
137 al. 1994, Lambertini et al. 2006). It was an uncommon species of wetland communities in North  
138 America for millennia but exhibited dramatic and rapid spread over the past ~150 years,  
139 particularly in the mid-Atlantic region of North America (Chambers et al. 1999). The rapid  
140 spread is attributed to the introduction of an invasive Eurasian genotype in mid 1800s (Chambers  
141 et al. 1999, Saltonstall 2002). Additional introduced genotypes have been discovered from the  
142 Atlantic and Gulf Coast regions of North America (Lambertini et al. 2012, Meyerson and Cronin  
143 2013). At least 14 genotypes of native *P. australis* lineage are distributed throughout North  
144 America (Saltonstall 2002, Meadows and Saltonstall 2007) and have been reported to be  
145 threatened by the spread of invasive *P. australis* genotypes (Meyerson et al. 2010). Phylogenetic  
146 analysis revealed that native and introduced genotypes belong to two different clades (hereafter,  
147 lineages) often regarded as separate subspecies (Saltonstall 2002). Hybridization is shown to  
148 occur between native and invasive genotypes in nature (Saltonstall 2003, Paul et al. 2010,  
149 Saltonstall et al. 2014).

150 Among the most common and important herbivores of *P. australis* in North America are  
151 several introduced species, including the mealy plum aphid *H. pruni* and several species of gall  
152 flies in the genus *Lipara* (Diptera: Chloropidae) (Tewksbury et al. 2002, Lambert et al. 2007,  
153 Allen et al. 2015, Cronin et al. 2015). Native *P. australis* genotypes suffer substantially greater  
154 herbivore damage than the invasive genotypes (Lambert and Casagrande 2007, Allen et al. 2015,  
155 Cronin et al. 2015). Moreover, native genotypes but not invasive genotypes exhibit latitudinal  
156 gradients in herbivore damage from the three major feeding guilds (sucking, stem-feeding  
157 [galling], and chewing) along the Atlantic and Gulf coasts of North America (Cronin et al. 2015).  
158 Leaf tissue loss from chewing herbivores and incidence of stem galls (primarily *Lipara* spp.)  
159 decreased with increasing latitude whereas densities of *H. pruni* increased with increasing  
160 latitude. These results suggest that native *P. australis* genotypes are more likely to have evolved  
161 latitudinal clines in response to herbivore pressure.

162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192

### **Plant defense, nutritional and palatability traits in common gardens**

We established replicate common gardens at Louisiana State University, Baton Rouge, LA (LSU: 30.35°, -91.14°) and the University of Rhode Island, Kingston, RI (URI: 41.49°, -71.54°, Fig. 1). A small clump of rhizome was collected from 12 native and 16 invasive patches from the Gulf and Atlantic Coasts of North America (Fig. 1, Appendix S1). The rhizome material was split between the two gardens and propagated. Henceforth, we refer to each rhizome source as a source genotype. Owing to sprouting failures, mortality during propagation, and slow growth of some genotypes, the number of genotypes and their identity did not perfectly match between gardens at the time of our experimental studies. Seven native and 13 invasive genotypes survived at the LSU garden whereas 10 native and 15 invasive genotypes survived at URI. Four native and eight invasive genotypes were common in both gardens in 2012 when the majority of the traits (defense, nutritional traits and palatability to *H. pruni*; see below) were quantified (Appendix S1). The URI garden represents a moderate temperate-zone climate, is roughly at the midpoint of the distribution of native genotypes used in our gardens, and is not far from where invasive *P. australis* first became established in North America (Saltonstall 2002). In contrast, the LSU garden is subtropical, close to the southern range limit of invasive genotypes, and is ~700 km south of our southernmost native population.

We used identical methods for the propagation of plants in each garden, including soil type and watering, fertilization and insecticidal treatment regimes. Rhizomes were planted in Metromix® soil in 19 liter nursery pots and maintained in outdoor plastic pools filled with fresh water. Plants were propagated vegetatively to get at least 32 pots per genotype. Therefore, all plants in both gardens that belong to a source genotype were clones. Source genotypes were randomly distributed within each garden. By growing the plants in the gardens for at least one year prior to the start of our study, maternal effects that might confound differences in plant traits were minimized. Plants were fertilized with Mega Green organic fertilizer upon detection of leaf yellowing and sprayed regularly with Safer® insecticidal soap (Woodstream Corp., Lititz, PA) to protect them from unwanted herbivores. Safer soap was used because it has a very short (< 2 week) residual time on the plants. *Hyalopterus pruni* was the most common pest in the garden and the insecticidal treatment was effective in keeping them at low abundances leading up to the start of the experiments. All palatability experiments described below were conducted with stems

193 that had no visible herbivore damage or traces. We therefore expected little to no effects of  
194 background herbivores on plant growth and fitness traits.

195 Because we prevented seed production by clipping and removing panicles before seeds  
196 were produced and only rhizomatous growth occurred in our gardens, it was not possible for the  
197 clonal populations to evolve in response to the local climate. Consequently, differences in mean  
198 trait values between plants (clones) from the same source genotype, but from gardens separated  
199 by 11° latitude, should be the result of phenotypic plasticity in that trait (Maron et al. 2004,  
200 Colautti et al. 2009, Woods et al. 2012).

201 Caging experiments were performed to assess the palatability of *P. australis* genotypes to  
202 herbivores from two feeding guilds: the mealy-plum aphid *H. pruni* and the fall armyworm *S.*  
203 *frugiperda*. *Hyalopterus pruni* is an invasive pest of dried plum (*Prunus domestica*) that was  
204 introduced from Europe and uses *P. australis* as a secondary host plant (Lozier et al. 2009). It is  
205 one of the most widespread herbivores of *P. australis* throughout North America and Europe and  
206 can produce massive outbreaks with severe damage to *P. australis* stands (Cronin et al. 2015).  
207 First recorded in California in 1881 (Smith 1936) it was probably introduced to North America  
208 after the introduction of invasive *P. australis* (Lozier et al. 2009). Consequently, the native and  
209 invasive lineages of *P. australis* in North America most likely had an equivalent amount of time  
210 to adapt to aphid herbivory and evolve clines. Therefore, we may expect parallel latitudinal  
211 gradients for traits associated with *H. pruni* herbivory between native and invasive *P. australis*.

212 *Spodoptera frugiperda* is native to North America and is a serious pest of cereal crops  
213 (Sparks 1979). It is known to feed on many grass species including *P. australis* (Sparks 1979,  
214 Bhattarai 2015). It overwinters only in the mild climates of southern US and reinvades most of  
215 the US and southern Canada every year (Sparks 1979). The annual northward invasion by this  
216 species is likely to result into a latitudinal gradient in duration and intensity of herbivory on the  
217 host plants. Native genotypes of *P. australis*, which have had a potentially long history of  
218 interaction with these herbivores, are expected to exhibit latitudinal gradients for traits associated  
219 with the *P. australis* - *S. frugiperda* interaction. In contrast, such gradients may not exist for the  
220 invasive genotypes due to their relatively short history of interaction.

221 Because the phenological state of *P. australis* is likely to affect plant physiology,  
222 nutritional condition, defense levels, and herbivory (e.g., Liu et al. 2011, Lehndal and Agren  
223 2015), our palatability experiments were designed to minimize differences in plant phenology

224 between gardens and among source genotypes within gardens. However, phenology is a difficult  
225 concept to apply to grasses in the middle of their growing season because they are constantly  
226 producing new shoots. For example, within a pot of *P. australis* during late spring or summer,  
227 there is considerable variability in plant phenological state. Consequently, metrics such as  
228 growing degree days (GDD; McMaster and Wilhelm 1997) have limited value in these cases.  
229 Therefore, to minimize variation among experimental plants, we did the following. First, the  
230 experiments were timed to coincide with when the aphid colonies were well established and *S.*  
231 *frugiperda* caterpillars were present in the region. Second, we chose mid-size (growing) stems  
232 that were between 0.75 and 1.25 m in height. Finally, within a stem, we chose the uppermost,  
233 fully open leaf for *H. pruni* or the upper 5-10 leaves for *S. frugiperda* cages.

234 *Palatability to H. pruni.* To assess the palatability of *P. australis* genotypes to aphids, we  
235 caged aphids on plants and measured colony growth. This bioassay (and other palatability  
236 metrics described below) integrates the effects of multiple plant defensive and nutritional traits  
237 on the herbivore's survival and reproduction (i.e., fitness) and is therefore a more holistic  
238 measure of plant resistance (Kim 2014). The source of aphids was a naturally occurring stand of  
239 *P. australis* within 80 km of each common garden. For ethical reasons and to minimize the  
240 genetic variation among aphids within each garden, we used a single source population of aphids  
241 for each garden. Cages for the aphid experiment were constructed using five cm lengths of clear  
242 acrylic tubing (2.8 cm in diameter, 0.1 cm in thickness) that were inserted over the *P. australis*  
243 leaves. The ends of the tubes were sealed with closed-cell foam plugs that had a small fine-  
244 screen window cut into their centers for air circulation (see Appendix S2: Fig. S1). Aphid  
245 colonies were initiated with two adult aphids caged on the youngest fully open leaf on a  
246 randomly selected stem from each pot. Aphids reproduce parthenogenetically and produce a  
247 colony within a few days. After 10 d, leaves with aphid colonies were collected, transported on  
248 ice to the laboratory, and stored in a freezer at -20° C. With a suitable host, aphid colonies can  
249 increase in size by 100-fold in 10 d without any evidence of leaf deterioration or resource  
250 depletion due to intraspecific competition (G.P. Bhattarai personal observation). Aphids per  
251 colony were enumerated and then dried at 40°C for two days. Dry mass of each colony was  
252 determined using a Mettler microbalance (0.1 mg precision). Because aphid mass was strongly  
253 correlated with aphid colony size ( $R = 0.878$ ,  $P < 0.0001$ ) we used only colony size in  
254 subsequent analyses. Aphid colony survival was determined as the proportion of cages per *P.*

255 *australis* source genotype that had a viable aphid colony after 10 d. Survivorship and colony size  
256 were used as indicators of plant palatability to the aphids. The aphid experiment was conducted  
257 at LSU from April 13-23, 2012 with nine introduced and five native genotypes (20 cages per  
258 genotype), and repeated at URI from June 17-27, 2012 with 12 introduced and six native  
259 genotypes (8-12 cages per genotype). The experiments were performed two months later in the  
260 URI than LSU garden so that plants were in comparable developmental stages (about two  
261 months old) in each garden.

262 *Palatability to S. frugiperda.* For *S. frugiperda*, we also performed a caging experiment  
263 and measured the performance of individual larvae feeding on plants from each source genotype.  
264 For both gardens, *S. frugiperda* were obtained as eggs from the same source (Benzon Research  
265 Inc., Carlisle, PA) and larvae were reared in the laboratory on artificial diet provided by  
266 Southland Products (Lake Village, AR). At the fourth instar stage, the mass of each larva was  
267 determined, and those larvae within 20-50 mg were selected for the experiment. Sleeve cages  
268 (45×60 cm<sup>2</sup> or 60×75cm<sup>2</sup> in size) were built using fine insect netting. The cage was inserted over  
269 the upper portion of a stem (containing 5-10 leaves) and enclosed around the stem at the bottom  
270 using a cable tie (see Appendix S2: Fig. S2). A single caterpillar was released into each cage  
271 through a hole cut open in the top of the cage. The hole was subsequently stapled closed. Within  
272 each pot, a single stem was selected at random for a cage. The experiment was terminated after 8  
273 d, before any caged plants had all available leaf material consumed by the *S. frugiperda* and  
274 before the larva could mature to the pupal stage. Each larva was collected, transported on ice to  
275 the laboratory and its fresh mass was determined using a Mettler microbalance. Larval growth  
276 was calculated as proportional change in fresh biomass during the experiment ( $\ln$  final mass [mg]  
277 -  $\ln$  initial mass [mg]). Survivorship of larvae on each source genotype of *P. australis* was  
278 determined as the proportion of cages with a live larva at the end of the experiment. All the cases  
279 in which the larva died before the termination of the trial were excluded from the analyses for  
280 palatability traits related to *S. frugiperda* (see below).

281 We took photographs of all the remaining leaves inside the cage to quantify leaf area  
282 consumed by each larva. Using ImageJ (Rasband 2014), we quantified the remaining leaf area  
283 (cm<sup>2</sup>) for each plant and estimated pre-consumption leaf area by extrapolation. Leaf area  
284 consumed by each larva was estimated as the pre-consumption leaf area – leaf area remaining  
285 after 8 d. We used leaf area consumed and not proportion leaf area consumed because the total

286 leaf area within a cage was variable and abundant (never being completely consumed), and  
287 therefore, the former measure is more informative. The amount of leaf area consumed by larvae  
288 could be an indicator of plant defense levels (e.g., Coley 1986). However, herbivores may also  
289 consume more to compensate for the lower quality of leaf tissues (Mattson 1980). Finally, we  
290 determined biomass conversion efficiency of larva (larval growth per unit [ $\ln$ ] area of leaf  
291 consumed). Leaf area measurements for plants with dead or missing larvae were excluded from  
292 the analyses. These four variables, larval survivorship and growth, leaf area consumed, and  
293 biomass conversion efficiency, were used as the measurements of plant palatability to chewing  
294 herbivores. The experiment was performed on May 23-31, 2013 including nine native and 13  
295 invasive genotypes (20 plants per genotype) at LSU garden, and on August 7-15, 2013 including  
296 six native and eight invasive genotypes (7-10 plants per genotype) at URI.

297 *Plant defense and nutritional traits.* Plant characteristics related to defense and nutritional  
298 quality were measured concurrently with the *H. pruni* experiment and from the same pots as  
299 those with cages. Water content of leaves has been shown to have a positive relationship with  
300 population growth rate of aphids (e.g., Johnson 2008, but see Woods et al. 2012) and  
301 lepidopteran larvae (Scriber and Feeny 1979). Water content of leaves was estimated as the  
302 proportion of water per unit fresh biomass of three newly opened leaves collected from each pot  
303 ( $n = 10$  per genotype). For nutrient analysis (percent carbon, percent nitrogen and CN ratio), the  
304 top three leaves were collected from a single plant per pot ( $n = 5$  pots per genotype). Leaves  
305 were lyophilized and ground to a fine powder in the laboratory. Leaf nutritional levels were  
306 assayed using an elemental analyzer at Brown University Environmental Chemistry Facilities  
307 (<http://www.brown.edu/Research/Evchem/facilities/>). Herbivores often prefer and perform better  
308 on plants with higher % nitrogen (Mattson 1980, Agrawal 2004). Also, carbon content of leaves  
309 has been shown to influence herbivore performance (Agrawal 2004, Cronin et al. 2015).

310 Leaf toughness (force [kg] required to push a blunt steel rod [4.8 mm in diameter]  
311 through the leaf) and total phenolics concentration were our measures of plant defenses against  
312 herbivores. As a member of the Poaceae, *P. australis* defenses are likely limited to structural  
313 defenses and phenolics (McNaughton 1979, Tschardtke and Greiler 1995, Strauss and Agrawal  
314 1999). In our field surveys (Cronin et al. 2015), leaf toughness and phenolics were negatively  
315 related to leaf-area consumed by chewing herbivores and aphid density, respectively. Using a  
316 penetrometer (Itin Scale Co., Inc., Brooklyn, NY), toughness was measured for the fully open

317 uppermost leaf from a randomly selected stem per pot. Leaf toughness was also measured for the  
318 top-most leaf inside each cage of the *S. frugiperda* experiment in 2013. Total phenolics (nM/g of  
319 dried leaf tissue) were estimated using a modified version of the Folin-Ciocalteu method  
320 (Waterman and Mole 1994, Cronin et al. 2015).

321

322

### Statistical methods

323 Our primary objective with this study was to determine if *P. australis* trait variation  
324 within a common garden is genetically based, and particularly whether this genetic variation was  
325 attributed to differences between lineages (native, invasive) and/or was related to latitude of  
326 origin of the genotype (i.e., a genetic-based cline). A secondary objective involved using source  
327 genotypes that were shared between common gardens to assess whether traits associated with *P.*  
328 *australis* – herbivore interactions are more phenotypically plastic for invasive than native  
329 lineages. For the primary objective, we used mixed-effects or general linear models to test  
330 whether each plant defense, nutritional, and palatability trait varied between gardens, lineages  
331 and along a latitudinal gradient. Garden (LSU, URI), lineage (native, invasive), and latitude of  
332 origin (hereafter, latitude effect) were treated as fixed effects, and source genotype was a random  
333 effect. With uniquely coded source genotypes (see Appendix S1), that belong to either native or  
334 invasive lineage, included in the model as a random effect our model structure was equivalent to  
335 a nested-model (i.e., genotypes within a lineage).

336 Several traits required a slightly different analytical approach. Aphid colony size (number  
337 per cage) was Poisson distributed. Therefore, for this trait, we used a generalized mixed effect  
338 model with Laplace estimation method and Poisson distribution of errors (Bolker et al. 2009). To  
339 account for overdispersion of the count data, an observation-level random effect was also added  
340 to the model (Bolker et al. 2009). Because we had a single estimate of *H. pruni* and *S. frugiperda*  
341 larval survivorship per source genotype, we could not assess a source-genotype effect for these  
342 two traits. Therefore, survivorship of aphids and larvae were analyzed using general linear  
343 models. Finally, because we measured leaf toughness in 2012 and 2013, we included year as a  
344 random variable in the linear mixed-effects model for this defense trait.

345 To help normalize data distributions and homogenize variances among categories (i.e.,  
346 gardens, lineages), % carbon, % nitrogen, CN ratio, total phenolics and leaf area were *ln*-  
347 transformed. Quantile-quantile plots were also used to identify potential outliers in the

348 distribution of trait estimates. However, in no case did the removal of these data points  
349 qualitatively change the conclusions of the model.

350 For each dependent variable, we used Akaike's Information Criteria corrected for finite  
351 sample size (AICc) to select the most informative model (Burnham and Anderson 2010). The  
352 full model included all random and fixed effects plus all two- and three-way interactions. In  
353 addition, we included a quadratic term ( $\text{latitude}^2$ ) to evaluate whether the relationship between  
354 the trait and latitude was nonlinear. Candidate models were constructed using all possible  
355 combinations of the variables including one with no fixed effect, but with two restrictions. First,  
356 interaction terms could only be present in the model if their main effects were also present in the  
357 model. Second, the random effects (source genotype and the repeated measure of toughness)  
358 were retained in every model combination. Without this underlying structure to the model, the  
359 design would be pseudoreplicated.

360 Candidate models were ranked by AICc from lowest to highest value and AICs with a  $\Delta_i$   
361 value ( $= \text{AICc}_i - \text{AICc}_{\min}$ ) of  $\leq 2$  were deemed to have substantial support (Burnham and  
362 Anderson 2010). We also report the AICc weights ( $w_i$ ) which indicate the weight of evidence (as  
363 a proportion) in favor of model  $i$  being the best model given the set of candidate models.  
364 Goodness of fit of each mixed effects model was reported as marginal ( $R^2_m$ , variance explained  
365 by fixed effects) and conditional  $R^2$  ( $R^2_c$ , variance explained by the entire model) that are  
366 comparable in interpretation to the coefficient of determination,  $R^2$ , for linear models (Nakagawa  
367 and Schielzeth 2013). All analyses were run in R 3.2.0 (R Development Core Team 2015) using  
368 lme4 (Bates et al. 2014) and AICcmodavg (Mazerolle 2015) packages.

369 The AICc-best models for each trait were used to evaluate the hypotheses outlined in the  
370 Introduction. The presence of a latitude effect in the model would suggest that there is a genetic-  
371 based latitudinal cline in *P. australis* (Hypothesis 1). The addition of a lineage  $\times$  latitude  
372 interaction in the best model would further indicate the existence of non-parallel latitudinal  
373 gradients between the native and invasive lineage (Hypothesis 2). Otherwise, the absence of this  
374 interaction term would suggest parallel latitudinal gradients (or, more appropriately, no evidence  
375 that gradients differ between lineages). A garden effect in the best model would suggest  
376 phenotypic plasticity in the trait (Hypothesis 3) and a garden  $\times$  latitude interaction would support  
377 Hypothesis 4 that the degree of plasticity for a trait (i.e., the difference between gardens) varies  
378 with latitude of origin of the plants. A three-way interaction between fixed effects (Garden  $\times$



379 Lineage  $\times$  Latitude) would support our Hypothesis 5 that the slopes of the latitudinal clines for  
380 each lineage differs between gardens (i.e., the latitudinal gradient for each lineage is  
381 phenotypically plastic). Finally, if a lineage effect is present in the best model and plant defense  
382 trait values are higher or palatability is lower for the native than invasive lineage, it would  
383 support the biotic-resistance hypothesis (Hypothesis 6).

384 In cases where the AICc-best model included a garden interaction term (Garden  $\times$   
385 Lineage, Garden  $\times$  Latitude and/or Garden  $\times$  Lineage  $\times$  Latitude interactions), we repeated the  
386 model-selection procedure for each garden to better elucidate the lineage and latitude effects on  
387 each trait. In this case, candidate models included all combinations of lineage, latitude and the  
388 lineage  $\times$  latitude interaction, as well as a model with no fixed effects. If latitude (or latitude and  
389 lineage  $\times$  latitude interaction) was in the AICc-best model for a particular garden, we performed  
390 mixed effect analyses to determine the relationship between latitude and the trait for each  
391 lineage.

392 We performed a separate correlation analysis for each garden to examine whether plant  
393 palatability traits were linearly related to defense and nutritional traits. Plant-level measures of  
394 putative defense (leaf toughness, total phenolics), nutritional (water content, % nitrogen, %  
395 carbon, C:N ratio) and palatability (aphid colony size, larval growth, leaf area chewed and  
396 biomass conversion efficiency) were used in the pairwise correlation analysis (Pearson's product  
397 moment correlation,  $R$ ). Nutrient concentrations and leaf area chewed were  $\ln$  transformed.  $P$ -  
398 values were adjusted for multiple comparisons using a Bonferroni correction.

399 For the second objective, we quantified phenotypic plasticity in defense, nutritional and  
400 palatability traits for *P. australis* genotypes that were present in both common gardens (4 native,  
401 8 invasive; Appendix S1). We could not assess plasticity in palatability to aphids because the  
402 aphids were obtained from different source populations located nearby each garden. As such,  
403 differences in palatability to aphids between gardens could also be due to genetic differences  
404 between aphid source populations (e.g., local co-adaptation between southern aphids and  
405 southern populations of *P. australis*). Plasticity for each trait and lineage was measured as the  
406 proportional difference in mean trait expression between gardens; e.g., effect size = [mean leaf  
407 toughness for native genotypes at LSU – mean leaf toughness for native genotypes at URI]/mean  
408 leaf toughness for native genotypes at LSU. Lineage means per garden for each trait were  
409 obtained from the least-squares means from the linear mixed-effects model outlined above (or

410 the alternative models for larval survivorship, and leaf toughness) that contained garden and  
411 latitude as fixed effects and genotype as a random effect. For all traits, the effect size for native  
412 lineage ( $x$ -axis) was plotted against the effect size for invasive lineage ( $y$ -axis). If the data point  
413 for a trait falls above the 1:1 line, it would indicate that plasticity for that trait is greater for the  
414 invasive than native lineage. Data points below the line would indicate the opposite. A pairwise  
415  $t$ -test was performed to assess whether plasticity for the native and invasive lineages differed for  
416 all traits combined (Hypothesis 3). Finally, we also computed plasticity for each source genotype  
417 of *P. australis* and used ANCOVA to examine the effects of lineage and latitude on plasticity for  
418 each defense and nutritional trait (Hypothesis 4).

419

420

## RESULTS

### *Defense and nutritional traits*

422 The phenotypic expression of *P. australis* defense and nutritional traits was strongly  
423 influenced by plant lineage and latitude of origin, and also was strongly modulated by where the  
424 study was conducted (LSU or URI). The best model, based on AICc weights, included latitude  
425 for all six traits (leaf toughness, total phenolics, water content, % nitrogen, % carbon, and CN  
426 ratio) and lineage for four traits (total phenolics, water content, % nitrogen, and CN ratio)(Table  
427 1, Appendix S3). Interestingly, in all cases where lineage was present in the best model, so was a  
428 lineage  $\times$  latitude interaction; an indication that native and invasive lineages exhibit nonparallel  
429 genetically based gradients in these traits. Finally, not only was the garden where the study was  
430 conducted important in all cases (Table 1, see Appendix S3: Table S1 for detailed information),  
431 there were many interactions involving garden, lineage and latitude of origin.

432 The defensive trait leaf toughness was negatively related to latitude, decreasing by 51%  
433 between our southernmost and northernmost genotypes (Fig. 2A, Table 1). Plants from the LSU  
434 garden exhibited steeper declines in toughness with increasing latitude than plants from the URI  
435 garden (garden  $\times$  latitude interaction). Because of the garden  $\times$  latitude interaction in the best  
436 model, we conducted separate model-selection procedures for each garden to explore the effects  
437 of lineage, latitude and their interactions on leaf toughness. At the LSU garden, native and  
438 invasive lineages had parallel latitudinal clines in leaf toughness (no lineage  $\times$  latitude  
439 interaction in the AICc-best model; Appendix S3: Table S2). There was a significant negative  
440 relationship between toughness and latitude for the invasive genotypes but the relationship was

441 not significant for the native genotypes (Fig. 2A, Appendix S3: Tables S2 and S3). In contrast,  
442 total phenolics were 34% higher for native than invasive genotypes and 96% higher for plants  
443 grown in the LSU vs. URI garden (Fig. 2B, Appendix S3: Tables S1, S4 and S5). At the LSU  
444 garden, total phenolics for invasive and native lineages did not vary with latitude of origin  
445 (Appendix S3: Table S2). However, at URI, total phenolics increased with latitude for native  
446 genotypes but did not vary with latitude for invasive genotypes (lineage  $\times$  latitude and non-  
447 parallel clines; Appendix S3: Tables S2 and S3).

448 Similar to total phenolics, the AICc best models to explain water content, % nitrogen and  
449 the CN ratio were the same and included all main factors and their 2- and 3-way interactions  
450 (Table 1, Appendix S3). AICc weights for these models (the likelihood of the model given the  
451 candidate models considered) were  $> 0.93$ . Water content was 8% higher for native than invasive  
452 genotypes and 4% higher for plants grown at URI vs. LSU (Fig. 2C, Appendix S3: Tables S4 and  
453 S5). At LSU, water content decreased with increasing latitude for both lineages but the  
454 latitudinal clines were non-parallel (lineage  $\times$  latitude interaction in the AICc-best model;  
455 Appendix S3: Tables S2 and S3). At URI, the invasive genotypes exhibited a significant negative  
456 latitudinal cline in water content but no cline was evident for the native genotypes. The absence  
457 of a lineage  $\times$  latitude interaction in the AICc-best models suggest that these two clines are not  
458 different (Appendix S3: Table S2). Lineage, latitude and their interaction were important factors  
459 contributing to variation in % nitrogen at URI but were unimportant to % nitrogen at LSU (Fig.  
460 2D, Appendix S3: Table S2). At the URI garden, % nitrogen increased significantly with latitude  
461 for the native genotypes but declined (non-significant) with latitude for the invasive genotypes  
462 (non-parallel cline; Appendix S3: Table S3). Overall, % nitrogen was similar between the native  
463 and invasive genotypes ( $3.13 \pm 1.04\%$  and  $3.10 \pm 1.02\%$ , respectively; mean  $\pm$  SE). For CN ratio,  
464 the slopes of the latitudinal clines at URI were in the opposite direction as for % nitrogen and  
465 was only significant for the native genotypes (Fig. 2F, Appendix S3: Tables S2 and S3). The  
466 difference in CN ratio between lineages changed between gardens (garden  $\times$  lineage interaction;  
467 Fig. 2F): the CN ratio for invasive genotypes was 12.1% greater at LSU and 13.3% less at URI  
468 than for native genotypes. Finally, the AICc best model for % carbon included only garden,  
469 latitude and their interaction (Table 1, Appendix S3). Mean % carbon was similar between native  
470 and invasive lineages ( $45.46 \pm 1.01\%$  and  $45.25 \pm 1.00\%$ , respectively). On average, the %  
471 carbon declined by 1.7% from the south to the north and the relationship was only evident for the

472 plants at the LSU garden (hence the garden  $\times$  latitude interaction; Fig. 2E, Appendix S3: Table  
473 S2). Only for the invasive lineage at LSU was there a significant latitudinal cline (Appendix S3:  
474 Table S3).

475  
476 *Palatability to herbivores*

477 Traits associated with *P. australis* palatability to *H. pruni* and *S. frugiperda* were strongly  
478 influenced by the garden in which the study was conducted and plant lineage, and to a lesser  
479 extent latitude. For aphid colony size, the best model included lineage, latitude, garden and a  
480 garden  $\times$  lineage interaction (AICc weight = 1.0, Appendix S3: Table S1). Aphid colony size  
481 was strongly affected by lineage and latitude at both gardens (both lineages; parallel negative  
482 latitudinal gradients) (Fig. 3A, Appendix S3: Tables S2 and S3). The garden  $\times$  lineage  
483 interaction was present in the model because there was a 26-fold difference in colony size  
484 between native and invasive genotypes at LSU but only a 5-fold difference at URI (Fig. 3A).  
485 After accounting for the garden effect, aphid colony size averaged 11 times larger for native than  
486 invasive genotypes (Appendix S3: Table S4). Based on paired measurements obtained from *P.*  
487 *australis* stems, aphid colony size was positively correlated with water content in both gardens  
488 (LSU:  $R = 0.39$ ,  $P < 0.01$ ; URI:  $R = 0.23$ ,  $P = 0.04$ ; Appendix S4) and total phenolics in the LSU  
489 garden ( $R = 0.53$ ,  $P < 0.01$ ; Appendix S4). No other nutritional or defense trait was correlated  
490 with this palatability measure.

491 Similar fixed effects were included in the best model for aphid survivorship (AICc  
492 weight = 0.51,  $R^2 = 0.628$ ; Table 1, Appendix S3). Aphid survivorship was 32% higher on native  
493 than invasive plants and declined with increasing latitude (Fig. 3B, Appendix S3: Table S4).  
494 Mean survivorship was estimated to be 94% in the southernmost genotypes and 70% in the  
495 northernmost genotypes (a 24% change). As with colony size, the difference in aphid  
496 survivorship between native and invasive genotypes at LSU (46%) is much greater than the  
497 difference at URI (18%). At URI, the native and invasive lineages exhibited parallel negative  
498 latitudinal clines for aphid survivorship but at LSU, the negative latitudinal clines were not  
499 significant (Appendix S3: Tables S2 and S3). Because we had only a single estimate of aphid  
500 survivorship per *P. australis* source genotype, we could not assess correlations between this trait  
501 and nutritional and defense levels on a per-plant basis.

502 For the four traits associated with *P. australis* palatability to *S. frugiperda*, the main  
503 sources of variation in the AICc-best models were garden and lineage (Appendix S3). For all  
504 traits, native genotypes were more palatable than invasive genotypes to *S. frugiperda* larvae.  
505 Larval growth rate, survivorship, leaf-area consumed and biomass conversion efficiency were  
506 61%, 11%, 114% and 33% higher on native than invasive genotypes, respectively (Fig. 3C-F,  
507 Appendix S3: Table S4). Interestingly, LSU plants were more palatable than URI plants. After  
508 accounting for the effects of lineage, larvae grew 142% larger, had 17% higher survivorship,  
509 consumed 65% more leaf tissue, and were 58% more efficient at converting plant biomass to  
510 larval biomass on LSU than URI plants (Fig. 3C-F, Appendix S3: Table S5). With regard to  
511 biomass conversion efficiency, the difference between lineages was only evident in the URI  
512 garden (garden  $\times$  lineage interaction) and the relationship between conversion efficiency and  
513 latitude was affected by both lineage and garden (lineage  $\times$  latitude and garden  $\times$  latitude  
514 interactions). At the LSU garden, there was no latitudinal gradient in conversion efficiency (Fig.  
515 3F, Appendix S3: Table S2). However, at the URI garden, conversion efficiency declined with  
516 increasing latitude but only for the invasive genotypes (but no lineage  $\times$  latitude interaction; Fig.  
517 3F, Appendix S3: Tables S2 and S3). Finally, *S. frugiperda* palatability was generally negatively  
518 correlated with leaf toughness and positively correlated with % nitrogen (Appendix S4). The  
519 correlations were strongest in the LSU garden where leaf toughness was significantly negatively  
520 related to larval growth rate ( $R = -0.21$ ,  $P < 0.01$ ) and leaf-area chewed ( $R = -0.20$ ,  $P < 0.01$ ) and  
521 nitrogen was significantly positively correlated with biomass conversion efficiency ( $R = 0.40$ ,  $P$   
522  $= 0.04$ ). At URI, the direction of the relationships between leaf toughness and % nitrogen and *S.*  
523 *frugiperda* palatability were similar to those found in the LSU garden but only one relationship  
524 was significant - larval growth rate and % nitrogen ( $R = 0.60$ ,  $P = 0.05$ ). Interestingly, water  
525 content was negatively correlated with leaf-area chewed in the LSU garden ( $R = -0.26$ ,  $P = 0.04$ ).

526

#### 527 *Plasticity in trait responses*

528 We found strong evidence that plant traits associated with herbivory in *P. australis* were  
529 phenotypically plastic, more so for invasive than for native genotypes. Averaged across all 10  
530 traits (aphid traits were excluded; see Methods), mean  $\pm$  SE plasticity (proportional difference  
531 between the two gardens) for the invasive genotypes was  $0.23 \pm 0.11$  and for the native was  $0.09$   
532  $\pm 0.08$ ; a statistically significant 156% difference ( $t_9 = 2.708$ ,  $P = 0.024$ , Fig. 4). These large

533 differences in trait expression between gardens, for both lineages, likely underlie the ubiquitous  
534 garden effect in the previous analyses (Appendix S3: Table S1).

535 Irrespective of *P. australis* lineage, plants reared at the URI garden were more nutritious  
536 and less well defended than those at LSU. Plants reared at URI had 7% greater leaf water content  
537 and had 24% more nitrogen than those at LSU, whereas plants at LSU produced 131% tougher  
538 leaves, 58% more total phenolics and 23% greater CN ratio than plants at URI (Appendix S5).  
539 Despite the higher defenses and lower nutrition at LSU, plants grown at LSU were more  
540 palatable to herbivores than those at URI. Aphid colony size was 57% greater at LSU than at  
541 URI (Appendix S5). *S. frugiperda* larvae consumed two times more leaf area and exhibited a  
542 244% higher growth rate at LSU than in URI (Appendix S5). Finally, biomass conversion  
543 efficiency of larvae was 137% greater at LSU than URI (Appendix S5).

544 When trait plasticity was measured for each *P. australis* genotype, we found that it varied  
545 with latitude for 4 of 6 traits in native genotypes and only 1 of 6 traits in invasive genotypes (Fig.  
546 5, Appendix S6). For native genotypes, plasticity in leaf toughness, water content, and %  
547 nitrogen decreased (Fig. 5A,C,D) and plasticity in the CN ratio increased (Fig. 5F) with  
548 increasing latitude of origin. For the invasive genotypes, leaf toughness was the only trait in  
549 which plasticity varied with latitude – a relationship that paralleled the one for native genotypes  
550 (Fig. 5A). The difference between lineages in the relationship between plasticity and latitude for  
551 total phenolics, water content, % nitrogen and CN ratio are indicated by a significant lineage  $\times$   
552 latitude interaction in the model ANCOVAs (Appendix S6). We also examined whether these  
553 latitudinal trends were driven by invasive genotypes collected from southern latitudes beyond the  
554 distributional range of native genotypes. Re-analyses of these data excluding those genotypes  
555 (collected from the sites south of 36° latitude) did not qualitatively alter the results.

556

557

## DISCUSSION

558 The juxtaposition of plant-species invasions with large-scale gradients in herbivore  
559 pressure and native plant-herbivore interactions is an important yet mostly unexplored issue in  
560 the field of invasion biology (Bezemer et al. 2014, Cronin et al. 2015). Our study is the first to  
561 demonstrate genetic-based latitudinal clines for traits related to plant-herbivore interactions  
562 involving sympatric invasive and native species (or lineages of the same species). These data  
563 suggest that clines in the invasive genotypes of *P. australis* evolved within the past ~ 100 years.

564 In 36% of the cases where clines were evident, the clines for native and invasive lineages were  
565 not parallel. Moreover, our study showed that the occurrence and direction of latitudinal clines in  
566 plant traits was commonly dependent on where the study was conducted (LSU or URI), strongly  
567 suggesting that environmental context is an important driver of the expression of plant traits and  
568 clines in those traits. Until now, this phenotypic plasticity in latitudinal clines for traits related to  
569 species interactions has never been reported. Finally, we found that invasive genotypes were 2.5  
570 times more plastic than native genotypes but the native genotypes were much more likely to  
571 exhibit significant latitudinal variability in phenotypic plasticity than the invasive genotypes.  
572 Overall, this study suggests that traits associated with *P. australis* interactions with its herbivores  
573 are under strong genetic and environmental controls and they vary between co-occurring native  
574 and invasive *P. australis* genotypes across their latitudinal range in eastern North America. Such  
575 spatial heterogeneity within and between lineages with respect to their interactions with  
576 herbivores has the potential to generate substantial spatial heterogeneity in biotic  
577 resistance/susceptibility that can have important implications for the establishment and spread of  
578 invasive genotypes and species.

579

#### 580 *Latitudinal clines in plant-herbivore traits*

581 ***Hypothesis 1: Genetic-based latitudinal clines for plant defenses and palatability to herbivores***  
582 ***are evident for native *P. australis* genotypes.***

583 ***Hypothesis 2: Because invasive genotypes may not have had sufficient time to evolve in***  
584 ***response to its novel environment, latitudinal clines in defense and palatability traits***  
585 ***will be absent or weak, and thus non-parallel to the gradients for the native genotypes.***

586

587 *Genetic-based clines.* In our study, support for the hypotheses that native genotypes  
588 should be more likely to exhibit genetic-based latitudinal clines than invasive genotypes  
589 (Hypotheses 1 and 2) was mixed. At least one cline was evident for each defense and nutritional  
590 trait, and for at least one trait associated with palatability to *H. pruni* and *S. frugiperda*.  
591 However, between the two gardens, there were twice as many latitudinal clines in plant-  
592 herbivore traits for the invasive genotypes compared to the native genotypes (8 vs 4 clines,  
593 respectively; Appendix S3: Table S3). Moreover, in only one instance was a latitudinal cline for  
594 a particular trait evident for both lineages (water content).

595           Although it was strongest in the LSU garden, leaf toughness of native and invasive  
596 genotypes generally increased with decreasing latitude. As herbivore pressure in naturally  
597 occurring patches of *P. australis* is higher at lower latitudes (Cronin et al. 2015), our common-  
598 garden study supports the presumed role of this trait as a defense against herbivores (e.g., Raupp  
599 1985, Salgado and Pennings 2005). Its role as a defense is further supported by the finding that  
600 palatability to *S. frugiperda* was generally negatively correlated with leaf toughness (see below).  
601 Our results with leaf toughness are also consistent with the prediction that longer lifespans of  
602 leaves in areas with longer growing seasons should favor tougher leaves (Coley and Aide 1991,  
603 Salgado and Pennings 2005). The lone positive latitudinal cline for total phenolics, evident for  
604 native genotypes at URI, was the opposite of what we observed in the field (Cronin et al. 2015)  
605 and is counter to our expectations if this trait is related to herbivore defense. However, total  
606 phenolics represent a broad class of compounds that serve other purposes for plants including  
607 protection against photodamage (Close and McArthur 2002). Moles et al. (2011) reported that ca.  
608 30% of the published studies found a significant latitudinal gradient in plant total phenolics.  
609 Interestingly, the majority of those significant gradients were in the direction observed for the  
610 native genotypes from the URI garden. Finally, these results are also consistent with the findings  
611 from the meta-analysis of common-garden studies by Colautti et al. (2009) in which there was no  
612 consistent directionality in latitudinal clines for plant defense traits, and that evidence for clines  
613 in the field do not always match up with clines for the same species in the common garden.

614           Latitudinal gradients in foliar nutrient levels are commonly reported (e.g., Siska et al.  
615 2002, Reich and Oleksyn 2004, Lovelock et al. 2007, De Frenne et al. 2013, He and Silliman  
616 2015). Between gardens, there was a trend toward decreasing % carbon with increasing latitude,  
617 the opposite pattern to that observed in nature (e.g., De Frenne et al. 2013). It is possible that  
618 plants adapted to the CO<sub>2</sub>-rich environments in the north may have evolved to be less efficient at  
619 uptaking or utilizing CO<sub>2</sub> for photosynthesis or storing carbon compounds into their tissues than  
620 plants from the relatively CO<sub>2</sub>-poor environments in the south (see e.g., Denning et al. 1995).  
621 For example, plant stomata are known to stay open longer and wider or the density of stomata  
622 increase when the concentration of atmospheric CO<sub>2</sub> decreases in order to maintain an adequate  
623 CO<sub>2</sub> gradient between the atmosphere and the leaf (Beerling et al. 1998). It is possible that *P.*  
624 *australis* adapted to their local CO<sub>2</sub> concentrations and consequently, southern plants were more  
625 efficient at producing or storing carbon compounds. Alternatively, carbon-rich tissues at



626 southern latitudes could serve as a defense against greater herbivore pressure in the south (Orians  
627 and Milewsky 2007).

628 We found evidence for latitudinal clines for % nitrogen, CN ratio, and water content but  
629 they varied between gardens and lineages. Although nitrogen content in coastal wetland plants  
630 tends to increase with increasing latitude (Siska et al. 2002, He and Silliman 2015), we found  
631 both positive (URI garden) and negative (LSU garden) latitudinal clines for % nitrogen in native  
632 *P. australis* genotypes. Similar results were found for the CN ratio and water content. These  
633 differences between lineages and gardens clearly indicate that latitudinal clines in *P. australis*  
634 traits are phenotypically plastic (Woods et al. 2012; see section “Phenotypic plasticity in defense  
635 and palatability traits”).

636 The genetic-based clines for palatability to *H. pruni* (aphid colony growth and  
637 survivorship decreased with increasing latitude; although more strongly for the invasive lineage)  
638 do not appear to be caused by latitudinal variation in the *P. australis* nutritional and defensive  
639 traits. Although aphid colony growth was positively correlated with total phenolics and water  
640 content of leaves, neither of those variables was positively correlated with latitude. Latitudinal  
641 gradients in palatability to aphids could be explained by the fact that aphid abundance on *P.*  
642 *australis* increases with increasing latitude in North America (Cronin et al. 2015). At northern  
643 latitudes, strong selection pressure by these herbivores may have resulted in higher resistance or  
644 lower palatability.

645 Because *H. pruni* was introduced to North America not long after the introduction of  
646 invasive genotypes of *P. australis* (Lozier et al. 2009), both native and invasive lineages likely  
647 had an equivalent amount of time to adapt to the gradient in aphid abundance. The existence of  
648 parallel clines between native and invasive genotypes suggests that both *P. australis* lineages  
649 evolved in similar ways to the latitudinal variation in aphid abundance. A few studies to date  
650 have reported the formation of clines along an environmental gradient (latitudinal or elevational)  
651 by an invasive species that parallels the clines in the native range (Maron et al. 2004, Alexander  
652 et al. 2009). However, most previous studies (e.g., Colautti et al. 2009, Maron et al. 2004)  
653 involved cross-continental comparisons between native and invasive species and so,  
654 environmental conditions, even at similar latitudes, may not be the same. In contrast, our native  
655 and invasive genotypes are found in the same marshes along the East and Gulf Coasts and  
656 experience the same environmental gradient.

657 With only one exception (negative correlation between biomass conversion efficiency  
658 and latitude for the invasive genotypes in the URI garden), there was very little evidence of  
659 genetic-based clines for palatability to *S. frugiperda*. In nature, chewing damage to native  
660 genotypes of *P. australis* is strongly negatively correlated with latitude (Cronin et al. 2015). In  
661 response to these selection pressures, we expected that the native genotypes would have evolved  
662 a positive latitudinal cline in palatability to native grass-feeding generalists like *S. frugiperda*.  
663 The fact that a gradient was evident for the invasive genotypes, and that it was in the opposite  
664 direction predicted for the native genotypes is surprising. Finally, we suggest that the negative  
665 relationship between leaf toughness and latitude (Fig. 2) could be the mechanism driving the  
666 negative latitudinal gradient in biomass conversion efficiency in the invasive lineage.

667 *Non-parallel latitudinal gradients between native and invasive genotypes.* One interesting  
668 and important finding regarding native and invasive genotypes of *P. australis* was that for 20%  
669 of the cases (5 of 24; 12 traits  $\times$  2 gardens), the relationship between latitude and trait level was  
670 different for native and invasive genotypes from the same garden (in support of Hypothesis 2).  
671 We observed non-parallel clines in water content, total phenolics, % nitrogen, and CN ratio (all  
672 but the first occurring in the URI garden). In the case of total phenolics, % nitrogen, and CN  
673 ratio, the slope of the relationship with latitude differed in sign between the native and invasive  
674 genotypes. These results suggest that the two lineages have evolved in different ways to the same  
675 environmental gradient. Although there are many studies that have examined latitudinal clines  
676 for species in their native and invasive ranges (Colautti et al. 2009), environmental differences  
677 make clinal comparisons questionable. Because the native and invasive ranges of *P. australis*  
678 overlap on the same continent, our study provides a much stronger example of clinal evolution in  
679 invasive taxa. Although co-occurring native and invasive genotypes across a broad latitudinal  
680 range is known only for *P. australis*, a number of other species have co-occurring and distinct  
681 invasive genotypes or native-invasive hybrids (Ayres et al. 2004, Lavergne and Molofsky 2007,  
682 Ciotir et al. 2013). These are potentially fertile systems for the study of cline formation and  
683 evolution. These non-parallel latitudinal clines between co-occurring native and invasive  
684 genotypes of *P. australis* can result in spatially varying degrees of local enemy release and biotic  
685 resistance (see “Implications for invasion success”).

686 *Origin of latitudinal clines.* We contend that the clines described above for the invasive  
687 genotypes of *P. australis* must have arisen *de novo* while in North America. A pre-existing cline,

688 i.e., one that evolved in Europe and subsequently transferred virtually intact to North America, is  
689 possible but unlikely. For this to occur, there would have to have been multiple introduction  
690 events in which individuals from one latitude in Europe colonized at a similar latitude in North  
691 America. There is no evidence for this sort of parallel invasion process with *P. australis*. Based  
692 on an analysis of chloroplast DNA from herbarium specimens by Saltonstall (2002), European *P.*  
693 *australis* became established in the mid-Atlantic region at least 150 years ago and spread rapidly  
694 north and south (Saltonstall 2002). It is highly suggestive of one or a few major founding events  
695 followed by range expansion. Although other European genotypes are present in North America,  
696 they appear restricted to very northern and southern latitudes (Lambertini et al. 2012, Meyerson  
697 and Cronin 2013). Therefore, our study supports the growing body of literature that invasive  
698 species evolve relatively quickly (< 100 years) in response to an environmental gradient (e.g.,  
699 Maron et al. 2004, Alexander et al. 2009, Li et al. 2015). Furthermore, the absence of a humped  
700 or u-shaped relationship between latitude and each trait, centering on the origin of the invasion  
701 (between Delaware and Connecticut; Saltonstall 2002), suggests that the time since invasion is  
702 not an important driver of the evolution of these gradients in the invasive lineage.

703 Our finding that invasive genotypes were twice as likely as native genotypes to exhibit a  
704 latitudinal cline for plant-herbivore traits is somewhat surprising. Such a result would suggest  
705 that the invasive genotypes in their new range were more evolutionarily responsive to the same  
706 environmental conditions faced by the native genotypes. For two main reasons, this seems  
707 unlikely. First, local adaptation and cline formation requires limited gene flow among  
708 populations distributed along the latitudinal gradient (Slatkin 1985). Populations of native  
709 genotypes are quite rare and isolated in comparison to the extremely widespread and abundant  
710 invasive populations (Chambers et al. 1999, Saltonstall 2002). The native genotypes would seem  
711 much more likely to exhibit local adaptation and genetic isolation-by-distance than invasive  
712 genotypes. This prediction is supported by Kettenring and Mock (2012) who found greater  
713 genetic homogeneity among invasive than native populations of *P. australis*. Second, the  
714 selection pressures from North American herbivores on the invasive genotypes is considerably  
715 weaker than on native genotypes. Based on our latitudinal field surveys of herbivory, invasive  
716 genotypes suffered 70% to 650% lower levels of herbivory from all herbivore guilds (chewers,  
717 gallers, suckers) than native genotypes and that herbivory levels for invasive genotypes did not  
718 vary significantly with latitude (Cronin et al. 2015). Moreover, this study demonstrated that two

719 herbivores, *H. pruni* and *S. frugiperda*, had lower growth and survivorship on the invasive  
720 genotypes. If anything, the high levels and strong latitudinal gradients in herbivory of the native  
721 genotypes should have favored a high frequency of genetic-based latitudinal clines for this  
722 lineage.

723  
724 *Trait variation between replicate common gardens.* One of our strongest and most  
725 obvious finding was that for all traits considered in this study, a garden effect was present in  
726 every model; indicative of significant trait plasticity. Also, interactions involving garden (e.g.,  
727 garden  $\times$  lineage, garden  $\times$  latitude, garden  $\times$  lineage  $\times$  latitude were quite common (see  
728 Appendix S3). These types of garden effects are a frequent occurrence in studies involving  
729 multiple common gardens (e.g., De Frenne et al. 2012, Woods et al. 2012, Zhou et al. 2014).  
730 Also, in their meta-analysis, Colautti et al. (2009) found numerous examples of a garden  $\times$   
731 latitude interaction for studies conducted in both the native and invasive ranges of a plant  
732 species. In some studies, reversals of latitudinal trends were evident between gardens (Chapin  
733 and Chapin 1981, Santamaría et al. 2003, Maron et al. 2004). As the same species or genotype is  
734 represented in each garden, these garden  $\times$  latitude interactions imply plasticity in the regulation  
735 of latitudinal clines (Richards et al. 2006, Woods et al. 2012, Zhou et al. 2014).

736 One possible reason for the substantial differences in trait expression between gardens is  
737 that the gardens reside in very different parts of the range of the two *P. australis* lineages. The  
738 URI garden is near the center of the coastal range for both native and invasive lineages, whereas  
739 the LSU garden is near the southern extent of the invasive lineage and ~700 km south of our  
740 southernmost native population. Plants in the southern garden, particularly for the native lineage,  
741 may be at their thermal limits (e.g., Drake et al. 2015) which could result in altered expression of  
742 genetic-based clines in these plant nutritional traits. Several studies have examined latitudinal  
743 clines in plant traits in replicate common-garden experiments located within, at the boundary or  
744 beyond the limits of the species range (e.g., De Frenne et al. 2012, Woods et al. 2012, Zhou et al.  
745 2014). For example, in their study of plant growth and defense traits in common milkweed  
746 (*Asclepias syriaca*), Woods et al. (2012) found a cline in milkweed phenology in their garden at  
747 the southern range limit (North Carolina) but no cline in their garden at the northern range limit  
748 (New Brunswick, Canada). Woods et al. (2012) also evaluated whether plant defense trait  
749 expression was dependent on the proximity of the milkweed source populations to the center of

750 the species range. According to the ‘range-center’ hypothesis (Alexander et al. 2007, Woods et  
751 al. 2012), because plant abundance and herbivore pressure is expected to be higher near the  
752 range center than range margins, plant defenses are predicted to be humped shaped (peaking at  
753 the range center) and palatability to herbivores should be u-shaped (trough at the range center).  
754 Woods et al.’s (2012) study did not support this hypothesis as traits were linearly related to  
755 latitude. We also find no support for the range-center hypothesis because none of our models  
756 supported a curvilinear relationship between plant traits and latitude (i.e., the quadratic term for  
757 latitude was never retained in our model-selection procedures). One important implication from  
758 our study, and those of De Frenne et al. (2012), Woods et al. (2012), Zhou et al. (2014), is that  
759 because latitudinal clines in plant traits are phenotypically plastic and dependent on climatic  
760 conditions, future climate change may fundamentally alter latitudinal gradients in nature.  
761 Understanding the biogeography of plant-herbivore interactions in the face of climate change is  
762 going to be a daunting task if latitudinal clines in species-interaction traits are universally plastic  
763 as these studies suggest.

764

#### 765 *Phenotypic plasticity in defense and palatability traits*

766 *Hypothesis 3: Invasive genotypes are more phenotypically plastic than native genotypes.*

767 *Hypothesis 4: Plasticity varies with latitude.*

768 *Hypothesis 5: Latitudinal clines are phenotypically plastic.*

769

770 A fundamental question in the field of invasion biology is what traits promote invasion  
771 success. Phenotypic plasticity is thought to be beneficial to founding populations by increasing  
772 niche breadth (e.g., Richards et al. 2006, Chun 2011, Davidson et al. 2011) or plasticity may be a  
773 trait that evolves in response to the novel environments (Richards et al. 2006, Lavergne and  
774 Molofsky 2007). In agreement with the meta-analysis by Davidson et al. (2011), we found that  
775 invasive genotypes were 2.5 times more plastic than native genotypes (supporting Hypothesis 3).  
776 Because we prevented these genotypes from evolving in response to the biotic and/or abiotic  
777 environment (see Methods “Plant defense, nutritional and palatability traits in common  
778 gardens”), these differences in trait expression between gardens are attributed to phenotypic  
779 plasticity. We also found that trait plasticity varied linearly with latitude (supporting Hypothesis  
780 4), although interestingly, it was primarily for the native genotypes (4 of 6 traits for the native

781 and 1 of 6 traits for the invasive genotypes). Similar linear relationships between latitude and  
782 plasticity have been reported elsewhere (e.g., Maron et al. 2004, De Frenne et al. 2011, Woods et  
783 al. 2012). It is possible that native genotypes under stress from high thermal limits or high  
784 herbivore pressure (Cronin et al. 2015) near the southern range limit of the lineage, had to evolve  
785 to be more plastic, supporting Chevin and Lande's (2011) argument that plasticity should be  
786 greater at the range margins. In contrast, the southern range limit for invasive *P. australis* is the  
787 Gulf of Mexico and geographic barriers may limit their southern range, not high temperature.  
788 Coupled with low levels of herbivory, the invasive genotypes may not be as stressed as native  
789 genotypes at lower latitudes. Finally, as we have discussed previously ("Trait variation between  
790 replicate common gardens"), we found evidence that for a specific lineage, latitudinal clines  
791 differed between gardens, supporting Hypothesis 5 that clines are phenotypically plastic.  
792 Latitudinal cline plasticity was evident for total phenolics, water content, % nitrogen and CN  
793 ratio (see Fig. 2). Latitudinal cline plasticity appears to be a common occurrence in studies  
794 involving multiple common gardens (see Colautti et al. 2009). However, this is the first study to  
795 show clinal plasticity at the sub-specific level.

796

797

#### *Implications for invasion success*

798

***Hypothesis 6: Across a broad latitudinal range in North America, European genotypes of *P.****

799

***australis are successful invaders because they are better defended and less palatable to***

800

***herbivores than native genotypes.***

801

802

Enemy release, i.e., reduced pressure by natural enemies in the introduced relative to the  
803 native range (Keane and Crawley 2002), is one of the more prominent hypotheses invoked to  
804 explain invasion success (e.g., Mitchell and Power 2003, Liu and Stiling 2006). Biogeographic  
805 studies comparing enemy pressure on a plant species in its native and invaded range support this  
806 hypothesis (e.g., Mitchell and Power 2003, Liu and Stiling 2006). However, invasion success is  
807 also likely to be dependent on the invasive species being less vulnerable to natural enemies than  
808 sympatric native species (i.e., "local enemy release"; Zheng et al. 2012), and support for this  
809 outcome is mixed (Colautti et al. 2004, Chun et al. 2010). Our field work with the European  
810 genotypes of *P. australis* revealed that not only was herbivory significantly lower in their  
811 invaded than native range, but also that they suffer 70% - 650% less herbivory in North America

812 as compared to North American native genotypes (Cronin et al. 2015). In addition, our common-  
813 garden study demonstrates an underlying genetic basis to lower levels of herbivory of European  
814 genotypes of *P. australis* – they are less palatable to *H. pruni* and *S. frugiperda*. Invasive  
815 genotypes had aphid colonies that were 91% smaller, and had *S. frugiperda* that consumed 54%  
816 less leaf material and exhibited 38% lower growth rates than native genotypes (supporting  
817 Hypothesis 6). Contrary to hypothesis 6, we did not find evidence that leaf toughness and total  
818 phenolics, our two putative measures of *P. australis* resistance, were greater for invasive than  
819 native genotypes. In fact, native genotypes had more total phenolics (Appendix S3: Table S4).  
820 One possible explanation for the lower palatability of invasive plants is that leaf water content is  
821 8% lower than for native plants. High water content is beneficial to many insect species (e.g.,  
822 Huberty and Denno 2004). Finally, not only are invasive genotypes of *P. australis* less palatable  
823 to herbivores than native genotypes, they are also more plastic with regard to the traits that likely  
824 influence palatability, including total phenolics, water content, % nitrogen and CN ratio. Flexible  
825 genotypes may not only be an adaptation to a variable environment (e.g., Agrawal 2001) but may  
826 also make it more difficult for herbivores to adapt to its host or track host resources in space  
827 (Denno 1983). Given the levels of herbivory observed for *P. australis* in North America, and the  
828 potential impact on plant fitness (Lambert and Casagrande 2007, Lambert et al. 2007, Cronin et  
829 al. 2015, Bhattarai et al. 2016, Cronin et al. 2016), the invasive genotypes of *P. australis* are at a  
830 significant advantage over native genotypes. It is no surprise that wetland communities in North  
831 America did not resist invasion by European genotypes of *P. australis*.

832 There is an important biogeographic component to enemy-release for invasive *P.*  
833 *australis*. A strong negative relationship between latitude and herbivory for the native genotypes  
834 but no latitudinal gradient for the invasive genotypes results in the strength of enemy release  
835 being greater at lower than higher latitudes (Cronin et al. 2015). Although it remains to be tested,  
836 we hypothesized that the likelihood of establishment and rate of spread of invasive *P. australis*  
837 genotypes should be greatest at lower latitudes (but see Bhattarai and Cronin 2014). We also  
838 suggested that non-parallel gradients in herbivory between sympatric native and invasive species  
839 or genotypes are likely to be a common occurrence, especially for recent invaders.

840 Based on our replicated common-garden study, the biogeography of this plant-herbivore  
841 interaction is much more complicated than previously envisioned. Latitudinal clines in traits  
842 potentially important to the interaction between *P. australis* and its herbivores evolved quickly

843 ( $\approx 100$  years) for the invasive genotypes, there is plasticity in the regulation of those clines, and  
844 in a number of cases, the clines run counter to those for the native genotypes. All of this adds up  
845 to substantial heterogeneity in the interactions between native and invasive *P. australis* and their  
846 shared herbivores. For example, Bhattarai et al. (in review) found significant variability along the  
847 Atlantic Coast of the United States in the strength of apparent competition (mediated by their  
848 shared herbivores) between native and invasive *P. australis* genotypes. Because studies with  
849 replicate common gardens often have found similar plasticity in latitudinal clines of invasive  
850 species (Richards et al. 2006, Woods et al. 2012, Zhou et al. 2014), we argue that large-scale  
851 plant invasions are characterized by considerable spatial heterogeneity in herbivore impact on  
852 native and invasive species (or genotypes). This heterogeneity may also explain why support for  
853 local enemy release is inconsistent among studies (see e.g., Colautti et al. 2004, Chun et al.  
854 2010).

855

856

### Concluding remarks

857 Although this study was conducted at the sub-specific level (i.e., lineages of the same  
858 species), we strongly believe that our results and conclusions are applicable to any native-  
859 invasive plant system, whether the participants differ at the species, genus or higher taxonomic  
860 level. If anything, we would have expected differences in plant traits, clines and plasticity at the  
861 sub-specific level to be much more difficult to detect than at higher taxonomic levels. As such,  
862 our study suggests that these biogeographic differences between native and invasive taxa,  
863 particularly early on in the establishment and spread of invasive taxa, are likely to be common  
864 and significant. Lastly, by conducting our study at the sub-specific level, phylogenetic  
865 differences between native and invasive taxa that could underlie differences in biogeographic  
866 patterns, are less likely to be an issue. This represents one of the great strengths of the *P.*  
867 *australis* study system and is one reason why we consider this a model system for studying  
868 species invasions (Meyerson et al. 2016).

869 Finally, replicate garden studies have proven to be very informative to understanding  
870 plant-herbivore interactions. Those positioned in climatically different environments or at  
871 different locations within the species range have been especially fruitful (e.g., Woods et al.  
872 2012). Continued research in this area should focus on experimental manipulations at key  
873 locations within, at the boundary, and beyond the invasion range, particularly focusing on the



874 impact of natural enemies (herbivores and pathogens) on local plant fitness, the role of generalist  
875 and specialist natural enemies, interactions with other sympatric native and invasive plant  
876 species and higher trophic levels. As our most pernicious invaders typically have broad invasion  
877 fronts, the biogeography of their interactions with native species cannot be ignored if we wish to  
878 understand what has led to their success.

879

#### 880 **ACKNOWLEDGEMENTS**

881 We thank A. Accamando, B. Bhandari, L. Brus, B. Chelette, A. Chow, J. Croy, E.  
882 Francine, C. Lee, P. Mooney, P. Panta, S. Rinehart, T. Scavuzzo-Duggan, V. Tran, D. Troiano  
883 and R. Young for assistance in common garden experiments. M. Stout assisted us with the  
884 quantification of total phenolics. B. Elderd and F. Dilleuth provided advice on rearing fall  
885 armyworm caterpillars. We thank J. Reeve for providing help on Statistics, and K. Brown, K.  
886 Harms, R. Stevens and six anonymous reviewers for valuable suggestions on the manuscript. The  
887 research was funded by NSF research grants 1050084 to JTC and 1049914 to LAM. DC was  
888 supported through an NSF REU fellowship to LAM. Additional support was provided to GPB  
889 from grants from the Louisiana Environmental Education Commission and LSU Biograds.

890

#### 891 **LITERATURE CITED**

- 892 Agrawal, A. A. 2001. Phenotypic plasticity in the interaction and evolution of species. *Science*  
893 **294**:321-326.
- 894 Agrawal, A. A. 2004. Plant defense and density dependence in the population growth of  
895 herbivores. *American Naturalist* **164**:113-120.
- 896 Alexander, H. M., S. Price, R. Houser, D. Finch, and M. Tourtellot. 2007. Is there reduction in  
897 disease and pre-dispersal seed predation at the border of a host plant's range? Field and  
898 herbarium studies of *Carex blanda*. *Journal of Ecology* **95**:446-457.
- 899 Alexander, J. M., P. J. Edwards, M. Poll, C. G. Parks, and H. Dietz. 2009. Establishment of  
900 parallel altitudinal clines in traits of native and introduced forbs. *Ecology* **90**:612-622.
- 901 Allen, W. J., R. E. Young, G. P. Bhattarai, J. R. Croy, A. M. Lambert, L. A. Meyerson, and J. T.  
902 Cronin. 2015. Multirophic enemy escape of invasive *Phragmites australis* and its  
903 introduced herbivores in North America. *Biological Invasions* **12**:3419-3432.

- 904 Andrew, N. R., and L. Hughes. 2005. Herbivore damage along a latitudinal gradient: relative  
905 impacts of different feeding guilds. *Oikos* **108**:176-182.
- 906 Ayres, D. R., D. L. Smith, K. Zaremba, S. Klohr, and D. R. Strong. 2004. Spread of exotic  
907 cordgrasses and hybrids (*Spartina* sp.) in the tidal marshes of San Francisco Bay,  
908 California, USA. *Biological Invasions* **6**:221-231.
- 909 Bates, D., M. Maechler, B. Bolker, and S. Walker. 2014. lme4: Linear mixed-effects models  
910 using Eigen and S4. R package version 1.1-7. <http://CRAN.R-project.org/package=lme4>.
- 911 Beerling, D. J., J. C. McElwain, and C. P. Osborne. 1998. Stomatal responses of the 'living  
912 fossil' *Ginkgo biloba* L. to changes in atmospheric CO<sub>2</sub> concentrations. *Journal of*  
913 *Experimental Botany* **49**:1603-1607.
- 914 Bezemer, T. M., J. A. Harvey, and J. T. Cronin. 2014. Response of native insect communities to  
915 invasive plants. *Annual Review of Entomology* **59**:119-141.
- 916 Bhattarai, G. P. 2015. Biogeographical approaches for studying species invasion. Louisiana State  
917 University, Baton Rouge, Louisiana, USA.
- 918 Bhattarai, G. P., and J. T. Cronin. 2014. Hurricane activity and the large-scale pattern of spread  
919 of an invasive plant species. *Plos One* **9**:e98478.
- 920 Bhattarai, G. P., L. A. Meyerson, and J. T. Cronin. 2016. Geographical variation in apparent  
921 competition between native and invasive *Phragmites australis*. *Ecology in review*.
- 922 Bhattarai, G. P., W. J. Allen, J. T. Cronin, E. Kiviat, and L. A. Meyerson. 2016. Response to  
923 Blossey and Casagrande – Ecological and evolutionary processes make host specificity at  
924 the subspecies level exceedingly unlikely. *Biological Invasions* **18**:2757-2758.
- 925 Bolker, B. M., M. E. Brooks, C. J. Clark, S. W. Geange, J. R. Poulsen, M. H. H. Stevens, and J.-  
926 S. S. White. 2009. Generalized linear mixed models: a practical guide for ecology and  
927 evolution. *Trends in ecology & evolution* **24**:127-135.
- 928 Bolser, R. C., and M. E. Hay. 1996. Are tropical plants better defended? Palatability and  
929 defenses of temperate vs. tropical seaweeds. *Ecology* **77**:2269-2286.
- 930 Burnham, K. P., and D. R. Anderson. 2010. *Model Selection and Multimodel Inference: A*  
931 *Practical Information-Theoretic Approach*. 2 edition. Springer.
- 932 Chambers, R. M., L. A. Meyerson, and K. Saltonstall. 1999. Expansion of *Phragmites australis*  
933 into tidal wetlands of North America. *Aquatic Botany* **64**:261-273.

- 934 Chapin, F. S., and M. C. Chapin. 1981. Ecotypic differentiation of growth processes in *Carex*  
935 *aquatilis* along latitudinal and local gradients. *Ecology* **62**:1000-1009.
- 936 Chevin, L. M., and R. Lande. 2011. Adaptation to marginal habitats by evolution of increased  
937 phenotypic plasticity. *Journal of Evolutionary Biology* **24**:1462-1476.
- 938 Chun, Y. J. 2011. Phenotypic plasticity of introduced versus native purple loosestrife: univariate  
939 and multivariate reaction norm approaches. *Biological Invasions* **13**:819-829.
- 940 Chun, Y. J., M. Van Kleunen, and W. Dawson. 2010. The role of enemy release, tolerance and  
941 resistance in plant invasions: linking damage to performance. *Ecology Letters* **13**:937-  
942 946.
- 943 Ciotir, C., H. Kirk, J. R. Row, and J. R. Freeland. 2013. Intercontinental dispersal of *Typha*  
944 *angustifolia* and *T. latifolia* between Europe and North America has implications for  
945 *Typha* invasions. *Biological Invasions* **15**:1377-1390.
- 946 Close, D. C., and C. McArthur. 2002. Rethinking the role of many plant phenolics - protection  
947 from photodamage not herbivores? *Oikos* **99**:166-172.
- 948 Colautti, R. I., A. Ricciardi, I. A. Grigorovich, and H. J. MacIsaac. 2004. Is invasion success  
949 explained by the enemy release hypothesis? *Ecology Letters* **7**:721-733.
- 950 Colautti, R. I., J. L. Maron, and S. C. Barrett. 2009. Common garden comparisons of native and  
951 introduced plant populations: latitudinal clines can obscure evolutionary inferences.  
952 *Evolutionary Applications* **2**:187-199.
- 953 Coley, P. D. 1986. Costs and benefits of defense by tannins in a neotropical tree. *Oecologia*  
954 **70**:238-241.
- 955 Coley, P. D., and T. Aide. 1991. Comparison of herbivory and plant defenses in temperate and  
956 tropical broad-leaved forests. Pages 25-49 in P. W. Price, T. M. Lewinsohn, G. W.  
957 Fernandes, and W. W. Benson, editors. *Plant-animal interactions: evolutionary ecology in*  
958 *tropical and temperate regions*. Wiley-Interscience.
- 959 Cronin, J. T., E. Kiviat, L. A. Meyerson, G. P. Bhattarai, and W. J. Allen. 2016. Biological  
960 control of invasive *Phragmites australis* will be detrimental to native *P. australis*.  
961 *Biological Invasions* **18**:2749-2752.
- 962 Cronin, J. T., G. P. Bhattarai, W. J. Allen, and L. A. Meyerson. 2015. Biogeography of a plant  
963 invasion: plant-herbivore interactions. *Ecology* **96**:1115-1127.

964 Davidson, A. M., M. Jennions, and A. B. Nicotra. 2011. Do invasive species show higher  
965 phenotypic plasticity than native species and, if so, is it adaptive? A meta-analysis.  
966 Ecology Letters **14**:419-431.

967 De Frenne, P., B. J. Graae, F. Rodriguez-Sanchez, A. Kolb, O. Chabrierie, G. Decocq, H. De  
968 Kort, A. De Schrijver, M. Diekmann, O. Eriksson, R. Gruwez, M. Hermy, J. Lenoir, J.  
969 Plue, D. A. Coomes and K. Verheyen. 2013. Latitudinal gradients as natural laboratories  
970 to infer species' response to temperature. Journal of Ecology 101:784-795.

971 De Frenne, P., B. J. Graae, J. Brunet, A. Shevtsova, A. De Schrijver, O. Chabrierie, S. A. O.  
972 Cousins, G. Decocq, M. Diekmann, M. Hermy, T. Heinken, A. Kolb, C. Nilsson, S.  
973 Stanton, and K. Verheyen. 2012. The response of forest plant regeneration to temperature  
974 variation along a latitudinal gradient. Annals of Botany **109**:1037-1046.

975 De Frenne, P., J. Brunet, A. Shevtsova, A. Kolb, B. J. Graae, O. Chabrierie, S. A. Cousins, G.  
976 Decocq, A. N. De Schrijver, M. Diekmann, R. Gruwez, T. Heinken, M. Hermy, C.  
977 Nilsson, S. Stanton, W. Tack, J. Willaert, and K. Verheyen. 2011. Temperature effects on  
978 forest herbs assessed by warming and transplant experiments along a latitudinal gradient.  
979 Global Change Biology **17**:3240-3253.

980 Denning, A. S., I. Y. Fung, and D. Randall. 1995. Latitudinal gradient of atmospheric CO<sub>2</sub> due  
981 to seasonal exchange with land biota. Nature **376**:240-243.

982 Denno, R. F. 1983. Tracking variable host plants in space and time. Pages 291-241 in R. F.  
983 Denno and M. S. McClure, editors, Variable plants and herbivores in natural and  
984 managed systems. Academic Press, New York.

985 Dobzhansky, T. 1950. Evolution in the tropics. American Scientist **38**:209-221.

986 Drake, J. E., M. J. Aspinwall, S. Pfautsch, P. D. Rymer, P. B. Reich, R. A. Smith, K. Y. Crous,  
987 D. T. Tissue, O. Ghannoum, and M. G. Tjoelker. 2015. The capacity to cope with climate  
988 warming declines from temperate to tropical latitudes in two widely distributed  
989 *Eucalyptus* species. Global Change Biology **21**:459-472.

990 He, Q., and B. R. Silliman. 2015. Biogeographic consequences of nutrient enrichment for plant-  
991 herbivore interactions in coastal wetlands. Ecology Letters **18**:462-471.

992 Hillebrand, H. 2004. On the generality of the latitudinal diversity gradient. American Naturalist  
993 **163**:192-211.

- 994 Huberty, A. F. and R. F. Denno. 2004. Plant water stress and its consequences for herbivorous  
995 insects: A new synthesis. *Ecology* **85**:1383-1398.
- 996 Johnson, M. T. 2008. Bottom-up effects of plant genotype in aphids, ants, and predators.  
997 *Ecology* **89**:145-154.
- 998 Keane, R. M., and M. J. Crawley. 2002. Exotic plant invasions and the enemy release hypothesis.  
999 *Trends in Ecology and Evolution* **17**:164-170.
- 1000 Kettenring, K. M., and K. E. Mock. 2012. Genetic diversity, reproductive mode, and dispersal  
1001 differ between the cryptic invader, *Phragmites australis*, and its native conspecific.  
1002 *Biological Invasions* **14**:2489-2504.
- 1003 Kim, T. N. 2014. Plant damage and herbivore performance change with latitude for two old-field  
1004 plant species, but rarely as predicted. *Oikos* **123**:886-896.
- 1005 Kozlov, M. V., A. V. Stekolshchikov, G. Söderman, E. S. Labina, V. Zverev, and E. L. Zvereva.  
1006 2015. Sap-feeding insects on forest trees along latitudinal gradients in northern Europe: a  
1007 climate-driven patterns. *Global Change Biology* **21**:106–116.
- 1008 Lambert, A. M., and R. A. Casagrande. 2007. Susceptibility of native and non-native common  
1009 reed to the non-native mealy plum aphid (Homoptera: Aphididae) in North America.  
1010 *Environmental Entomology* **36**:451-457.
- 1011 Lambert, A. M., K. Winiarski, and R. A. Casagrande. 2007. Distribution and impact of exotic  
1012 gall flies (*Lipara* sp.) on native and exotic *Phragmites australis*. *Aquatic Botany* **86**:163-  
1013 170.
- 1014 Lambertini, C., I. A. Mendelssohn, M. H. Gustafsson, B. Olesen, R. Tenna, B. K. Sorrell, and H.  
1015 Brix. 2012. Tracing the origin of Gulf Coast *Phragmites* (Poaceae): a story of long-  
1016 distance dispersal and hybridization. *American Journal of Botany* **99**:538-551.
- 1017 Lambertini, C., M. H. G. Gustafsson, J. Frydenberg, J. Lissner, M. Speranza, and H. Brix. 2006.  
1018 A phylogeographic study of the cosmopolitan genus *Phragmites* (Poaceae) based on  
1019 AFLPs. *Plant Systematics and Evolution* **258**:161-182.
- 1020 Lavergne, S., and J. Molofsky. 2007. Increased genetic variation and evolutionary potential drive  
1021 the success of an invasive grass. *Proceedings of the National Academy of Sciences*  
1022 **104**:3883-3888.

- 1023 Lehndal, L. and J. Agren. 2015. Latitudinal variation in resistance and tolerance to herbivory in  
1024 the perennial herb *Lythrum salicaria* is related to intensity of herbivory and plant  
1025 phenology. *Journal of Evolutionary Biology* **28**:576-589.
- 1026 Levine, J. M., P. B. Adler, and S. G. Yelenik. 2004. A meta-analysis of biotic resistance to exotic  
1027 plant invasions. *Ecology Letters* **7**:975-989.
- 1028 Li, X.-M., D.-Y. She, D.-Y. Zhang, and W.-J. Liao. 2015. Life history trait differentiation and  
1029 local adaptation in invasive populations of *Ambrosia artemisiifolia* in China. *Oecologia*  
1030 **177**:669-677.
- 1031 Liu, H., and P. Stiling. 2006. Testing the enemy release hypothesis: a review and meta-analysis.  
1032 *Biological Invasions* **8**:1535-1545.
- 1033 Liu, Y. Z., P. B. Reich, G. Y. Li, and S. C. Sun. 2011. Shifting phenology and abundance under  
1034 experimental warming alters trophic relationships and plant reproductive capacity.  
1035 *Ecology* **92**:1201-1207.
- 1036 Lovelock, C. E., I. C. Feller, M. C. Ball, J. Ellis, and B. Sorrell. 2007. Testing the growth rate vs.  
1037 geochemical hypothesis for latitudinal variation in plant nutrients. *Ecology Letters*  
1038 **10**:1154-1163.
- 1039 Lozier, J. D., G. K. Roderick, and N. J. Mills. 2009. Tracing the invasion history of mealy plum  
1040 aphid, *Hyalopterus pruni* (Hemiptera: Aphididae), in North America: a population  
1041 genetics approach. *Biological Invasions* **11**:299-314.
- 1042 Marks, M., B. Lapin, and J. Randall. 1994. *Phragmites australis* (*P. communis*): threats,  
1043 management and monitoring. *Natural Areas Journal* **14**:285-294.
- 1044 Maron, J. L., M. Vilà, R. Bommarco, S. Elmendorf, and P. Beardsley. 2004. Rapid evolution of  
1045 an invasive plant. *Ecological Monographs* **74**:261-280.
- 1046 Mattson, W. J. 1980. Herbivory in relation to plant nitrogen content. *Annual Review of Ecology*  
1047 *and Systematics* **11**:119-161.
- 1048 Mazerolle, M. J. 2015. AICcmodavg: Model selection and multimodel inference based on  
1049 (Q)AIC(c). R package version 2.0-3. <http://CRAN.R-project.org/package=AICcmodavg>.
- 1050 McMaster, G. S., and W. W. Wilhelm. 1997. Growing degree-days: one equation, two  
1051 interpretations. *Agricultural and Forest Meteorology* **87**: 291-300.
- 1052 McNaughton, S. J. 1979. Grazing as an optimization process: grass-ungulate relationships in the  
1053 Serengeti. *American Naturalist* **113**:691-703.

- 1054 Meadows, R. E., and K. Saltonstall. 2007. Distribution of native and introduced *Phragmites*  
1055 *australis* in freshwater and oligohaline tidal marshes of the Delmarva peninsula and  
1056 southern New Jersey. *The Journal of the Torrey Botanical Society* **134**:99-107.
- 1057 Meyerson, L. A., A. M. Lambert, and K. Saltonstall. 2010. A tale of three lineages: expansion of  
1058 common reed (*Phragmites australis*) in the US Southwest and Gulf Coast. *Invasive Plant*  
1059 *Science and Management* **3**:515-520.
- 1060 Meyerson, L. A., and J. T. Cronin. 2013. Evidence for multiple introductions of *Phragmites*  
1061 *australis* to North America: detection of a new non-native haplotype. *Biological*  
1062 *Invasions* **15**:2605-2608.
- 1063 Meyerson, L. A., J. T. Cronin, and P. Pysek. 2016. *Phragmites australis* as a model organism for  
1064 studying plant invasions. *Biological Invasions* **18**:2421–2431.
- 1065 Mitchell, C. E., and A. G. Power. 2003. Release of invasive plants from fungal and viral  
1066 pathogens. *Nature* **421**:625-627.
- 1067 Moles, A. T., S. P. Bonser, A. G. B. Poore, I. R. Wallis, and W. J. Foley. 2011. Assessing the  
1068 evidence for latitudinal gradients in plant defence and herbivory. *Functional Ecology*  
1069 **25**:380-388.
- 1070 Nakagawa, S., and H. Schielzeth. 2013. A general and simple method for obtaining  $R^2$  from  
1071 generalized linear mixed-effects models. *Methods in Ecology and Evolution* **4**:133-142.
- 1072 Novy, A., S. L. Flory, and J. M. Hartman. 2013. Evidence for rapid evolution of phenology in an  
1073 invasive grass. *Journal of Evolutionary Biology* **26**:443-450.
- 1074 Orians, G. H., and A. V. Milewski. 2007. Ecology of Australia: the effects of nutrient-poor soils  
1075 and intense fires. *Biological Reviews* **82**:393-423.
- 1076 Paul, J., N. Vachon, C. J. Garroway, and J. R. Freeland. 2010. Molecular data provide strong  
1077 evidence of natural hybridization between native and introduced lineages of *Phragmites*  
1078 *australis* in North America. *Biological Invasions* **12**:2967-2973.
- 1079 Pennings, S. C., C.-K. Ho, C. S. Salgado, K. Więski, N. Davé, A. E. Kunza, and E. L. Wason.  
1080 2009. Latitudinal variation in herbivore pressure in Atlantic Coast salt marshes. *Ecology*  
1081 **90**:183-195.
- 1082 Pennings, S. C., E. L. Siska, and M. D. Bertness. 2001. Latitudinal differences in plant  
1083 palatability in Atlantic coast salt marshes. *Ecology* **82**:1344-1359.

1084 R Development Core Team. 2015. R: A language and environment for statistical computing. R  
1085 Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.  
1086 Rasband, W. S. 2014. ImageJ. U. S. National Institutes of Health, Bethesda, Maryland, USA,  
1087 <http://imagej.nih.gov/ij/>, 1997-2014.  
1088 Raupp, M. J. 1985. Effects of leaf toughness on mandibular wear of the leaf beetle *Plagioder*  
1089 *versicolora*. *Ecological Entomology* **10**:73-79.  
1090 Reich, P. B., and J. Oleksyn. 2004. Global patterns of plant leaf N and P in relation to  
1091 temperature and latitude. *Proceedings of the National Academy of Sciences of the United*  
1092 *States of America* **101**:11001-11006.  
1093 Richards, C. L., O. Bossdorf, N. Z. Muth, J. Gurevitch, and M. Pigliucci. 2006. Jack of all trades,  
1094 master of some? On the role of phenotypic plasticity in plant invasions. *Ecology Letters*  
1095 **9**: 981-993.  
1096 Rosenzweig, M. L. 1995. *Species diversity in space and time*. Cambridge University Press.  
1097 Salgado, C. S., and S. C. Pennings. 2005. Latitudinal variation in palatability of salt-marsh  
1098 plants: are differences constitutive? *Ecology* **86**:1571-1579.  
1099 Saltonstall, K. 2002. Cryptic invasion by a non-native genotype of the common reed, *Phragmites*  
1100 *australis*, into North America. *Proceedings of the National Academy of Sciences*  
1101 **99**:2445-2449.  
1102 Saltonstall, K. 2003. Microsatellite variation within and among North American lineages of  
1103 *Phragmites australis*. *Molecular Ecology* **12**:1689-1702.  
1104 Saltonstall, K., H. E. Castillo, and B. Blossey. 2014. Confirmed field hybridization of native and  
1105 introduced *Phragmites australis* (Poaceae) in North America. *American Journal of*  
1106 *Botany* **101**:211-215.  
1107 Santamaría, L., J. Figuerola, J. J. Pilon, M. Mjelde, A. J. Green, T. De Boer, R. A. King, and R.  
1108 J. Gornall. 2003. Plant performance across latitude: the role of plasticity and local  
1109 adaptation in an aquatic plant. *Ecology* **84**:2454-2461.  
1110 Schemske, D. W., G. G. Mittelbach, H. V. Cornell, J. M. Sobel, and K. Roy. 2009. Is there a  
1111 latitudinal gradient in the importance of biotic interactions? *Annual Review of Ecology*  
1112 *and Systematics* **40**:245-269.  
1113 Scriber, J. M., and P. Feeny. 1979. Growth of herbivorous caterpillars in relation to feeding  
1114 specialization and to the growth form of their food plants. *Ecology* **60**:829-850.



- 1115 Siska, E. L., S. C. Pennings, T. L. Buck, and M. D. Hanisak. 2002. Latitudinal variation in  
1116 palatability of salt-marsh plants: which traits are responsible? *Ecology* **83**:3369-3381.
- 1117 Slatkin, M. 1985. Gene flow in natural populations. *Annual Review of Ecology and Systematics*  
1118 **16**:393-430.
- 1119 Smith, L. M. 1936. Biology of the mealy plum aphid, *Hyalopterus pruni* (Geoffroy). *Hilgardia*  
1120 **10**:167-211.
- 1121 Sparks, A. N. 1979. A review of the biology of fall armworm. *Florida Entomologist* **62**:82-87.
- 1122 Strauss, S. Y. and A. A. Agrawal. 1999. The ecology and evolution of plant tolerance to  
1123 herbivory. *Trends in Ecology and Evolution* **14**:179-185.
- 1124 Tewksbury, L., R. Casagrande, B. Blossey, P. Häfliger, and M. Schwarzländer. 2002. Potential  
1125 for Biological Control of *Phragmites australis* in North America. *Biological control*  
1126 **23**:191-212.
- 1127 Tscharrntke, T. and Greiler 1995. Insect communities, grasses and grasslands. *Annual Review of*  
1128 *Entomology* **40**:535-558.
- 1129 Waterman, P. G., and S. Mole. 1994. Analysis of plant metabolites. *Methods in ecology series*.  
1130 Blackwell Scientific Publications, Oxford, United Kingdom.
- 1131 Woods, E. C., A. P. Hastings, N. E. Turley, S. B. Heard, and A. A. Agrawal. 2012. Adaptive  
1132 geographical clines in the growth and defense of a native plant. *Ecological Monographs*  
1133 **82**:149-168.
- 1134 Zheng, Y. L., Y. L. Feng, R. F. Wang, X. D. Shi, Y. B. Lei, and L. H. Han. 2012. Invasive  
1135 *Eupatorium adenophorum* suffers lower enemy impact on carbon assimilation than native  
1136 congeners. *Ecological Research* **27**:867-872.
- 1137 Zhou, W., Z. Wang, W. Xing, and G. Liu. 2014. Plasticity in latitudinal patterns of leaf N and P  
1138 of *Oryza rufipogon* in China. *Plant Biology* **16**:917-923.

#### SUPPORTING INFORMATION

1141 Supporting information may be found online at: <http://onlinelibrary.wiley.com/doi/xx>

#### DATA AVAILABILITY

1144 Data associated with this paper have been deposited in Dryad:  
1145 <http://dx.doi.org/10.5061/dryad.r8d1m>

1146 **Table 1.** Best models (based on AICc model-selection procedure for the effects of garden (G),  
 1147 lineage (S), and latitude (L) and all possible two- and three-way interactions on each *P. australis*  
 1148 plant defense, nutrition and palatability traits. Detailed information about these models is  
 1149 provided in Appendix S3: Table S1.

Dependent variable	Model
<i>Plant defense traits</i>	
Leaf toughness	G L G×L
Phenolics	G S L G×S G×L S×L G×S×L
<i>Plant palatability traits</i>	
Water content	G S L G×S G×L S×L G×S×L
% nitrogen	G S L G×S G×L S×L G×S×L
% carbon	G L G×L
C:N	G S L G×S G×L S×L G×S×L
Aphid colony size*	G S L G×S
Aphid survivorship**	G S L G×S
Larval growth	G S
Larval survivorship**	G S
Leaf area chewed	G S
Larval biomass conversion efficiency	G S L G×S G×L

1150 *Notes:* Nutrient concentrations (% carbon, % nitrogen, C:N, and phenolics) and leaf area chewed  
 1151 (cm<sup>2</sup>) were *ln* transformed.

1152 \*Generalized linear mixed effect model (GLMM) with Poisson family was used in the analysis.

1153 \*\*General linear models were used in these analyses.

1154

1155

#### FIGURE LEGENDS

1156 **Fig. 1.** Map of the location of *P. australis* source genotypes used in this study and the location of  
1157 the common gardens at Louisiana State University (LSU) and University of Rhode Island (URI).

1158 **Fig. 2.** Effects of lineage, latitude and garden on defense and nutritional traits of native and  
1159 invasive *P. australis* grown in common gardens at Louisiana State University and the University  
1160 of Rhode Island. Symbols in the shaded portion of the graph are the least-squares means ( $\pm$  SE)  
1161 for different lineages in different gardens. The relationship between a plant trait and latitude is  
1162 shown by a line fit by least-squares regression, only for cases in which the AICc-best model  
1163 included the effect of latitude (either a Latitude or Lineage  $\times$  Latitude or Garden  $\times$  Latitude).  
1164 Thick lines denote significant trait-latitude relationships ( $P \leq 0.05$ ) and thin lines represent non-  
1165 significant relationships ( $P > 0.05$ ; see Appendix S3). Slopes ( $\beta_{LSU}$  and  $\beta_{URI}$ ) are reported for the  
1166 significant relationships.


1167 **Fig. 3.** Effects of lineage, latitude and garden on palatability traits of native and invasive *P.*  
1168 *australis* grown in common gardens at Louisiana State University and the University of Rhode  
1169 Island. Symbols in the shaded portion of each graph are least-squares means ( $\pm$  SE) for native  
1170 and invasive lineages in each garden. In cases where Latitude or Lineage  $\times$  Latitude or Garden  $\times$   
1171 Latitude was a factor in the AICc-best model, the relationship between the plant trait and latitude  
1172 is shown by a line fit by least-squares regression. Thick lines denote significant trait-latitude  
1173 relationships ( $P \leq 0.05$ ) and thin lines represent non-significant relationships ( $P > 0.05$ ; see  
1174 Appendix S3). Slopes ( $\beta_{LSU}$  and  $\beta_{URI}$ ) are reported for the significant relationships.

1175 **Fig. 4.** Phenotypic plasticity in defense and palatability traits for native and invasive genotypes  
1176 of *P. australis*. Plasticity in plant traits ( $n = 10$ ) was estimated as the proportional difference in  
1177 mean trait expression between gardens (i.e., [mean trait value for a lineage at LSU – mean trait  
1178 value for that lineage at URI]/mean trait value for LSU). Symbols above the 1:1 line depict  
1179 greater plasticity in invasive genotypes than the natives. Inset bar graph shows least squares  
1180 mean  $\pm$  SE plasticity for native (Nat) and invasive (Inv) genotypes. A statistically significant  
1181 difference between means (pairwise *t*-test;  $P = 0.024$ ) is represented by an asterisk.

1182 **Fig. 5.** The relationship between trait plasticity (proportional difference in mean trait expression  
1183 between gardens) and latitude for native and invasive *P. australis* genotypes. Each point

1184 represents a single genotype that was represented in each garden. Lines are fit by least-squares  
1185 regression and were provided only for those traits in which a significant latitude or lineage  $\times$   
1186 latitude interaction was detected in an ANCOVA (see Appendix S6). Slopes ( $\beta$ ) are reported for  
1187 the significant relationships.

Author Manuscript




This article is protected

Figure 1

● Native LSU    ○ Native URI    ■ Invasive LSU    □ Invasive URI


ecm\_1233\_f2.pdf



This article is protected by copyright. All rights reserved


Figure 2

Native LSU Native URI Invasive LSU Invasive URI



This article is protected by copyright. All rights reserved


Figure 3



This article is protected

Figure 4





This article is protected by copyright. All rights reserved

Figure 5