

2-7-2000

A density functional study of open-shell cyclopentadienyl-molybdenum(II) complexes. A comparison of stabilizing factors: Spin-pairing, Mo-X π bonding, and release of steric pressure

Ivo Cacelli
Scuola Normale Superiore di Pisa

Rinaldo Poli
Universite de Bourgogne

Elsje Alessandra Quadrelli
Ist. Chim. Quantistica ed E.

Antonio Rizzo
Ist. Chim. Quantistica ed E.

Kevin M. Smith
Universite de Bourgogne

Follow this and additional works at: https://repository.lsu.edu/chemistry_pubs

Recommended Citation

Cacelli, I., Poli, R., Quadrelli, E., Rizzo, A., & Smith, K. (2000). A density functional study of open-shell cyclopentadienyl-molybdenum(II) complexes. A comparison of stabilizing factors: Spin-pairing, Mo-X π bonding, and release of steric pressure. *Inorganic Chemistry*, 39 (3), 517-524. <https://doi.org/10.1021/ic990875i>

This Article is brought to you for free and open access by the Department of Chemistry at LSU Scholarly Repository. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Scholarly Repository. For more information, please contact ir@lsu.edu.



HAL
open science

A Density Functional Study of Open-Shell Cyclopentadienyl–Molybdenum(II) Complexes. A Comparison of Stabilizing Factors: Spin-Pairing, Mo–X π Bonding, and Release of Steric Pressure

Ivo Cacelli, Rinaldo Poli, Elsje Alessandra Quadrelli, Antonio Rizzo, Kevin
Smith

► To cite this version:

Ivo Cacelli, Rinaldo Poli, Elsje Alessandra Quadrelli, Antonio Rizzo, Kevin Smith. A Density Functional Study of Open-Shell Cyclopentadienyl–Molybdenum(II) Complexes. A Comparison of Stabilizing Factors: Spin-Pairing, Mo–X π Bonding, and Release of Steric Pressure. *Inorganic Chemistry*, American Chemical Society, 2000, 39 (3), pp.517-524. 10.1021/ic990875i . hal-03296084

HAL Id: hal-03296084

<https://hal.archives-ouvertes.fr/hal-03296084>

Submitted on 29 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A Density Functional Study of Open-shell Cyclopentadienyl-molybdenum(II) Complexes. A Comparison of Stabilizing Factors: Spin-pairing, Mo-X π Bonding, and Release of Steric Pressure

Ivo Cacelli,^a Rinaldo Poli,*^b Elsje Alessandra Quadrelli,^c Antonio Rizzo,^c and Kevin M. Smith^b

^aScuola Normale Superiore, Piazza dei Cavalieri 7, I-56126 Pisa, Italy.

^bLaboratoire de Synthèse et d'Electrosynthèse Organométalliques, Faculté des Sciences "Gabriel", Université de Bourgogne, 6 Boulevard Gabriel, F-21000 Dijon, France

^cIstituto di Chimica Quantistica ed Energetica Molecolare del Consiglio Nazionale delle Ricerche, Via Risorgimento 35, I-56126 Pisa, Italy.

Abstract

The dissociation of PH_3 from the 18-electron system $\text{CpMoX}(\text{PH}_3)_3$ to afford the corresponding 16-electron $\text{CpMoX}(\text{PH}_3)_2$ fragment has been investigated theoretically by DFT for $\text{X} = \text{H}, \text{CH}_3, \text{F}, \text{Cl}, \text{Br}, \text{I}, \text{OH},$ and PH_2 . The product is found to prefer a triplet spin state for all X ligands except PH_2 , the singlet-triplet gap varying between 1.7 kcal/mol for OH to 8.7 kcal/mol for F . The Mo-PH_3 bond dissociation energy to the 16-electron ground state varies dramatically across the series, from 4.5 kcal/mol for OH to 23.5 kcal/mol for H and correlates with experimental observations on trisubstituted phosphine derivatives. Geometry-optimized spin doublet $\text{CpMo}(\text{PH}_3)_3$, on the other hand, has a Mo-PH_3 BDE of 24.3 kcal/mol. The modulation of the Mo-PH_3 BDE by the introduction of X is analyzed in terms of three effects which stabilize the 16-electron product relative to the 18-electron starting complex: (i) adoption of the higher (triplet) spin state by release of pairing energy; (ii) Mo-X π interactions; (iii) release of steric pressure. A computational model for the approximate separation and evaluation of these three stabilizing effects is presented. According to the results of these calculations, the relative importance of the three effects depends on various factors related to the nature of X . For double-sided π donor X ligands, the larger triplet-singlet gap is provided by the more electronegative atoms ($\text{F} > \text{Cl} > \text{Br} > \text{I}$), while single-sided π donors favor the singlet state. The π stabilization ability goes in the order $\text{PH}_2 > \text{OH} > \text{F} > \text{other halogens} > \text{H}$. Finally, the major steric interaction appears to be associated with the presence of inactive lone-pairs and by their orientation/proximity to the PH_3 ligands ($\text{Cl}, \text{Br} > \text{I}, \text{OH} > \text{F}, \text{PH}_2, \text{H}, \text{CH}_3$). The 16-electron methyl system establishes a marked α -agostic interaction in the singlet state, which remains nevertheless unfavored relative to an undistorted triplet configuration.

Introduction

The chemistry of organometallic compounds is dominated by the 18-electron rule and by diamagnetism.¹ This is the direct consequence of three factors: the high bond covalency in this realm of chemistry, the ability of the ligands to establish π back-bonding interactions with the central metallic element, and the relatively low electron-electron repulsion enabling the establishment of spin-paired configurations. Electronically unsaturated (open-shell) configurations are, however, frequently associated with reaction intermediates (this being the basis of Tolman's "16 and 18-electron rule"),² and can lead to the isolation of stable compounds under favorable circumstances. A clear understanding of the factors at work in the stabilization of open-shell structures relative to saturated counterparts is therefore fundamental for the rationalization of reaction rates and catalytic activity.

Complexes having a 16-electron configuration may be accessed from closed-shell precursors in a number of ways, *e.g.* ligand dissociation, reductive elimination, migratory insertion, and so on. Four mechanisms may be distinguished for their energetic stabilization relative to the saturated precursor: (i) release of steric pressure associated with the decrease of interligand repulsive van der Waals interactions;³ (ii) intervention of ligand lone pairs (π -donation);^{4,5} (iii) release of pairing energy (this playing a role only when the open-shell and saturated species have different spin states);⁶ (iv) interactions, including agostic ones, with other donor molecules (*e.g.* the solvent) or groups (*e.g.* dangling donor functions from ligands). The fourth mechanism effectively consists of the temporary saturation of the open coordination site (replacement of a ligand with a more labile one), rather than the relative stabilization of an unsaturated structure. It is responsible, for instance, for the acceleration of many reactions involving a rate-determining dissociative step when these are carried out in donor solvents. The present contribution focuses only on the analysis of the first three effects, although the intervention of intramolecular α -agostic interactions will be highlighted in a particular case.

The relative importance of each of these effects cannot be easily determined or estimated. We will be concerned here with the formation of half-sandwich 16-electron Mo(II) complexes by ligand

dissociation from 18-electron precursors, but the arguments may be extrapolated to any other reaction generating a 16-electron from an 18-electron compound or any $(n-2)$ -electron from an n -electron compound. We shall take $\text{CpMoX}(\text{PH}_3)_2$ as a model for the bis- PMe_3 complexes previously used by us in a number of studies.⁷⁻¹¹ The relevance of a sterics-related stabilization is experimentally indicated by the equilibrium between $\text{Cp}^*\text{MoCl}(\text{PMe}_3)_3$ and the mixture of $\text{Cp}^*\text{MoCl}(\text{PMe}_3)_2$ and PMe_3 at room temperature, or between $\text{CpMoCl}(\text{PMe}_2\text{Ph})_3$ and the mixture of $\text{CpMoCl}(\text{PMe}_2\text{Ph})_2$ and PMe_2Ph upon warming, whereas no phosphine loss was observed for $\text{CpMoCl}(\text{PMe}_3)_3$ under the same conditions.⁸ The absence of any $\text{CpMo}(\text{OH})(\text{PMe}_3)_3$ in equilibrium with a mixture of $\text{CpMo}(\text{OH})(\text{PMe}_3)_2$ and PMe_3 ,^{10,11} on the other hand, demonstrates the intervention of additional stabilizing factors (OH occupies less space than Cl). Both Cl and OH have lone pairs and are thus capable of providing π stabilization. It is generally believed that a hydroxo ligand is a stronger π donor than the halogens but quantitative assessment, to our knowledge, are not available. On the other hand, since the stable 16-electron systems $\text{Cp}^*\text{MoCl}(\text{PMe}_3)_2$ and $\text{CpMo}(\text{OH})(\text{PMe}_3)_2$ exhibit a spin triplet ground state, a stabilizing factor associated to the release of pairing energy must also play a role.

The quantitative evaluation of π bond strengths is a difficult exercise.⁴ It requires the separation of σ and π bonding components, which is experimentally impossible and theoretically nonrigorous.¹²⁻¹⁴ From the experimental point of view, qualitative evaluations of trends in M-X π bond strengths, these relating in most cases to bonds between a metal and the halogens, have been deduced from the analysis of NMR chemical shifts,¹⁵ EPR g values,¹⁶ IR carbonyl stretching frequencies,^{17,18} electrochemical data,^{19,20} UV-visible spectra,^{21,22} valence photoelectron spectra,²³ X-ray crystal structures,^{24,25} and rates of chemical and fluxional processes or even the simple observation that certain reactions take place whereas others do not.²⁶⁻²⁹ The use of such a wide array of techniques, each necessitating a different set of assumptions and approximations, has led to controversy. For instance, no universal agreement seems to have been achieved as to the whether a lighter donor atom is a weaker or a stronger π donor than a heavier congener (e.g. Cl vs. I or SR vs. TeR); the relative donor power may well depend on the nature of the metal as well as on the chemical environment.

From the point of view of theory, different localization and partition schemes have been employed^{14,30-40} but, to the best of our knowledge, these have not been applied to characterize the nature of the metal to ligand bonding in transition metal complexes of the kind studied here.

In a recent contribution,¹¹ we have presented a new computational model that has allowed us, within certain approximations, to tell apart the contributions of interligand repulsion, π donation, and electron pairing to the relative stabilization of the CpMoX(PH₃)₂ model systems. We have now refined this model and extended it to a wider range of X ligands. The results reported herein allow, among other things, a rough quantitative estimation of the Mo-X π bond strength.

Computational Details

All electronic structure and geometry optimization calculations were performed using Gaussian-94⁴¹ on an SGI Origin 200 workstation at the Université de Bourgogne and on both a Alpha Digital and a SGI Indigo 2 in Pisa. The three-parameter form of the Becke, Lee, Yang and Parr functional (B3LYP),⁴² was employed. The LanL2DZ basis set includes both Dunning and Hay's D95 sets for H, C, O and F⁴³ and the relativistic Electron Core Potential (ECP) sets of Hay and Wadt for the 10 inner electrons of Cl and P, the 18 inner electrons of Br, and the 28 inner electron of Mo and I.⁴⁴⁻⁴⁶ Unless otherwise stated, calculations were carried out without spatial symmetry constraints. The energies reported for the open shell systems correspond to unrestricted (UB3LYP) calculations. In our⁴⁷⁻⁴⁹ and other's⁵⁰⁻⁵² experience, this computational method correctly reproduces the experimentally observed spin state for open-shell systems. For open shell systems the mean value of the spin over the electronic density in an unrestricted calculations does not reproduce exactly the assigned spin multiplicity. In all our cases, though, it was considered to be suitable to identify unambiguously the spin state. Mean values of $\langle S^2 \rangle$ were in the narrow 2.012-2.015 range for triplets.

Results and Discussion

a. Geometries and Mo-PH₃ bond dissociation energies. Singlet vs. triplet CpMoX(PH₃). The considerations presented in this article are based on the energetics of the dissociation of one phosphine ligand on the model system CpMoX(PH₃)₃ (X = H, CH₃, F, Cl, Br, I, OH, PH₂), see Scheme 1. The Mo-PH₃ bond dissociation energy (BDE) relates the 18-electron CpMoX(PH₃)₃ complex and the combination of 16-electron CpMoX(PH₃)₂ and free PH₃, all geometry optimized. The BDE's relative to both singlet and triplet 16-electron products are collected in Table 1. While the 18-electron complex adopts a singlet ground state, the 16-electron complex has a lowest lying triplet configuration in all cases except for the phosphido derivative.

Scheme 1




Table 1. Relevant energetic parameters for the PH₃ dissociation from CpMoX(PH₃)₃ (see Schemes 1 and 2). All energies are in kcal/mol.

| X | BDE (Mo-PH ₃) | BDE (Mo-PH ₃) | ΔE_{S-T}^a | ΔE_{π}^b | ΔE_{π}^b | ΔE_{steric}^c | ΔE_{steric}^d |
|-----------------|---------------------------|---------------------------|--------------------|--------------------|--------------------|-----------------------|-----------------------|
| | (triplet) | (singlet) | | (singlet) | (triplet) | | (singlet) |
| H | 23.5 | 25.3 | 1.8 | -2.0 | 1.3 | 1.0 | -0.6 |
| Me | 13.2 | 16.3 | 3.1 | 7.0 | 11.7 | 1.0 | 0.3 |
| F | 6.3 | 14.9 | 8.7 | 7.7 | 19.3 | 1.7 | -0.2 |
| Cl | 5.9 | 13.4 | 7.5 | 2.8 | 14.3 | 8.1 | 5.2 |
| Br | 6.5 | 13.6 | 7.0 | 1.8 | 13.1 | 9.0 | 5.7 |
| I | 9.1 | 15.7 | 6.6 | 4.2 | 15.3 | 4.5 | 1.0 |
| OH | 4.5 | 6.2 | 1.7 | 14.1 | 18.8 | 4.1 | 2.1 |
| PH ₂ | 14.9 | 7.0 | -7.9 | 16.0 | 10.2 | 1.3 | 0.3 |

^a $\Delta E_{S-T} = E_{singlet} - E_{triplet}$. ^b $\Delta E_{\pi} = BET(Mo-X \sigma+\pi) - BET(Mo-X \sigma)$. ^c $\Delta E_{steric}(singlet) = -\Delta E_1 + \Delta E_2$. ^d $\Delta E_{steric}(triplet) = -\Delta E_1 + \Delta E_3$.

We have calculated the basis set superposition error (BSSE)⁵³ for the fluoride system using the counterpoise technique,⁵⁴ and found a decrease of 2.9 kcal/mol for the Mo-PH₃ BDE. Since the BSSE depends largely on the bond distance and since the Mo-PH₃ bond length is essentially independent of X, we presume that all BDE's reported in Table 1 should be scaled down by the same amount. A reduction of all BDE's may in fact lead to a better agreement with available experimental data, as the lowest BDE for the OH system may be compared with the stability of triplet CpMo(OH)(PMe₃)₂ in the presence of excess PMe₃.^{10,11}

The geometry optimized 18-electron and the triplet 16-electron systems correspond to essentially C_s-symmetric structures which can be described, respectively, as four- and three-legged piano stools (see **I** and **II**). The singlet 16-electron systems, on the other hand, have optimized geometries exhibiting a certain degree of asymmetry. Two different minima were located for each X, the lowest one corresponding in all cases except for X = H to a C₁ geometry where the ligand X has more or less deviated from the P-Mo-P bisector plane, as schematically illustrated in **III**. The second minimum (lowest for X = H) is an essentially C_s-symmetric structure where the P-Mo-P angle has more or less increased relative to the triplet **II**. The distortion from a regular (pseudo-octahedral) structure as found in the triplet can be assessed by two dihedral angles. The angle θ is the angle formed by the CNT-Mo-X plane and the plane passing through CNT, Mo, and the bisector of the two Mo-PH₃ bonds, whereas the angle ϕ is the dihedral P-Mo-CNT-P angle (CNT is the Cp ring centroid), see **III**. The values of the angles θ and ϕ for the lowest energy optimized CpMoX(PH₃)_n (n = 2, 3) are collected in Table 2 together with the Mo-X distances. In most cases, the Mo-CNT and Mo-PH₃ distances for **I** have intermediate values between those of **II** (longer) and **III** (shorter), which differ from each other by less than 3%.




Table 2. Selected optimized bond angles (°) and lengths (Å) for CpMoX(PH₃)_n (n = 3, 2)

| | X | H ^a | CH ₃ | F | Cl | Br | I | OH | PH ₂ |
|-------------------------------|----------------------|----------------|-----------------|-------|-------|-------|-------|-------|-----------------|
| n = 3 | Mo-X | 1.722 | 2.299 | 2.085 | 2.633 | 2.805 | 3.002 | 2.095 | 2.696 |
| n = 2 II (Triplet) | Mo-X | 1.739 | 2.193 | 1.993 | 2.472 | 2.627 | 2.806 | 2.012 | 2.565 |
| | ξ(Mo-X) ^b | -0.99 | 4.6 | 4.4 | 6.1 | 6.3 | 6.5 | 4.0 | 4.9 |
| | θ ^c | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 6.7 |
| | φ ^c | 159.0 | 130.7 | 123.4 | 120.0 | 119.0 | 117.2 | 123.6 | 129.0 |
| n = 2 III (Singlet) | Mo-X | 1.723 | 2.126 | 1.994 | 2.463 | 2.611 | 2.781 | 1.972 | 2.326 |
| | ξ(Mo-X) ^b | -0.06 | 7.5 | 4.4 | 6.5 | 6.9 | 7.4 | 5.9 | 13.7 |
| | θ ^c | 43.9 | 26.0 | 25.5 | 13.1 | 6.4 | 0.3 | 10.5 | 1.2 |
| | φ ^c | 105.7 | 102.0 | 102.2 | 102.4 | 102.3 | 101.9 | 102.7 | 102.6 |

^aThe lowest energy n=2 (singlet) structure (see Figure 1a) has Mo-H = 1.700 Å, ξ(Mo-X) = 1.2, θ = 1.4° and φ = 188.7°. ^b ξ(Mo-X) = {[Mo-X (n=3) - Mo-X (n=2)]/[Mo-X (n=3)]} · 100. ^cSee drawing **III**.

The global singlet minimum for $X = H$ is shown in Figure 1a. One can imagine this structure as derived from structure **I** by removal of the PH_3 ligand trans to the H atom, whereas structure **III** may be imagined as derived by removal of a cis PH_3 ligand. All X ligands used in this study, except for H, have a weaker trans influence than PH_3 , thus the Mo-PH₃ bond trans to X is stronger than the Mo-PH₃ bond trans to PH_3 . A special word must be said about the methyl system, for which the 16-electron global singlet minimum corresponds to a distorted geometry similar to **III** with an additional agostic interaction involving one of the methyl group H atoms, see Figure 1b. The agostic Mo...H distance in this optimized structure is 2.279 \AA and the Mo-C-H angle is 82.7° . The agostic C-H bond is slightly longer (1.137 \AA) than the other two C-H bonds (1.098 and 1.100 \AA). The preference for this agostic structure is well in agreement with the C-H oxidative addition reactivity established for complexes $\text{CpMo}(\text{CH}_3)(\text{PMe}_3)_3$ and for the Cp^* analogue.⁵⁵ An estimate of the strength of this agostic interaction was obtained by imposing identical Mo-C-H angles for the Mo-CH₃ moiety during the optimization. This structure was found to be 3.7 kcal/mol less stable than that of Figure 1b.




Figure 1. MacMoMo⁷⁰ view of the optimized structures of singlet $\text{CpMoH}(\text{PH}_3)_2$ (a) and $\text{CpMo}(\text{CH}_3)(\text{PH}_3)_2$ (b). The cyclopentadienyl H atoms are omitted for clarity.

Geometrical distortions similar to those observed here for singlet $\text{CpMoX}(\text{PH}_3)_2$ have previously been analyzed computationally for related electronically unsaturated systems, for instance 16-electron d^6 and d^4 CpML_n systems.^{25,56-58} In order to explain the geometries observed for the singlet d^4 $\text{CpMoX}(\text{PH}_3)_2$ systems, the Kohn and Sham orbitals of the highly

distorted hydride species were examined in detail. The use of these orbitals for both qualitative and quantitative assessments of chemical phenomena is well established.⁵⁹⁻⁶¹ The orbital energies and shapes of the frontier orbitals of singlet CpMoH(PH₃)₂ in various geometries are represented pictorially in Figure 2. In addition to the distorted optimized geometries (the higher energy local minimum, labeled **III**, and the global minimum **IV**), the orbitals of the undistorted, pseudo-octahedral (electronically speaking, when considering the Cp as occupying three mutually adjacent coordination positions) species were obtained via a single point calculation on an idealized three-legged piano stool geometry, labeled "**O_h**" in Figure 2.




Figure 2. Frontier orbital evolution upon distortion of singlet CpMoH(PH₃)₂ from an idealized three-legged piano stool geometry.

The "**O_h**" geometry has PH₃ ligands that lie partially along the lobes of the HOMO. In both optimized structures **III** and **IV**, the distortion moves the PH₃ ligands onto the nodal planes of this orbital, thereby lowering it in energy so that it becomes the SHOMO. The other doubly-occupied orbital, the HOMO for **III** and **IV** and the SHOMO for "**O_h**", has a nodal surface along the Mo-PH₃ and Mo-H bonds in all three cases, and so it remains essentially constant in energy. The same geometry change that stabilizes one of the filled orbitals causes the PH₃ groups to move onto the lobes of the LUMO, leading to the destabilization and rehybridization of this orbital in **III** and **IV**. This effect is accompanied by the stabilization of the corresponding metal-ligand bonding combination, which provides an additional, perhaps more important, stabilizing effect. The net result is that both distortions induce a significant stabilization of two filled orbitals and the increase of the HOMO-LUMO gap, thereby improving the relative stability of the spin-paired complexes.⁶² This orbital splitting is not desirable for the triplet spin state and triplet CpMoH(PH₃)₂ does not exhibit these deformations. Parallels may be drawn to the studies of Eisenstein and co-workers, who have demonstrated that although triplet d⁶ ML₅ compounds have undistorted trigonal bipyramidal structures,⁶³ two different deformation modes are observed for the singlet species, the relative energy of which are dependent on the π - and σ -bonding properties of the ancillary ligands.^{64,65}

Although the geometric distortions are most pronounced for X = H and Me (see Table 2), large effects are also displayed for the F, Cl and OH singlet complexes and the distortions are progressively smaller for Br, PH₂ and I systems. This variation with respect to the identity of the X ligand is presumably the result of competing σ - and π -effects. For the hydride and methyl, no significant π -donation interactions are possible, and so geometric distortions are the only available option to decrease the total energy. For X = π -donor ligand, the orbital energies are also influenced by π -donation,^{8,11} which is equally effective in a non-distorted geometry. Thus, although F and OH have stronger π donating properties (*vide infra*), they also provide a stronger σ component leading to significant distortions.


All optimized parameters compare rather well with those experimentally observed for available related systems. The calculations correctly reproduce the observed ground state for all those 16-electron systems that have been isolated or observed in solution: $\text{Cp}^*\text{Mo}(\text{PR}_2)(\text{PMe}_3)_2$ is diamagnetic,⁶⁶ while $\text{Cp}^*\text{MoClL}_2$ ($L = \text{PMe}_3, \text{PMe}_2\text{Ph}$ or $L_2 = \text{dppe}$)^{7,8} and $\text{CpMo}(\text{OH})(\text{PMe}_3)_2$ ^{10,11} have two unpaired electrons. The calculated Mo-PH₃ BDE's (Table 1) are also consistent with the observed relative stability of the various 18-electron systems: while no PMe_3 dissociation from $\text{CpMoH}(\text{PMe}_3)_3$ occurs under ambient conditions,¹¹ the same dissociation from $\text{CpMo}(\text{CH}_3)(\text{PMe}_3)_3$ and $\text{Cp}^*\text{Mo}(\text{CH}_3)(\text{PMe}_3)_3$ readily takes place at 40°C and room temperature, respectively, leading to metallation of the PMe_3 and Cp^* ligand, respectively.⁵⁵ In addition, $\text{Cp}^*\text{MoCl}(\text{PMe}_3)_3$ establishes an observable equilibrium with $\text{Cp}^*\text{MoCl}(\text{PMe}_3)_2$ and PMe_3 ,⁸ and finally $\text{CpMo}(\text{OH})(\text{PMe}_3)_3$ does not exist. Steric effects are certainly involved in further stabilizing the unsaturated structure with respect to the computed PH₃ model system, see below. In addition, the use of the PH₃ model usually leads to weaker Mo-P bonds relative to PMe_3 .^{67,68}

b. Extent of the Mo-X π stabilization. As qualitatively discussed before,⁷ an X ligand with two lone pairs (double-sided π donor) should interact with a 16-electron metal fragment by establishing one two-center two-electron π bond with the empty metal orbital in the singlet state and two two-center three-electron π interactions with the two singly-occupied metal orbitals in the triplet state. Thus, for both spin states, a double-sided π donor can transfer two π electrons overall to the metal center. In addition, a four-electron destabilizing interaction is present in both spin states. On the other hand, a single-sided π donor can donate both π electrons only in the singlet structure, while a one electron π -stabilization (one half bond order) occurs for the triplet structure. Finally, π neutral X ligands should establish only a pure σ interaction in both singlet and triplet 16-electron systems like in the 18-electron system.

Breaking the Mo-X bond in **I** gives the radical pair $\text{CpMo}(\text{PH}_3)_3^\bullet$ and X^\bullet , both in the doublet state (Scheme 2). Energies for these systems are calculated in the frozen geometry derived

from optimized CpMoX(PH₃)₃, providing the so-called Bond Energy Terms (BET)⁶⁹ which are a measure of the Mo-X σ bond strengths. The analogous process on **II** and **III** gives a BET which may be related to the strength of the ($\sigma+\pi$) interaction. The subtraction of the Mo-X BET (18-electron system) from the Mo-X BET (16-electron system) gives us an evaluation of the stabilization provided to the unsaturated system by the establishment of the π interaction (ΔE_{π}). The results obtained are listed in Table 1. These values are essentially unaffected by the BSSE, as the individual BET's are affected by BSSE by essentially the same amount. For instance, a counterpoise correction for the fluoride system has given BET decreases of 3.0, 2.6 and 2.6 kcal/mol for **I**, **II** and **III**, respectively, leading to correction of less than 0.5 kcal/mol for ΔE_{π} .

Scheme 2




It is important to realize that the calculated ΔE_π values for the triplet state are skewed by the influence of X on the electron pairing energies. In fact, removal of $\text{X}\cdot$ from triplet $\text{CpMoX(PH}_3)_2$ correlates with the spin quartet state, as shown in Scheme 3 (left). Therefore, there is a larger exchange interaction in this case relative to singlet system (Scheme 3, right). This is the reason for the much larger ΔE_π values calculated for the triplet systems (Table 1). Even without pairing energy problems (i.e. for the singlet state), the deduction of a “ π -bond strength” from the ΔE_π values requires a number of approximations: (i) considering the σ bond as having the same strength in **I** and **III**; (ii) neglecting the rehybridization of X; (iii) neglecting the rehybridization of M; (iv) neglecting variations in the other M-L bonds. A π interaction

compresses the Mo-X bond (lengths and shortening factors are shown in Table 2), weakening the σ component of the bond, although the bond becomes stronger overall. This approximation leads to an underestimation of ΔE_π . Single point calculations for **I** at the Mo-X distance of **III** show that this underestimation is 0.0 for H (as expected for a π neutral ligand) and up to 3.4 kcal/mol for the halides. The effect of the rehybridization of X is important only for PH₂, again leading to an underestimation of the π bond strength for this ligand. The effect of a metal rehybridization is clearly illustrated by the structural distortions on going from **I** to **II** and **III**. The other Mo-ligand bonds do not appear to vary greatly, but how these small variations reflect into energy changes cannot be evaluated easily. In the absence of all these effects, the ΔE_π value for the π neutral H ligand should be exactly zero. The small negative value calculated for X = H (see Table 1) is an estimate of the limitations of these approximations. It should also be kept in mind that there is a nonzero (destabilizing) π interaction between the X and the metal lone pairs in CpMoX(PH₃)₃. The calculated ΔE_π values actually reflect the combination of this *filled-filled* repulsion⁴ at the 18-electron level and the π stabilization at the 16-electron level.

The ΔE_π results in Table 1 lend themselves to several considerations. Although a Mo-X π -bond strength for the triplet state cannot be deduced from the data, the trend of the ΔE_π values are approximately the same for the singlet and triplet species. In addition, while ΔE_π (triplet) is greater than ΔE_π (singlet) for all double-sided π donors, it is smaller for the phosphido ligand, confirming the qualitative considerations made above. This is probably the main reason for the adoption of a singlet ground state by the phosphido complexes. The action of the X ligands as π donors in the triplet complexes can also be judged by the percent of Mo-X bond shortening (ξ , see Table 2), which is rather similar for triplet and singlet complexes containing double-sided π -donor ligands (*i.e.* the halogens). The PH₂ ligand, being able to provide only 1 π electron in the triplet state, leads to a significant increase of ξ on going from triplet **II** to singlet **III**. The OH case is interesting, ξ increasing by 50% on going from triplet to singlet. While the OH ligand acts as a double-sided π donor,¹¹ one orbital interaction is stronger than the other

one leading to a weaker interaction in the triplet state. As a result, $\xi(\text{F}) > \xi(\text{OH})$ for triplet **II** whereas the reverse holds true for singlet **III**.

Scheme 3



The ΔE_π value calculated for F is significantly greater than zero, in agreement with the notion that the fluoride ion is a good π donor ligand. The values calculated for the other halides are smaller but positive. Within the halogen group, however, the trend is not monotonic, iodide leading to a greater value than both chloride and bromide. Given the approximations alluded to above, we do not feel that too much emphasis should be placed on this trend. However, the results are in agreement with the halogens having a small but significant π donating capability. The methyl ligand shows a 7.5% contraction for the agostic structure of Figure 1b. This shortening accompanies the high ΔE_π value which is, obviously, a measure of the strength of the Mo-CH₃ agostic interaction rather than a π interaction. Finally, the much larger values calculated for OH and PH₂ agree well with the known strong π -donor power of hydroxide, alkoxides, aryloxides, and dialkyl- and diarylphosphido ligands.

Somewhat surprisingly, the percent of Mo-X bond shortening $\xi(\text{Mo-X})$ in Table 2 *does not* correlate with the ΔE_π values. The smallest change is seen, as expected, for the π neutral H ligand. For the halogen series, the percent contraction increases in the order $\text{F} < \text{Cl} < \text{Br} < \text{I}$ (in both spin states), while F has the greater ΔE_π value. Also, OH has a ΔE_π value similar to that of PH_2 (for the singlet) but a much smaller $\xi(\text{Mo-X})$ value, similar to those of the halogens. These data illustrate that the percent bond contraction should *not* be used in general as a measure of relative π bond strengths.

An independent evaluation of the π bond strength has been obtained for those ligands, namely OH and PH_2 , for which the strength of the π interaction depends on the angular orientation. For singlet $\text{CpMoX}(\text{PH}_3)_2$ ($\text{X} = \text{OH}, \text{PH}_2$), single point calculations have been carried out on geometries derived from the optimized minimum by gradually rotating the X ligand around the Mo-X bond. The results are shown in Figure 3.





Figure 3. Plot of the energy of $\text{CpMo}(\text{EH}_n)(\text{PH}_3)_2$ ($S=0$) vs. the CNT-Mo-E-H dihedral angle. All other geometrical parameters are those of the optimized geometry. Squares: $\text{E} = \text{P}$, $n = 2$; triangles: $\text{E} = \text{O}$, $n = 1$.

For the phosphido system, the maximum π overlap is achieved at the optimized geometry with $\text{CNT-Mo-P-H} = 0^\circ$ or 180° , while the orthogonal orientation gives a zero overlap (see **V** and **VI** in Scheme 4). The energy difference of 23.1 kcal/mole may be taken as an

overestimation of the Mo-PH₂ π bond strength, because of the unrelaxed nature of the CpMo(PH₃)₂ moiety in the rotated geometry. The window between this overestimated value and the underestimated one of Table 1 is relatively narrow.

Scheme 4



For the hydroxide system, the two C_s -symmetric structures (CNT-Mo-O-H angles of 0° , *endo*, see **VII**, and 180° , *exo*) are not identical, because of the greater filled-filled repulsion between the oxygen sp^x lone pair and the Mo d_{z^2} lone pair in the *exo* conformation. In the 90° conformation (**VIII**), Mo-O π bonding is not quite zero, since OH is still able to overlap with the Mo d_{xy} orbital (albeit not so as effectively) by using the sp^x lone pair. At any rate, the energy difference between the most stable *exo* structure and the 90° maximum is 18 kcal/mol and is again slightly higher than the ΔE_π values calculated for OH in Table 1.

c. Steric interactions. The bond dissociation energy of PH₃ from CpMoX(PH₃)₃ is modulated by π donation and by the release of pairing energy upon adoption of the triplet configuration, both being functions of the nature of X. However, differences in size and electronic environment for the various X ligands will also cause differences in the steric pressures exerted on the rest of the coordination sphere and provide an additional contribution

to affect the Mo-PH₃ bond dissociation energy. These differences may be evaluated by comparing the relaxation energies of the fragments that are obtained from the Mo-X bond breaking process for each system. Thus, $(-\Delta E_1 + \Delta E_2) = \Delta E_{\text{steric}}$ (triplet) is a measure of the release of steric pressure associated to the formation of the triplet 16-electron product, while $(-\Delta E_1 + \Delta E_3) = \Delta E_{\text{steric}}$ (singlet) measures the same parameter for the formation of the singlet product. These values are again collected in Table 1. Positive values indicate greater pressure on the more saturated system, as is expected. In some cases small values are obtained by subtraction of individual large relaxation energies. For each X ligand, the singlet state gives rise to a greater ΔE_{steric} value. This result correlates with the greater distortion that the CpMo(PH₃)₂ fragment has to withstand to lead to the triplet geometry upon binding X, thus more efficiently balancing out the greater relaxation experienced by the more saturated system.

We can immediately observe that the smallest X ligand, H, gives small steric contributions. The optimized geometries of both CpMo(PH₃)₃ and CpMo(PH₃)₂ fragments are quite close to those of the same fragments in the corresponding optimized hydrides, and the individual relaxation energies ΔE_1 , ΔE_2 and ΔE_3 (3.2, 2.0, and 1.4 kcal/mol, respectively) are smaller than those of the other X ligands, indicating the H has a small steric influence in both 18-electron and 16-electron structures. It is notable that ΔE_{steric} is relatively large for Cl and Br, while it is smaller for I and F. It is unexpectedly small for PH₂, especially considering that the size of P is similar to that of Cl, and it is even smaller for CH₃. These values suggest to us that the major steric repulsion derives from the interaction between the X lone pairs and the neighboring PH₃ ligands. The Cl and Br atoms have two *p* lone pairs oriented perpendicularly to the Mo-X bond, extending in the region of space occupied by the PH₃ ligands in the 18-electron complex (see Figure 4). On an energy scale, these two orbitals are placed immediately underneath the two filled metal orbitals. Correspondingly, OH has only one *p* lone pair (the second lone pair is a hybrid orbital pointing away from the PH₃ ligands,¹¹ as shown in VII), while PH₂ and CH₃ have none (the only P lone pair is a hybrid orbital with a large amount of *s* character, pointing away from the adjacent PH₃ ligands). This is a manifestation of an interligand *filled-filled*

repulsion. The *filled-filled* repulsion between the X lone pairs and the filled metal orbitals, on the other hand, is accounted for in the ΔE_π term.

It is interesting to observe the trend of ΔE_{steric} values for the halogen series in the order (for both spin states) $F < \text{Cl} \sim \text{Br} > \text{I}$. This result is determined for the most part by the trend in the rearrangement factor ΔE_1 for the 18-electron compounds (7.11, 12.74, 13.35, and 8.7 kcal/mol for F, Cl, Br and I, respectively). The small value of ΔE_{steric} observed for F may be attributed to the small size of the atom and of its orbitals, leading to a small interaction with the adjacent PH_3 ligands. The smaller value for I relative to Cl and Br, on the other hand, may be attributed to the longer distance and greater diffuseness of the orbitals. These considerations appear consistent with the shape of the orbitals as shown in Figure 4.




Figure 4. MOLDEN views of the 3rd (left) and 4th (right) highest MO for $\text{CpMoX}(\text{PH}_3)_3$.

The orbital contour lines correspond to 4 % of the maximum electron density.

d. A comparison of stabilizing effects. We have examined, in the three preceding sections, the effects of the singlet - triplet conversion, of the π -stabilization, and of the release of steric pressure to the relative stabilization of the open-shell system deriving from dissociation of a PH_3 ligand from $\text{CpMoX}(\text{PH}_3)_3$ (Schemes 1 and 2). The correction of the Mo-PH₃ BDE (for either the singlet or the triplet 16-electron product) by the ΔE_π and the ΔE_{steric} parameters (Table 1) provides the BDE for the Mo-PH₃ bond in geometry optimized spin doublet $\text{CpMo}(\text{PH}_3)_3$ (relative to optimized doublet and quartet $\text{CpMo}(\text{PH}_3)_2$, respectively). These values are independent on the nature of X and are calculated as 24.3 and 25.4 kcal/mol, respectively, see eqs. 1 and 2.

$$\text{BDE (singlet)} + \Delta E_\pi (\text{singlet}) + \Delta E_{\text{steric}} (\text{singlet}) = 24.3 \text{ kcal/mol} \quad (1)$$

$$\text{BDE (triplet)} + \Delta E_\pi (\text{triplet}) + \Delta E_{\text{steric}} (\text{triplet}) = 25.4 \text{ kcal/mol} \quad (2)$$

These values are very close to each other because the optimized doublet and quartet $\text{CpMo}(\text{PH}_3)_2$ species have very similar energies, and can be taken as measures of the “intrinsic” Mo-PH₃ bond strength (for each spin state) in the absence of any stabilizing factor introduced by the presence of X. The introduction of X modulates these BDE’s by stabilizing the less saturated and/or by destabilizing the more saturated structure by steric and π bonding effects. The introduction of X also has a large effect on the electron pairing energy, as shown by the variable singlet-triplet gap (varying from 8.7 kcal/mol for F to -7.9 kcal/mol for PH_2) for the 16-electron $\text{CpMoX}(\text{PH}_3)_2$, as compared with the doublet-quartet gap of -1.10 kcal/mol for the 15-electron $\text{CpMo}(\text{PH}_3)_2$.

It is impossible to completely separate the effects caused by these stabilizing factors when one wishes to analyze the stabilization of *triplet* $\text{CpMoX}(\text{PH}_3)_2$ (*i.e.* the ground state in all cases except for $\text{X} = \text{PH}_2$) and free PH_3 relative to *singlet* $\text{CpMoX}(\text{PH}_3)_3$, because these effects are linked to the structural changes associated with the change of spin state. We have already

pointed out that pairing energy effects and π bonding effects are combined in the ΔE_{π} term in equation 2. A convenient, though admittedly limited, approach is given in equation 3.

$$\text{BDE (triplet)} + \Delta E_{\text{S-T}} + \Delta E_{\pi} (\text{singlet}) + \Delta E_{\text{steric}} (\text{singlet}) = 24.30 \text{ kcal/mol} \quad (3)$$

This approach corresponds to the following stepwise process. The optimized $\text{CpMo}(\text{PH}_3)_3$ on one side and the combination of optimized *doublet* $\text{CpMo}(\text{PH}_3)_2$ and free PH_3 on the other side (relative energy 24.30 kcal/mol) are distorted to the geometries of the corresponding fragments in the adducts with X [$\Delta E_{\text{steric}} (\text{singlet})$]. Subsequently, X^{\bullet} is added to form the Mo-X bond, involving BET (σ) for $\text{CpMoX}(\text{PH}_3)_3$ and BET ($\sigma+\pi$) for $\text{CpMoX}(\text{PH}_3)_2$ [$\Delta E_{\pi} (\text{singlet})$]. Finally, singlet $\text{CpMoX}(\text{PH}_3)_2$ is relaxed to the triplet ground state ($\Delta E_{\text{S-T}}$). While this scheme has limitations due to the interrelated nature of these effects, the same would hold true for possible alternative schemes.

The comparison of the $\Delta E_{\text{S-T}}$, singlet ΔE_{π} and singlet ΔE_{steric} values in Table 1 for each X give interesting indications on the relative importance of spin-pairing effects, ligand π donation, and release of steric pressure for the relative stabilization of the 16-electron $\text{CpMoX}(\text{PH}_3)_2$ and therefore for the dependence of the Mo-PH₃ bond dissociation energy on the nature of X. The hydride system has a relatively unencumbering ($\Delta E_{\text{steric}} = 1.0$ kcal/mol) and π neutral ($\Delta E_{\pi} = -2.0$ kcal/mol) X group. Since the triplet state is only slightly stabilized relative to the singlet one ($\Delta E_{\text{S-T}} = 1.8$ kcal/mol), the Mo-PH₃ BDE is quite close to the intrinsic bond strength found in $\text{CpMo}(\text{PH}_3)_3$ and is the highest found for the series of X systems investigated in this work. For all other X systems, the Mo-PH₃ BDE drops dramatically, but for different reasons. As discussed earlier, the steric stabilization is mostly related to the presence of inactive lone pairs. For the π neutral CH₃ system, there is essentially no sterics-related stabilization, while a significant ΔE_{π} stabilization is related to the establishment of an agostic Mo-C-H interaction. The change of spin state also contributes to a large extent to the stabilization of the 16-electron methyl system. We cannot rationalize the difference between $\Delta E_{\text{S-T}}$ for H and CH₃. The

halogen systems are stabilized to a significant extent by each of the three effects, the ΔE_{S-T} stabilization being nearly the same for all four (in the 8.7 - 6.6 kcal/mol interval, predictably in the order $F > Cl > Br > I$), while the other two effects vary as a function of X in a way as to provide an approximately identical total contribution (in the 9-11 kcal/mol range; mostly ΔE_{π} for F, mostly ΔE_{steric} for Cl and Br, and about 50:50 for I). The OH system experiences an even greater overall stabilization, which is mostly due to O-M π donation, but ΔE_{S-T} and ΔE_{steric} also play a role, especially the latter one. Finally, the PH₂ system (which is the only one examined here to adopt a singlet ground state) is almost entirely stabilized by π effects, while the steric factor is nearly zero.

Conclusions

We have examined the relative importance of three difference factors (pairing energy associated to a spin state change, π donation from ligand lone pairs, and release of steric pressure) in the relative stabilization of the PH₃ dissociation product from CpMoX(PH₃)₃ as a function of X (X = H, CH₃, F, Cl, Br, I, OH and PH₂). These are model systems for a class of phosphine-substituted half-sandwich Mo(II) compounds which exhibit richness in structure, magnetic properties, and chemical reactivity. The study has shown that the three effects can display all possible orders of relative importance depending on the nature of the donor atom, on the presence of inactive lone pairs, and on the number of π orbitals available (π neutral, single- or double-sided π donor). The spin state change plays a dominant role for the stabilization of the halide derivatives and the methyl derivative. The study has further shown a contribution to stabilization from the CH₃ group attributable to the establishment of an α -agostic interaction.

Acknowledgement. We are grateful to the Conseil Régional de Bourgogne for support and to the European Commission for a Marie Curie postdoctoral fellowship to K.M.S. We also thank Odile Eisenstein for helpful discussion.

References

- (1) Collman, J. P.; Hegedus, L. S.; Norton, J. R.; Finke, R. G. *Principles and Applications of Organotransition Metal Chemistry*; University Science Books: Mill Valley, CA, 1987.
- (2) Tolman, C. A. *Chem. Soc. Rev.* **1972**, *1*, 337-353.
- (3) Tolman, C. A. *Chem. Rev.* **1977**, *77*, 313-348.
- (4) Caulton, K. G. *New J. Chem.* **1994**, *18*, 25-41.
- (5) Ashby, M. T. *Comments Inorg. Chem.* **1990**, *10*, 297-313.
- (6) Poli, R. *Chem. Rev.* **1996**, *96*, 2135-2204.
- (7) Abugideiri, F.; Keogh, D. W.; Poli, R. *J. Chem. Soc., Chem. Commun.* **1994**, 2317-2318.
- (8) Abugideiri, F.; Fettinger, J. C.; Keogh, D. W.; Poli, R. *Organometallics* **1996**, *15*, 4407-4416.
- (9) Keogh, D. W.; Poli, R. *J. Am. Chem. Soc.* **1997**, *119*, 2516-2523.
- (10) Fettinger, J. C.; Kraatz, H.-B.; Poli, R.; Quadrelli, E. A. *Chem. Commun.* **1997**, 889-890.
- (11) Poli, R.; Quadrelli, E. A. *New J. Chem.* **1998**, *22*, 435-450.
- (12) Chisholm, M. H.; Clark, D. L. *Comments Inorg. Chem.* **1987**, *6*, 23-40.
- (13) Kapp, J.; Schade, C.; El-Nahasa, A. M.; von Ragué Scheleyer, P. *Angew. Chem., Int. Ed. Engl.* **1996**, *35*, 2236-2237.
- (14) González-Blanco, O.; Brachadell, V.; Monteyne, K.; Ziegler, T. *Inorg. Chem.* **1998**, *37*, 1744-1748.
- (15) Gross, Z.; Mahammed, A.; Barzilay, C. M. *Chem. Commun.* **1998**, 1105-1106.
- (16) Lukens, W. W.; Smith, M. R., III; Andersen, R. A. *J. Am. Chem. Soc.* **1996**, *118*, 1719-1728.
- (17) Poulton, J. T.; Folting, K.; Streib, W. E.; Caulton, K. G. *Inorg. Chem.* **1992**, *31*, 3190-3191.

- (18) Poulton, J. T.; Sigalas, M. P.; Folting, K.; Streib, W. E.; Eisenstein, O.; Caulton, K. G. *Inorg. Chem.* **1994**, *33*, 1476-1485.
- (19) Hogarth, G.; Konidaris, P. C.; Saunders, G. C. *J. Organomet. Chem.* **1991**, *406*, 153-157.
- (20) Tilset, M.; Hamon, J.-R.; Hamon, P. *J. Chem. Soc., Chem. Commun.* **1998**, 765-766.
- (21) Lever, A. B. P. *Inorganic Electronic Spectroscopy*; 2nd ed.; Elsevier: New York, 1984.
- (22) Bendix, J.; Bøgevig, A. *Inorg. Chem.* **1998**, *37*, 5992-6001.
- (23) Lichtenberger, D. L.; Rai-Chauduri, A.; Seidel, M. J.; Gladysz, J. A.; Agbossou, S. K.; Igau, A.; Winter, C. H. *Organometallics* **1991**, *10*, 1355-1364.
- (24) Chisholm, M. H.; Hammond, C. E.; Huffman, J. C. *Polyhedron* **1989**, *8*, 1419-1423.
- (25) Johnson, T. J.; Folting, K.; Streib, W. E.; Martin, J. D.; Huffman, J. C.; Jackson, S. A.; Eisenstein, O.; Caulton, K. G. *Inorg. Chem.* **1995**, *34*, 488-499.
- (26) Atwood, J. D.; Brown, T. L. *J. Am. Chem. Soc.* **1976**, *98*, 3160-3166.
- (27) Bryndza, H. E.; Domaille, P. J.; Paciello, R. A.; Bercaw, J. E. *Organometallics* **1989**, *8*, 379-385.
- (28) Buhro, W. E.; Chisholm, M. H.; Folting, K.; Huffman, J. C.; Martin, J. D.; Streib, W. E. *J. Am. Chem. Soc.* **1992**, *114*, 557-570.
- (29) Cooper, A. C.; Bollinger, J. C.; Caulton, K. G. *New J. Chem.* **1998**, 473-480.
- (30) Edmiston, C.; Ruedenberg, K. *Rev. Mod. Phys.* **1963**, *35*, 457.
- (31) Edmiston, C.; Ruedenberg, K. *J. Chem. Phys.* **1965**, *43*, S97-S116.
- (32) Edmiston, C.; Ruedenberg, K. In *Quantum Theory of Atoms, Molecules and Solid State. A Tribute to John C. Slater*; P.-O. Löwdin, Ed.; Academic Press: New York, 1966; pp 263-280.
- (33) Pipek, J.; Mezey, P. G. *J. Chem. Phys.* **1989**, *90*, 4916-4926.
- (34) Umeyama, H.; Morokuma, K. *J. Am. Chem. Soc.* **1976**, *98*, 7208-7220.
- (35) Kitaura, K.; Morokuma, K. *Int. J. Quantum. Chem.* **1976**, *10*, 325-340.
- (36) Bagus, P. S.; Hermann, K.; Bauschlicher C. W., J. *J. Chem. Phys.* **1984**, *80*, 4378-4386.

- (37) Bagus, P. S.; Hermann, K.; Bauschlicher, C. W., Jr. *J. Chem. Phys.* **1984**, *81*, 1966-1974.
- (38) Bauschlicher, C. W., Jr. *Chem. Phys.* **1986**, *106*, 391-398.
- (39) Ziegler, T.; Rauk, A. *Theoret. Chim. Acta* **1977**, *46*, 1-10.
- (40) Ziegler, T.; Rauk, A. *Inorg. Chem.* **1979**, *18*, 1558-1565.
- (41) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Gill, P. M. W.; Johnson, B. G.; Robb, M. A.; Cheeseman, J. R.; Keith, T. A.; Petersson, G. A.; Montgomery, J. A.; Raghavachari, K.; Al-Laham, M. A.; Zakrzewski, V. G.; Ortiz, J. V.; Foresman, J. B.; Cioslowski, J.; Stefanov, B. B.; Nanayakkara, A.; Challacombe, M.; Peng, C. Y.; Ayala, P. Y.; Chen, W.; Wong, M. W.; Andres, J. L.; Replogle, E. S.; Gomperts, R.; Martin, R. L.; Fox, D. J.; Binkley, J. S.; Defrees, D. J.; Baker, J.; Stewart, J. P.; Head-Gordon, M.; Gonzales, C.; Pople, J. A. *Gaussian 94 (Revision E.1)*; Gaussian Inc.: Pittsburgh, PA, 1995.
- (42) Becke, A. D. *J. Chem. Phys.* **1993**, *98*, 5648-5652.
- (43) Dunning, T. H., Jr.; Hay, P. J. In *Modern Theoretical Chemistry*; H. F. Schaefer, III, Ed.; Plenum Press: New York, 1976; pp 1-28.
- (44) Hay, P. J.; Wadt, W. R. *J. Chem. Phys.* **1985**, *82*, 270-283.
- (45) Wadt, W. R.; Hay, P. J. *J. Chem. Phys.* **1985**, *82*, 284-298.
- (46) Hay, P. J.; Wadt, W. R. *J. Chem. Phys.* **1985**, *82*, 299-310.
- (47) Cacelli, I.; Keogh, D. W.; Poli, R.; Rizzo, A. *J. Phys. Chem. A* **1997**, *101*, 9801-9812.
- (48) Poli, R.; Smith, K. M. *Eur. J. Inorg. Chem.* **1999**, 877-880.
- (49) Poli, R.; Smith, K. M. *Eur. J. Inorg. Chem.*, in press.
- (50) Siegbahn, P. E. M. *J. Am. Chem. Soc.* **1996**, *118*, 1487-1496.
- (51) Wang, W.; Weitz, E. *J. Phys. Chem. A* **1997**, *101*, 2358-2363.
- (52) Yang, H.; Asplund, M. C.; Kotz, K. T.; Wilkens, M. J.; Frei, H.; Harris, C. B. *J. Am. Chem. Soc.* **1998**, *120*, 10154-10165.
- (53) Davidson, E. R.; Feller, D. *Chem. Rev.* **1986**, *86*, 681-696.
- (54) Boys, S. F.; Bernardi, F. *Mol. Phys.* **1970**, *19*, 553-566.

- (55) Abugideiri, F.; Kelland, M. A.; Poli, R.; Rheingold, A. L. *Organometallics* **1992**, *11*, 1303-1311.
- (56) Kubáček, P.; Hoffmann, R.; Havlas, Z. *Organometallics* **1982**, *1*, 180-188.
- (57) Ward, T. R.; Schafer, O.; Daul, C.; Hofmann, P. *Organometallics* **1997**, *16*, 3207-3215.
- (58) Smith, K. M.; Poli, R.; Legzdins, P. *Chem. Commun.* **1998**, 1903-1904.
- (59) Baerends, E. J.; Gritsenko, O. V.; Van Leeuwen, R. In *Chemical Application of Density-Functional Theory*; B. B. Laird, Ed.; American Chemical Society: Washington D.C., 1996; Vol. 629; pp 42.
- (60) Baerends, E. J.; Gritsenko, O. V. *J. Phys. Chem* **1997**, *101*, 5383-5403.
- (61) Stowasser, R.; Hoffmann, R. *J. Am. Chem. Soc.* **1999**, *121*, 3414-3420.
- (62) Kubáček, P.; Hoffmann, R. *J. Am. Chem. Soc.* **1981**, *103*, 4320-4332.
- (63) Riehl, J.-F.; Jean, Y.; Eisenstein, O.; Péliissier, M. *Organometallics* **1992**, *11*, 729-737.
- (64) El-Idrissi, I.; Eisenstein, O.; Jean, Y. *New J. Chem.* **1990**, *14*, 671-677.
- (65) Albinati, A.; Bakhmutov, V. I.; Caulton, K. G.; Clot, E.; Eckert, J.; Eisenstein, O.; Gusev, D. G.; Grushin, V. V.; Hauger, B. E.; Klooster, W. T.; Koetzle, T. F.; McMullan, R. K.; O'Laughlin, T. J.; Péliissier, M.; Ricci, J. S.; Sigalas, M. P.; Vymenits, A. B. *J. Am. Chem. Soc.* **1993**, *115*, 7300-7312.
- (66) Baker, R. T.; Calabrese, J. C.; Harlow, R. L.; Williams, I. D. *Organometallics* **1993**, *12*, 830-841.
- (67) Schmid, R.; Herrmann, W. A.; Frenking, G. *Organometallics* **1997**, *16*, 701-708.
- (68) González-Blanco, O.; Branchadell, V. *Organometallics* **1997**, *16*, 5556-5562.
- (69) Martinho Simões, J. A.; Beauchamp, J. L. *Chem. Rev.* **1990**, *90*, 629-688.
- (70) Dobler, M. In *Laboratory for organic chemistry, ETH Zürich, Switzerland*, 1996.