

1-1-1998

Carbon monoxide oxidation by bacteria associated with the roots of freshwater macrophytes

Jeremy J. Rich
University of Maine

G. M. King
University of Maine

Follow this and additional works at: https://repository.lsu.edu/biosci_pubs

Recommended Citation

Rich, J., & King, G. (1998). Carbon monoxide oxidation by bacteria associated with the roots of freshwater macrophytes. *Applied and Environmental Microbiology*, 64 (12), 4939-4943. <https://doi.org/10.1128/aem.64.12.4939-4943.1998>

This Article is brought to you for free and open access by the Department of Biological Sciences at LSU Scholarly Repository. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Scholarly Repository. For more information, please contact ir@lsu.edu.

Carbon Monoxide Oxidation by Bacteria Associated with the Roots of Freshwater Macrophytes†

JEREMY J. RICH‡ AND G. M. KING*

Darling Marine Center, University of Maine, Walpole, Maine 04573

Received 23 June 1998/Accepted 22 September 1998

The potential rates and control of aerobic root-associated carbon monoxide (CO) consumption were assessed by using excised plant roots from five common freshwater macrophytes. Kinetic analyses indicated that the maximum potential uptake velocities for CO consumption ranged from 0.4 to 2.7 μmol of CO g (dry weight)⁻¹ h⁻¹ for the five species. The observed rates were comparable to previously reported rates of root-associated methane uptake. The apparent half-saturation constants for CO consumption ranged from 50 to 370 nM CO; these values are considerably lower than the values obtained for methane uptake. The CO consumption rates reached maximum values at temperatures between 27 and 32°C, and there was a transition to CO production at $\geq 44^\circ\text{C}$, most likely as a result of thermochemical organic matter decomposition. Incubation of roots with organic substrates (e.g., 5 mM syringic acid, glucose, alanine, and acetate) dramatically reduced the rate of CO consumption, perhaps reflecting a shift in metabolism by facultative CO oxidizers. Based on responses to a suite of antibiotics, most of the CO consumption (about 90%) was due to eubacteria rather than fungi or other eucaryotes. Based on the results of acetylene inhibition experiments, methanotrophs and ammonia oxidizers were not active CO consumers.

Carbon monoxide (CO) is one of the most important chemical reactants in the troposphere. It influences the fate of methane and ozone by removing the major atmospheric oxidizing agent, hydroxyl radical (29). The background CO mixing ratios range from about 50 to 300 ppb and vary considerably over time and space (33). Fossil fuel use, biomass burning, and oxidation of atmospheric hydrocarbons (methane and other compounds) account for most of the CO source strength (1,200 to 2,850 Tg year⁻¹), and direct natural emissions from oceans and vegetation (80 to 360 Tg year⁻¹ [34]) account for 10 to 15% of the total. However, each of these components of the CO budget is uncertain (34), thus limiting an accurate assessment of the effects of CO on atmospheric chemistry and climate change.

The role of wetlands in the global CO cycle has not been adequately investigated. Several limited studies have indicated that wetlands are not important. Conrad et al. (17) and Khalil and Rasmussen (23) measured CO emissions from rice paddies and extrapolated the results globally to predict that wetland CO emissions are a minor part of the CO budget at most. However, these results have not been validated with natural wetland data, and emissions may have been underestimated since the roles of photochemical reactions with dissolved organic carbon (9, 43, 46) and vegetation (17, 39, 42) were ignored. In addition, Funk et al. (20) showed that CO emissions were considerably higher than expected based on extrapolations made by Conrad et al. (17).

CO emissions from wetlands and their relative importance in the global CO budget may be determined in part by populations of root-associated CO oxidizers. As is the case for methane, CO produced in sediments may be transported through roots and stems to the atmosphere. Both aerobic and

anaerobic CO oxidizers growing on or in plant roots could significantly attenuate such a flux. The aerobic CO oxidizers include carboxydrotrophs that grow chemolithotrophically on CO or heterotrophically on various organic compounds (30–32, 37, 45); the significance of carboxydrotrophs in situ is unknown but has been questioned (13). Methanotrophs, ammonia oxidizers, and certain fungi also consume CO (5, 39). Although each of these groups may contribute to CO oxidation in soil (10a, 12), the role of these groups in wetlands has not been addressed.

The objective of the study described here was to characterize aerobic root-associated CO consumption. Potential rates and controls of activity were assessed in vitro following using methods previously used to characterize root-associated methanotrophy (25). Kinetic analyses of CO consumption revealed that there were substantial variations among plant taxa and root types. Eubacteria were primarily responsible for CO consumption, but methanotrophs and ammonium oxidizers were probably not involved. Organic substrates limited CO consumption, and uptake had a mesophilic response to temperature.

MATERIALS AND METHODS

Sampling and root processing. Plants were obtained from a wetland near Bristol, Maine (26). As described previously, intact plants (roots and sediment) were removed with a spade and transported to the laboratory for analysis (25). Sediment was washed from the plants and intact root systems with tap water. Roots were cut from rhizomes, carefully teased apart, rinsed again, and sorted. Subsamples of individual roots (length, up to 45 cm) were obtained after extraneous particulate organic matter was removed. The sampling procedure minimized damage to the roots and any associated loss of soluble organic compounds. *Pontederia cordata* roots were used for all temperature response, organic carbon, and antibiotic assays (which were conducted in August through October 1997). Wet roots were blotted gently with paper towels and sealed in 50-ml culture tubes containing 1 to 2 ml of deionized water and having neoprene stoppers. Root volume was determined after kinetic analyses by water displacement; the root volume was $4.0 \pm 0.2 \text{ cm}^3$ (mean \pm standard error; $n = 24$). All experiments were initiated within 10 h of the time that plant samples were obtained, and clean latex gloves were worn when the roots were handled. The roots were incubated with ambient laboratory lighting during time course assays.

Differences in CO consumption as a function of root type were determined by separating *P. cordata* roots into four groups based on the following simple visual distinctions: color, form, thickness, fine root growth, and position on the plant. The root types distinguished were “feeder,” “shallow-anchor,” “deep-anchor,”

* Corresponding author. Mailing address: Darling Marine Center, University of Maine, Walpole, ME 04573. Phone: (207) 563-3146, ext. 207. Fax: (207) 563-3119. E-mail: gking@maine.edu.

† Contribution 321 from the Darling Marine Center.

‡ Present address: Department of Crop and Soil Science, Oregon State University, Corvallis, OR 97331-7306.

TABLE 1. Kinetic constants for CO oxidation by macrophyte roots

Source of roots	V_{\max} (μmol of CO g [dry wt] of root ⁻¹ h ⁻¹) ^a	K_{app} (nM CO) ^a
<i>Pontederia cordata</i>		
Feeder roots	1.2 \pm 0.2	50 \pm 5 A ^b
Shallow-anchor roots	1.1 \pm 0.1	150 \pm 30
Miscellaneous roots	1.1 \pm 0.2	180 \pm 10
Deep-anchor roots	1.2 \pm 0.4	260 \pm 40 B
<i>Sagittaria latifolia</i>		
	2.7 \pm 0.1B	370 \pm 60 A
<i>Sparganium eurycarpum</i>		
	0.9 \pm 0.2A	230 \pm 20
<i>Elymus riparius</i>		
	0.4 \pm 0.1	180 \pm 60
<i>Typha latifolia</i>		
	0.5 \pm 0.2	140 \pm 40

^a Data are means \pm standard deviations ($n = 3$). We assumed that the solubility coefficient was 0.018 cm³ of CO ml of H₂O⁻¹ at 20°C (36).

^b Values followed by different letters in the same column are significantly different as determined by Tukey's test ($P < 0.05$); the data for *P. cordata* and the data for the four other taxa were treated as separate data sets for the statistical analysis.

and "miscellaneous" roots. The feeder roots had extensive lateral fine root growth (twofold more than anchor roots on a dry weight basis). Anchor roots had thicker tap roots. Deep-anchor roots were coated with brown iron oxide plaques, and shallow-anchor roots were white or purple. Miscellaneous roots had combinations of features. Feeder roots grew at the sediment-water interface, while shallow- and deep-anchor roots penetrated into the sediment matrix.

Differences in CO consumption among taxa were determined by using triplicate samples of five species (Table 1) that grew under similar conditions (i.e., relatively dense colonies within an approximately 30-m radius; *Sagittaria latifolia* occurred as solitary plants). The roots of each plant were sorted to ensure that samples were monospecific.

Kinetic analyses. Kinetic data were obtained by using triplicate samples consisting of approximately 3 g (fresh weight) of roots obtained during early autumn. CO consumption was assayed at headspace CO concentrations of <30 ppm in order to obtain a first-order rate constant (k), and this was followed immediately by an assay at a headspace CO concentration of approximately 1% to obtain the maximum potential uptake velocity (V_{\max}). V_{\max} and k were used to approximate the apparent half-saturation constant of the Michaelis-Menten equation (K_{app}) from the relationship $k = V_{\max} K_{\text{app}}^{-1}$. This approach, which was based on the Michaelis-Menten model, was chosen in order to minimize the variability inherent in the use of replicate samples for a series of concentration-versus-rate assays and because use of individual roots for a complete concentration series (i.e., subsaturating to saturating concentrations) would have required excessively long incubations.

Temperature analysis. k values were obtained at 23 to 25°C by using headspace CO concentrations of <20 ppm for triplicate 0.7-g (fresh weight) samples of *P. cordata* feeder roots. After this initial assay, which lasted about 7 h, the roots were vented with ambient air and incubated for 2 to 4 h at temperatures ranging from 2 to 44°C without CO. After CO (<20 ppm) was added, a second uptake assay was used to determine k as a function of temperature.

Response to added organic substrates. Triplicate samples consisting of approximately 1.5 g (fresh weight) of *P. cordata* deep-anchor roots were incubated horizontally at 30°C on a rotary shaker (100 rpm) for 5 days with a headspace containing 2,000 ppm of CO and an aqueous phase containing 20 ml of either syringic acid, glucose, alanine, or acetate (5 mM). The controls were treated with deionized water and 2,000 ppm of CO. On day 5, the incubation medium was decanted and centrifuged (15,000 \times g, 30 min), and the pellet was resuspended in 2 ml of fresh substrate, which was transferred to 30-ml tubes. The roots were blotted with paper towels and returned to the culture tubes. The solution and roots were assayed for CO consumption rates with a headspace containing approximately 1,000 ppm of CO. A linear function best described CO consumption at this mixing ratio.

Antibiotic assay. Antibiotics at the following concentrations (in deionized water) were used to differentiate among fungi, bacteria, and plant cells: streptomycin, 100 μg ml⁻¹; ampicillin, 50 μg ml⁻¹; chlortetracycline, 20 μg ml⁻¹; nystatin, 50 μg ml⁻¹; and cycloheximide, 50 μg ml⁻¹. Twenty milliliters of each fresh antibiotic solution and 500 ppm of CO were added to triplicate samples consisting of approximately 2 g (fresh weight) of *P. cordata* deep-anchor roots incubated horizontally at 30°C on a rotary shaker (100 rpm). The incubation medium was decanted after 24 h, and the excess liquid was blotted from the roots with paper towels. CO consumption rates were obtained by using a headspace CO concentration of approximately 1,000 ppm. Immediately after this initial assay, fresh antibiotic solution was added to the roots, and the preparations were incubated for an additional 4 days. On day 5, the CO consumption assay was repeated. The controls were treated with only deionized water and CO.

Acetylene inhibition. The potential role of methanotrophic and ammonia-oxidizing bacteria was assessed by using acetylene, which is a well-known non-specific irreversible inhibitor of methane and ammonia monooxygenases. Washed, sediment-free roots (about 1 g [fresh weight]) of *Sparganium eurycarpum*, *S. latifolia*, and *Typha latifolia* were incubated with anoxic headspace in triplicate in 40-cm³ tubes containing 1 ml of deionized water. The tubes were sealed with neoprene stoppers, and the headspaces of three tubes for each root type were amended with 1% acetylene; unamended tubes served as controls. All of the tubes received 0.5 μCi (18.5 kBq) of ¹⁴C prepared by dehydrating [¹⁴C]formic acid (60 mCi mmol⁻¹; 2.2 GBq mmol⁻¹; New England Nuclear) as described by Bartholomew and Alexander (2). The total headspace CO concentrations were approximately 1 ppm. Headspace subsamples (0.5 cm³) were removed at intervals to assay for ¹⁴CO₂ as an index of CO oxidation. The ¹⁴CO₂ was trapped in 2 ml of 1 N KOH, and then the amount of ¹⁴CO₂ was determined by liquid scintillation counting with an LKB model 1217 counter (LKB-Wallac Instruments, Inc.).

Gas chromatography. Headspace CO concentrations less than 30 ppm were measured by using 1-cm³ gas samples injected into a model RGA3 instrument (Trace Analytical, Inc.) equipped with a mercury vapor detector. The instrument was fitted with a gas sampling valve and a 1-m Molecular Sieve type 5A column operated at 105°C with an airflow rate of 20 cm³ min⁻¹. The detector was operated at 265°C. The detection limit was <10 ppb of CO. The retention times for hydrogen and CO were 0.5 and 1.5 min, respectively, and the peaks were well separated. The instrument was standardized by using a certified 91.9-ppb CO standard (National Oceanic and Atmospheric Administration) a 10.8-ppm CO standard (Maine Oxy Supply Co.), and laboratory dilutions (1 to 40 ppm) of certified 996-ppm CO (Maine Oxy Supply Co.). The instrument response at concentrations above about 2 to 5 ppm was nonlinear, and the relationship between detector response and concentration was established by nonlinear curve fitting.

Headspace samples (volume, 0.1 cm³) with CO concentrations of >0.01% were analyzed with a model 3700 gas chromatograph (Varian Instruments, Inc.) equipped with a methanizer (SRI Instruments, Inc.) and a flame ionization detector. The columns used for gas separation were a Poropak Q column (length, 1 m; outside diameter, 0.125 in.; inside diameter, 0.0625 in.) in series with a silica gel column (Davison Grade 12; 80/100; Supelco, Inc.) to retain hydrocarbons and a Molecular Sieve type 5A column (length, 1 m; outside diameter, 0.125 in.; inside diameter, 0.0625 in.) to retain CO₂. The columns were operated at 60°C, and the air, H₂, and N₂ carrier gas flow rates were approximately 30, 300, and 30 cm³ min⁻¹, respectively. The flame ionization detector, injector, and methanizer were operated at 200, 80, and 380°C, respectively. The instrument was standardized with a certified 996-ppm CO standard (Maine Oxy Supply Co.) and laboratory dilutions (0.25 to 1.5% CO) of pure 99.3% CO (Scott Specialty Gases, Inc.).

Statistical analyses. Headspace CO mixing ratios were plotted as a function of time for individual replicates; rate constants (i.e., k and V_{\max}) were determined from linear and nonlinear regression analyses by using Kaleidagraph software (Synergy, Inc.). The significance of regression analyses and the significance of differences among means were determined by using analysis of variance (Systat); Tukey's test for mean multiple comparisons was performed to define significant groups of samples.

RESULTS

Rate estimates. Roots under aerobic conditions (Fig. 1) rapidly and efficiently consumed CO. The kinetic constants for the various plant taxa differed significantly (Table 1). The V_{\max} for *S. latifolia* (2.7 μmol of CO g [dry weight]⁻¹ h⁻¹) exceeded by a factor of more than 2 the V_{\max} values for *S. eurycarpum* and *P. cordata* (0.9 to 1.1 μmol of CO g [dry weight]⁻¹ h⁻¹) and by a factor of 5 the V_{\max} values for *Elymus riparius* and *T. latifolia*, (0.4 to 0.5 μmol of CO g [dry weight]⁻¹ h⁻¹) (Table 1). In contrast, the V_{\max} values for the various *P. cordata* root types were very similar (Table 1).

The K_{app} values differed significantly as a function of root type and plant taxa. The feeder roots exhibited the highest affinity for CO, with a K_{app} of 54 \pm 5 nM CO (67 \pm 6 ppm of CO) (mean \pm standard deviation). The K_{app} values for shallow and miscellaneous roots were intermediate, 154 \pm 25 to 182 \pm 13 nM CO (192 \pm 32 to 226 \pm 16 ppm of CO), while the K_{app} values for deep-anchor roots were 258 \pm 40 nM CO (321 \pm 50 ppm of CO). The K_{app} values for *S. latifolia*, 367 \pm 63 nM CO (457 \pm 78 ppm of CO), were significantly different from the K_{app} values for *T. latifolia*, *E. riparius*, and *S. eurycarpum*, which were 137 \pm 40 to 229 \pm 21 nM CO (171 \pm 50 to 285 \pm 27 ppm of CO). In general, the K_{app} values (0.1 to 0.4 μM CO) were




FIG. 1. Root-associated CO oxidation by different *P. cordata* root types (see text) (A) and different plant taxa (B). (A) Symbols: ●, feeder roots; ○, shallow-anchor roots; □, miscellaneous roots; ■, deep-anchor roots. (B) Symbols: ●, *S. latifolia*; ○, *S. eurycarpum*; □, *T. latifolia*; ■, *E. riparius*. The data are means ± 1 standard error ($n = 3$) fit by exponential (A) and linear (B) regression.

less variable than the V_{maxp} values (0.4 to 2.7 μmol of CO g [dry weight] $^{-1}$ h $^{-1}$).

The biomass of each of the four root types was measured, and most of the roots (45% ± 10% [mean ± standard deviation; $n = 2$]) were miscellaneous roots, which were roots that had combinations of features. The relative biomasses of the feeder, shallow-anchor, and deep-anchor roots were 6% ± 2%, 22% ± 9%, and 27% ± 1% ($n = 2$), respectively.

Temperature analysis. CO consumption by roots increased with temperature from 2°C, reaching a maximum at temperatures between 27 and 32°C (Fig. 2) and decreasing at higher temperatures. Net CO production was evident at temperatures




FIG. 2. Carbon monoxide consumption as a function of temperature for roots of *P. cordata*. The curve is a weighted fit, and the data are means ± 1 standard error ($n = 3$). gdw, gram (dry weight).




FIG. 3. Carbon monoxide consumption after 5-days of incubation with 5 mM organic substrate and a headspace CO concentration of 2,000 ppm. Organic compounds were not added to the control. Data are means ± 1 standard error ($n = 3$). gdw, gram (dry weight).

of $\geq 44^\circ\text{C}$ (Fig. 2). The activation energy (E_a) for CO oxidation determined from an Arrhenius plot for temperatures from 2 to 29°C was 59 kJ mol $^{-1}$ ($P < 0.0001$) (data not shown). The Q_{10} value from 2 to 29°C was approximately 2.

Response to added organic substrates. The rates of CO consumption decreased significantly compared to controls with roots incubated with a headspace containing 2,000 ppm of CO and solutions containing various simple organic compounds at a concentration of 5 mM (Fig. 3). All of the compounds assayed yielded similar results, which did not differ statistically. Negligible CO consumption was observed in the incubation medium after root tissue was removed.

Antibiotic assay. The antifungal agents nystatin and cycloheximide had no effect on CO consumption, while two of the antibacterial agents significantly reduced activity (Fig. 4). Streptomycin was the most effective inhibitor, decreasing the CO consumption rate by 90% compared to controls; ampicillin inhibited activity to a lesser extent. Inhibition by streptomycin and ampicillin increased over time and was not significant until day 5. Chlortetracycline, a general antimicrobial agent, had no effect on CO consumption. Although the antifungal agents and chlortetracycline appeared to slightly stimulate CO consumption, the difference was insignificant ($P \geq 0.052$).




FIG. 4. Carbon monoxide consumption after incubation with antibiotics and 500 ppm of CO for 1 day (open bars) and 5 days (solid bars). See the text for the concentrations of antibiotics used. Amp, ampicillin; CTC, chlortetracycline; Cyc, cycloheximide; Nys, nystatin; Str, streptomycin. The following values were significantly different ($P \leq 0.007$): day 5 control value and day 5 streptomycin value; day 5 control value and day 5 ampicillin value; and day 5 streptomycin value and day 1 streptomycin value. Data are means ± 1 standard error ($n = 3$). gdw, gram (dry weight).




FIG. 5. Time course for accumulation of $^{14}\text{CO}_2$ from ^{14}CO oxidation by roots of *S. eurycarpum* (circles) and *T. latifolia* (squares). The open and solid symbols indicate the data for controls and 1% acetylene treatment, respectively. gfw, gram (fresh weight).

Acetylene inhibition. ^{14}CO was oxidized rapidly (Fig. 5), and the maximum relative extent of oxidation approached 70% during a 24-h incubation. In these assays, *S. eurycarpum* control roots exhibited the greatest rates of oxidation. The time course of $^{14}\text{CO}_2$ accumulation in controls was consistent with exponential CO uptake since the data fit a model having the following form: $[\text{}^{14}\text{CO}_{2(t)}/\text{}^{14}\text{CO}_{(t0)}] = 1 - \text{}^{14}\text{CO}_{(t0)}e^{-kt}$. CO oxidation was inhibited completely immediately after 1% acetylene was added and for 15 to 25 h thereafter. However, during extended incubation in the presence of acetylene, the extent of oxidation increased significantly (Fig. 5).

DISCUSSION

CO consumption by macrophyte roots apparently can contribute significantly to the fate of wetland CO . A comparison of CO and methane consumption rates is illustrative. The V_{maxp} values for methane oxidation range from 0.3 to 27 μmol of CH_4 g [dry weight] of root $^{-1}$ h $^{-1}$ for a variety of macrophytes (25), including rice (5 μmol of CH_4 g [dry weight] $^{-1}$ h $^{-1}$ [8]). The V_{maxp} values for CO oxidation, 0.5 to 2.7 μmol of CO g (dry weight) of root $^{-1}$ h $^{-1}$ (Table 1), are about 10-fold lower for the same taxa (10). However, the methane concentrations in pore water are more than 100 times greater than the CO concentrations (27). Thus, CO consumption may have an impact similar to that of CH_4 consumption, which reduces potential emissions by 25 to 50% (10, 26).

The V_{maxp} values for CO oxidation varied significantly among plant taxa but not as a function of root type for *P. cordata*. Differences among taxa may reflect differences in root aeration (10, 35, 38, 40) that affect the oxygen and CO concentrations in the rhizosphere, as well as the densities of CO oxidizers. The average K_{app} values ranged from 50 to 370 nM CO among taxa and root types (Table 1). The differences may reflect variations in the root microenvironment. The K_{app} values varied substantially and independently of the V_{maxp} values in a comparison of *P. cordata* roots. In contrast, the V_{maxp} values varied substantially more among taxa than the K_{app} values varied.

Root K_{app} values are greater than soil K_{app} values (4 to 41 nM CO [3, 15, 18, 41]) and water column K_{app} values (7 to 9 nM CO [13]) but less than carboxydrotrophic culture K_{app} values (370 to 890 nM CO [15]). This may indicate that distinct populations of CO oxidizers dominate roots. However, half-saturation constants are not fixed parameters for cultures (7), and differences among roots and other systems may simply

reflect adaptation to different concentration regimes by similar populations.

The E_a for root CO consumption at temperatures from 2 to 29°C was 59 kJ mol $^{-1}$ or about 60% of the value obtained for root-associated methane oxidation (97 kJ mol $^{-1}$ [25]). The lower E_a suggests that CO consumption is less sensitive to temperature; alternatively, simultaneous CO production may reduce the influence of temperature. In support of the latter hypothesis, net CO production was observed at $\geq 44^\circ\text{C}$; this production may have been due to abiological processes similar to those that occur in soil (14). Humic substances, such as those associated with plant roots, contribute to CO formation in soils and may contribute to CO formation in wetlands, although the specific mechanisms involved remain unknown (14).

Preliminary estimates of pore water CO concentrations, which range from about 10 to 100 nM CO (27), are at the lower limit of root K_{app} values (Table 1). This suggests that CO concentrations could limit the rate of CO oxidation in situ. In contrast, pore water CH_4 is less of a limiting factor for root-associated methane oxidation since the K_{app} for this process (3 to 6 μM CH_4 [25]) is considerably less than the in situ concentrations (>100 μM CH_4 [10]). For methane oxidation, oxygen (i.e., root aeration) is a more important control (10). Oxygen may also limit CO consumption but may be less important than other factors, such as the availability of alternative carbon sources.

The dramatic decrease in CO consumption rates in roots incubated with exogenous organic substrates (Fig. 3) may be attributed to the facultative nature of at least some CO -oxidizing bacteria. In general, carboxydrotrophic bacteria express CO dehydrogenase only when CO is the primary growth substrate; heterotrophically grown cells either do not oxidize CO or exhibit a reduced rate of oxidation (15, 24, 30). In the root assays, exogenous organic substrates may have allowed opportunistic heterotrophs to outgrow CO oxidizers or may have decreased the expression of CO dehydrogenase. Abiological CO production from exogenous substrates probably does not explain the results since controls revealed no significant production from the compounds assayed (27).

Streptomycin was the most effective inhibitor of CO consumption, decreasing the rate of uptake by 90%. In contrast, the antifungal agents nystatin and cycloheximide had no effect (Fig. 4). Although the efficacies of the various antibiotics are uncertain, the results suggest that bacteria may be responsible for most of the root-associated CO oxidation and that neither root cells nor fungi are important. In contrast, Conrad and Seiler (12) showed that fungi consumed the majority of CO in certain soils, while bacteria dominated in other soils. Since only one root type was examined in this study, it is not known whether the observed response to antibiotics is typical.

The results of the assay performed with acetylene (Fig. 5) indicate that methanotrophs probably contribute little to root CO consumption despite the fact that they are active on most aquatic plant roots (10, 25, 26) and are capable of consuming CO (5). Acetylene effectively and irreversibly inhibits methane monooxygenase, the enzyme used by methanotrophs to oxidize CO . Thus, the initial inhibition of root CO consumption by acetylene is consistent with methanotrophic activity, but the subsequent reversal of inhibition in the continued presence of acetylene is not. Based on analogous reasoning, a significant role for ammonia oxidizers is also unlikely. Similar results were obtained in experiments in which CO consumption by soils was examined (27); in these experiments reversible inhibition by acetylene and the absence of any effect of methyl fluoride also suggested that methanotrophs and ammonia oxidizers played a

negligible role. The results obtained with both roots and soils indicate that acetylene might be a short-term (albeit nonspecific) inhibitor of CO consumption. However, neither the mechanism of inhibition nor the reason for the reversal observed is clear.

In conclusion, preliminary kinetic and pore water data suggest that root-associated CO oxidizers have the potential to consume significant amounts of CO produced in sediments but may be limited in situ by the presence of organic substrates and the absence of oxygen. Assuming that $V_{\max p}$ is a surrogate for population size and K_{app} is a marker for different microbial populations, the number of root-associated CO-oxidizing bacteria appears to vary among plant species, while specific populations may vary as a function of root type. Thus, multiple populations of CO oxidizers may be present on roots of individual plants. Differences in densities and types of root-associated CO oxidizers may contribute to differences among plant taxa in the extent to which CO is transported from peats or sediments through stems and to the atmosphere. Understanding the characteristics and control of root-associated CO consumption should not only help explain wetland CO emissions but also contribute to a greater understanding of the interactions among microbial populations, plant species, and root types.

ACKNOWLEDGMENTS

This work was supported in part by grant DEB-9528552 from the National Science Foundation.

We thank K. Hardy for technical assistance.

REFERENCES

1. **Armstrong, W.** 1978. Root aeration in the wetland condition, p. 269–297. *In* D. D. Hook and R. M. M. Crawford (ed.), *Plant life in anaerobic environments*. Ann Arbor Science, Ann Arbor, Mich.
2. **Bartholomew, G. W., and M. Alexander.** 1979. Microbial metabolism of carbon monoxide in culture and in soil. *Appl. Environ. Microbiol.* **37**:932–937.
3. **Bartholomew, G. W., and M. Alexander.** 1981. Soil as a sink for atmospheric carbon monoxide. *Science* **212**:1389–1391.
4. **Bartholomew, G. W., and M. Alexander.** 1982. Microorganisms responsible for the oxidation of carbon monoxide in soil. *Environ. Sci. Technol.* **16**:300–301.
5. **Bedard, C., and R. Knowles.** 1989. Physiology, biochemistry, and specific inhibitors of CH_4 , NH_4^+ , and CO oxidation by methanotrophs and nitrifiers. *Microbiol. Rev.* **53**:68–84.
6. **Bender, M., and R. Conrad.** 1994. Microbial oxidation of methane, ammonium, and carbon monoxide, and turnover of nitrous oxide and nitric oxide in soils. *Biogeochemistry* **27**:97–112.
7. **Benstead, J., G. M. King, and H. G. Williams.** 1998. Methanol promotes atmospheric methane oxidation by methanotrophic cultures and soils. *Appl. Environ. Microbiol.* **64**:1091–1098.
8. **Bosse, U., and P. Frenzel.** 1997. Activity and distribution of methane-oxidizing bacteria in flooded-rice soil microcosms and in rice plants (*Oryza sativa*). *Appl. Environ. Microbiol.* **63**:1199–1207.
9. **Bourbonniere, R. A., W. L. Miller, and R. G. Zepp.** 1997. Distribution, flux, and photochemical production of carbon monoxide in a boreal beaver impoundment. *J. Geophys. Res.* **102**(D):29321–29329.
10. **Calhoun, A., and G. M. King.** 1997. Regulation of root-associated methanotrophy by oxygen availability in the rhizosphere of two aquatic macrophytes. *Appl. Environ. Microbiol.* **63**:3051–3058.
- 10a. **Castro, M. S., W. T. Peterjohn, J. M. Mellilo, P. A. Steudler, H. L. Gholz, and D. Lewis.** 1994. Effects of nitrogen fertilization on the fluxes of N_2O , CH_4 and CO_2 from soils in a Florida slash pine plantation. *Can. J. For. Res.* **24**:9–13.
11. **Conrad, R.** 1988. Biogeochemistry and ecophysiology of atmospheric CO and H_2 . *Adv. Microb. Ecol.* **10**:231–283.
12. **Conrad, R., and W. Seiler.** 1980. Role of microorganisms in the consumption and production of atmospheric carbon monoxide by soil. *Appl. Environ. Microbiol.* **40**:437–445.
13. **Conrad, R., and W. Seiler.** 1982. Utilization of traces of carbon monoxide by aerobic oligotrophic microorganisms in ocean, lake and soil. *Arch. Microbiol.* **132**:41–46.
14. **Conrad, R., and W. Seiler.** 1985. Characteristics of abiological carbon monoxide formation from soil organic matter, humic acids, and phenolic compounds. *Environ. Sci. Technol.* **19**:1165–1169.
15. **Conrad, R., O. Meyer, and W. Seiler.** 1981. Role of carboxydobacteria in the consumption of atmospheric carbon monoxide by soil. *Appl. Environ. Microbiol.* **42**:211–215.
16. **Conrad, R., M. Aragno, and W. Seiler.** 1983. Production and consumption of carbon monoxide in a eutrophic lake. *Limnol. Oceanogr.* **28**:42–49.
17. **Conrad, R., H. Schütz, and W. Seiler.** 1988. Emission of carbon monoxide from submerged rice fields into the atmosphere. *Atmos. Environ.* **22**:821–823.
18. **Duggin, J. A., and D. A. Cataldo.** 1985. The rapid oxidation of atmospheric CO to CO_2 by soils. *Soil Biol. Biochem.* **17**:469–474.
19. **Epp, M. A., and J. P. Chanton.** 1993. Rhizospheric methane oxidation determined via the methyl fluoride inhibition technique. *J. Geophys. Res.* **98**(D):18413–18422.
20. **Funk, D. W., E. R. Pullman, K. M. Peterson, P. M. Crill, and W. D. Billings.** 1994. Influence of water table on carbon dioxide, carbon monoxide, and methane fluxes from taiga bog microcosms. *Global Biogeochem. Cycles* **8**:271–278.
21. **Hino, S., and H. Tauchi.** 1987. Production of carbon monoxide from aromatic amino acids by *Morganella morganii*. *Arch. Microbiol.* **148**:167–171.
22. **Jones, R. D., and R. Y. Morita.** 1983. Carbon monoxide oxidation by chemolithotrophic ammonium oxidizers. *Can. J. Microbiol.* **29**:1545–1551.
23. **Khalil, M. A. K., and R. A. Rasmussen.** 1990. Emissions of trace gases from Chinese rice fields and biogas generators: CH_4 , N_2O , CO, CO_2 , chlorocarbons, and hydrocarbons. *Chemosphere* **20**:207–226.
24. **Kiessling, M., and O. Meyer.** 1982. Profitable oxidation of carbon monoxide or hydrogen during heterotrophic growth of *Pseudomonas carboxydoflava*. *FEMS Microbiol. Lett.* **13**:333–338.
25. **King, G. M.** 1994. Associations of methanotrophs with the roots and rhizomes of aquatic vegetation. *Appl. Environ. Microbiol.* **60**:3220–3227.
26. **King, G. M.** 1996. In situ analyses of methane oxidation associated with the roots and rhizomes of a bur reed, *Sparganium eurycarpum*, in a Maine wetland. *Appl. Environ. Microbiol.* **62**:4548–4555.
27. **King, G. M.** 1998. Unpublished results.
28. **Loewus, M. W., and C. C. Delwiche.** 1963. Carbon monoxide production by algae. *Plant Physiol.* **38**:371–374.
29. **Logan, J. A., M. J. Prather, S. C. Wofsy, and M. B. McElroy.** 1981. Tropospheric chemistry: a global perspective. *J. Geophys. Res.* **86**(C):7210–7254.
30. **Meyer, O., and H. G. Schlegel.** 1978. Reisolation of the carbon monoxide utilizing hydrogen bacterium *Pseudomonas carboxydovorans* (Kistner), comb. nov. *Arch. Microbiol.* **118**:35–43.
31. **Meyer, O., K. Frunzke, and G. Mörsdorf.** 1993. Biochemistry of the aerobic utilization of carbon monoxide, p. 433–459. *In* J. C. Murrell and D. P. Kelly (ed.), *Microbial growth on C_1 compounds*. Intercept Limited, Andover, United Kingdom.
32. **Mörsdorf, G., K. Frunzke, D. Gadhari, and O. Meyer.** 1992. Microbial growth on carbon monoxide. *Biodegradation* **3**:61–82.
33. **Novelli, P. C., K. A. Masarie, P. P. Tans, and P. M. Lang.** 1994. Recent changes in atmospheric carbon monoxide. *Science* **263**:1587–1590.
34. **Prather, M., R. Derwent, D. Ehhalt, P. Fraser, E. Sanhueza, and X. Zhou.** 1995. Other trace gases and atmospheric chemistry, p. 73–126. *In* J. T. Houghton et al. (ed.), *Climate change 1994. Radiative forcing of climate change*. Intergovernmental Panel on Climate Change. University Press, Cambridge, United Kingdom.
35. **Sand-Jensen, K., C. Prahl, and H. Stokholm.** 1982. Oxygen release from roots of submerged aquatic macrophytes. *Oikos* **38**:349–354.
36. **Schmidt, U.** 1979. The solubility of carbon monoxide and hydrogen in water and sea-water at partial pressures of about 10^{-5} atmospheres. *Tellus* **31**:68–74.
37. **Schübel, U., M. Kraut, G. Mörsdorf, and O. Meyer.** 1995. Molecular characterization of the gene cluster *coxMSL* encoding the molybdenum-containing carbon monoxide dehydrogenase of *Oligotropha carboxydovorans*. *J. Bacteriol.* **177**:2197–2203.
38. **Sebacher, D. L., R. C. Harris, and K. B. Bartlett.** 1985. Methane emissions to the atmosphere through aquatic plants. *J. Environ. Qual.* **14**:40–46.
39. **Seiler, W.** 1978. The influence of the biosphere on the atmospheric CO and H_2 cycles, p. 773–810. *In* W. E. Krumbain (ed.), *Environmental biogeochemistry and geomicrobiology*. Ann Arbor Science, Ann Arbor, Mich.
40. **Smits, A. J. M., P. Laan, and R. H. Thier.** 1990. Root aerenchyma, oxygen leakage patterns and alcoholic fermentation ability of the roots of some nymphaeid and isoetid macrophytes in relation to the sediment type of their habitat. *Aquat. Bot.* **38**:3–17.
41. **Spratt, H. G., and J. S. Hubbard.** 1981. Carbon monoxide metabolism in roadside soils. *Appl. Environ. Microbiol.* **41**:1192–1201.
42. **Tarr, M. A., W. L. Miller, and R. G. Zepp.** 1995. Direct carbon monoxide production from plant matter. *J. Geophys. Res.* **100**(D):11403–11413.
43. **Valentine, R. L., and R. G. Zepp.** 1993. Formation of carbon monoxide from the photodegradation of terrestrial dissolved organic carbon in natural waters. *Environ. Sci. Technol.* **27**:409–412.
44. **Wolff, D. G., and W. R. Bidlack.** 1976. The formation of carbon monoxide during the peroxidation of microsomal lipids. *Biochem. Biophys. Res. Commun.* **73**:850–857.
45. **Zavarzin, G. A., and A. N. Nozhevnikova.** 1977. Aerobic carboxydobacteria. *Microbiol. Ecol.* **3**:305–326.
46. **Zuo, Y., and R. D. Jones.** 1997. Photochemistry of natural dissolved organic matter in lakes and wetland waters—production of carbon monoxide. *Water Res.* **31**:850–858.