

1-1-1990

A new chloroplast protein is induced by growth on low CO₂ in *Chlamydomonas reinhardtii*

Catherine B. Mason
Louisiana State University

Livingston J. Manuel
Louisiana State University

James V. Moroney
Louisiana State University

Follow this and additional works at: https://repository.lsu.edu/biosci_pubs

Recommended Citation

Mason, C., Manuel, L., & Moroney, J. (1990). A new chloroplast protein is induced by growth on low CO₂ in *Chlamydomonas reinhardtii*. *Plant Physiology*, 93 (2), 833-836. <https://doi.org/10.1104/pp.93.2.833>

This Article is brought to you for free and open access by the Department of Biological Sciences at LSU Scholarly Repository. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Scholarly Repository. For more information, please contact ir@lsu.edu.

Communication

A New Chloroplast Protein Is Induced by Growth on Low CO₂ in *Chlamydomonas reinhardtii*¹

Catherine B. Mason, Livingston J. Manuel, and James V. Moroney*

Department of Botany, Louisiana State University, Baton Rouge, Louisiana 70803

ABSTRACT

The biosynthesis of a 36 kilodalton polypeptide of *Chlamydomonas reinhardtii* was induced by photoautotrophic growth on low CO₂. Fractionation studies using the cell-wall-deficient strain of *C. reinhardtii*, CC-400, showed that this polypeptide was different from the low CO₂-induced periplasmic carbonic anhydrase. In addition, the 36 kilodalton polypeptide was found to be localized in intact chloroplasts isolated from low CO₂-adapting cultures. This protein may, in part, account for the different inorganic carbon uptake characteristics observed in chloroplasts isolated from high and low CO₂-grown *C. reinhardtii* cells.

The unicellular green alga *Chlamydomonas reinhardtii* can grow photoautotrophically at very low CO₂ concentrations due to its ability to concentrate C_i², internally to levels much higher than could be obtained by diffusion (2). This ability to concentrate C_i is a major photosynthetic adaptation that is seen in many unicellular algae, both eukaryotic and prokaryotic (1). However, the mechanism by which unicellular algae concentrate C_i is still poorly characterized.

In *C. reinhardtii*, the CO₂ concentrating mechanism is inducible, in that if the alga is grown on elevated CO₂ (1% [v/v] in air or higher), it exhibits a relatively low affinity for external C_i (2, 16). If, however, the alga is grown on low CO₂ concentrations, it acquires a very high affinity for C_i and has an extremely low CO₂ compensation point (2, 16). During adaptation to low CO₂, a number of proteins are preferentially made (3, 4, 10, 15). This induction requires both light and low CO₂ to be complete and involves an increase in transcription (19). To date, only the periplasmic carbonic anhydrase has been identified and characterized (1, 5, 19). Previous work has implied that a fundamental change in the C_i uptake characteristics of the chloroplast also occurs, as chloroplasts isolated from low CO₂ grown cells accumulate C_i to a greater extent than chloroplasts isolated from high CO₂ grown cells

¹ Supported by National Science Foundation grant DMB-8703462 and Louisiana Board of Regents Contract LEQSF (86-89)-RD-A-03.

² Abbreviations: C_i, inorganic carbon; low CO₂, air containing ambient (350 ppm) CO₂; high CO₂, air supplemented with CO₂ so that the final CO₂ concentration is 5% (v/v); Min medium, the phosphate buffered medium described by Sueoka (17) that contains no carbon source other than CO₂.

(12, 18). In this communication we describe an additional low CO₂ induced protein that is located within the chloroplast.

MATERIALS AND METHODS

Algal Strains and Culture Conditions

The wild-type strain of *Chlamydomonas reinhardtii*, 137 mt⁺, has been maintained in R. K. Togasaki's laboratory. The cell-wall-deficient mutant, CC-400 cw-15 mt⁺, was obtained from the Duke University *Chlamydomonas* Culture Collection. In liquid culture, the strains were grown in minimal medium (17), aerated with 5% CO₂ in air, and illuminated with 300 μE m⁻² s⁻¹ of white light. For ³⁵SO₄⁻² labeling, the algal cells were switched to minimal medium with one-tenth the normal amount of sulfur 48 h prior to the experiment.

³⁵S Labeling of Wild Type and CC-400 Cells

The labeling procedures used are similar to those previously published (10). In brief, harvested cells were resuspended in minimal medium lacking sulfate (Min-S), pelleted again by centrifugation, resuspended in Min-S (3 mL), and the Chl concentration was determined. The cells were adjusted to 25 μg Chl/mL and divided into six 150-mL flasks. Three flasks were bubbled with air and three with air supplemented with 5% CO₂. After a 30 min adaptation to the low and high CO₂ regimes, the cultures were labeled with carrier-free ³⁵SO₄⁻² (1000 Ci/mmol) in the light (3 h for wild-type cells and 4 h for CC-400).

Cell Fractionation

All steps were carried out at 0 to 4°C. After incubating the cells for the appropriate time, triplicate samples were combined, and 100 mL of cells were withdrawn from each treatment (air versus CO₂). The cells were harvested by centrifugation, washed twice with 50 mL ice-cold 20 mM Tris-HCl (7.5), resuspended in 250 μL of 20 mM Tris-HCl (pH 7.5), quick-frozen, and stored at -20°C.

Cells from the remaining 350 mL of each sample were collected by centrifugation, and the culture supernatant was brought to 70% saturation with (NH₄)₂SO₄. The (NH₄)₂SO₄ precipitates were collected by centrifugation, and the pellets were taken up in 1 mL 20 mM Tris-HCl (pH 7.5) and dialyzed overnight against 20 mM Tris-HCl (7.5). The resulting dialy-

sate (extracellular protein fraction) was quick-frozen and stored at -20°C .

The pelleted cells from the 350 mL harvest were washed twice with 100 mL ice-cold 20 mM Tris-HCl (pH 7.5) and recollected by centrifugation. The pellets were resuspended in 12 mL cold fractionation buffer (20 mM Tris-HCl [pH 7.5], 150 mM sucrose, 50 mM NaCl, 0.4 mM benzamidine, and 0.4 mM aminocaproic acid), and passed two times through a cell disruption bomb (PARR 4639) at 1800 psi. Unbroken cells were pelleted by centrifugation in a Sorvall Hb-4 rotor for 2 min at 670g (2000 rpm) and discarded. The supernatant was centrifuged at 6800g for 10 min and the pellet was taken up in 250 μL 20 mM Tris-HCl and quick-frozen (low speed pellet). The supernatant was then centrifuged for 2 h at 156,000g. The pellet was resuspended up in 250 μL 20 mM Tris-HCl (pH 7.5) and quick-frozen (high speed pellet). The supernatant was brought up to 70% saturation with $(\text{NH}_4)_2\text{SO}_4$, collected by centrifugation and the pellet was resuspended in 1 mL 20 mM Tris-HCl (pH 7.5) and dialyzed overnight (intracellular soluble protein fraction).

³⁵S-Labeling of Intact Chloroplasts

Chlamydomonas strain CC-400 cultures were grown in Min medium (3.0 L) on 12 h light/12 h dark regime to synchronize growth. Cells switched to minimal one-tenth sulfur media were harvested in the middle of the third light period by centrifugation, and the pellets were washed two times in 100 mL of 20 mM HEPES-KOH (pH 7.5) and were resuspended in 5 mL breaking buffer (300 mM sorbitol, 50 mM HEPES-KOH [pH 7.2], 2 mM Na-EDTA, 1 mM MgCl_2 , 1% BSA). The cell number was adjusted to 5×10^7 cells/mL and 20 mL aliquots were passed once through a cell disruption bomb (Parr 4639) at a pressure of 500 psi. The lysate was centrifuged in a Sorvall Hb-4 rotor for 2 min at 760g (2000 rpm) to pellet whole cells and intact chloroplasts. This pellet was resuspended in 2 mL breaking buffer and layered onto discontinuous Percoll gradients (20, 40, 60% Percoll) prepared according to Price and Reardon (14). A 15 min centrifugation of the gradients was carried out in a Sorvall Hb-4 rotor at 4200g. The 40 to 60% interface was collected and diluted fourfold with breaking buffer. Intact chloroplasts were collected by centrifugation at 670g for 1 min, and resuspended in 250 μL of 50 mM HEPES-KOH (pH 8.0), 0.3 M sorbitol. These intact chloroplasts retained galactosyltransferase activity and had similar physical and photosynthetic properties as *C. reinhardtii* chloroplasts previously reported (6, 9, 11, 12, 18).

Other Methods

Carbonic anhydrase was assayed as previously described (8). Chl concentrations were determined spectrophotometrically. Proteins estimations and SDS-PAGE was performed as previously described (10). The immunoblot assay was performed according to the protocol from Bio-Rad Laboratories except that 5% nonfat dry milk was used to block the nitrocellulose.

Materials

Goat anti-rabbit IgG(H+L) Horseradish Peroxidase conjugate and HRP color development reagent were purchased from Bio-Rad Laboratories. Carrier-free $\text{H}_2^{35}\text{SO}_4$ was obtained from ICN.

RESULTS

Chlamydomonas reinhardtii synthesizes at least four polypeptides preferentially when grown under limiting CO_2 (4, 10, 15). One of the labeled polypeptides had a molecular mass of 36 kD, similar to that of the 37 kD periplasmic carbonic anhydrase (5, 19). However, when fractionation studies were done with ³⁵S-labeled *C. reinhardtii* cells, the labeled 36 kD protein was found predominantly in the membrane fraction while the carbonic anhydrase activity was found mostly in the soluble protein fraction. Spalding and Jeffrey (15) had done fractionation studies on *cw-15 mt+* cells and reported membrane-associated proteins with molecular masses of 35 and 36 kD. This cell-wall deficient strain excretes the periplasmic carbonic anhydrase into the medium (5). We performed cell fractionation experiments on CC-400 cells (*cw-15 mt+*) and recovered both the periplasmic carbonic anhydrase from the cell supernatant and isolated membrane fractions. Labeled proteins of about 36 kD were present in both the extracellular protein fraction (Fig. 1, A and B, lanes 9 and 10), and the low speed membrane fraction (Fig. 1, A and B, lanes 1 and 2). However, only the extracellular protein was recognized by the antibody raised against the periplasmic carbonic anhydrase indicating that the membrane-associated, ³⁵S-labeled band was distinct from the periplasmic carbonic anhydrase (Fig. 1C). That this polypeptide is distinct from the periplasmic carbonic anhydrase is further supported by the observation that greater than 95% of the cells carbonic anhydrase activity is in the extracellular protein fraction (1316 units mg Chl^{-1}) from CC-400 cultures, and less than 1% of the activity was in the low speed pellet (5 units mg Chl^{-1}). Hence, at least five distinct proteins (indicated with asterisks in Fig. 1B), including a 36 kD polypeptide that is membrane bound, are specifically induced in a low CO_2 environment.

We further localized this protein by isolating intact chloroplasts from both low and high CO_2 -grown CC-400 cells. These cells were grown synchronously in Min $\frac{1}{10}$ S medium for 2 d on high CO_2 , and then labeled for 4 h on low or high CO_2 before isolating chloroplasts. Figure 2 shows that the 36 kD low- CO_2 -induced protein is only present in isolated chloroplasts from the low- CO_2 -adapted cultures. The soluble proteins induced by photoautotrophic growth on low CO_2 are not present in these chloroplast preparations (Fig. 2). Furthermore, an immunoblot of chloroplast proteins confirms the absence of the periplasmic carbonic anhydrase in these preparations (data not shown) and indicates that the 36 kD chloroplast protein is distinct from the periplasmic carbonic anhydrase. These data indicate that there is a low- CO_2 -induced protein present on intact chloroplasts that may play a role in the ability of low- CO_2 -grown cells to accumulate inorganic carbon.




Figure 1. ³⁵S-Labeled protein and immunoblot analysis of protein fractions from CC-400 cells. The labeled cells were (100 μg/lane) subjected to SDS-PAGE, stained with Coomassie blue (A), subjected to autoradiography (B), or probed with an antibody raised against the periplasmic CA (C). In lanes 1, 3, 5, 7, and 9 cells were grown on high CO₂, while lanes 2, 4, 6, 8, and 10 were from cells grown on low CO₂. Lanes 1 and 2, low speed membrane fraction; lanes 3 and 4, high speed membrane fraction; lanes 5 and 6, intracellular soluble protein fraction; lanes 7 and 8, total cellular protein fraction; lanes 9 and 10, extracellular protein fraction. The stained proteins seen in lanes 9 and 10 in A also include proteins from cells that have lysed during the initial harvest. These lysed cells do not incorporate ³⁵S, however (B).

DISCUSSION

Chlamydomonas reinhardtii possesses a CO₂ concentrating mechanism that involves at least five inducible proteins. These proteins include the periplasmic carbonic anhydrase (5, 19; Fig. 1, B and C, lanes 9 and 10), and soluble proteins with molecular masses of 46 and 44 kD (4, 10, 15; Fig. 1B, lanes 5, 6, 7, and 8). In addition, Spalding and Jeffrey (15) reported on low CO₂-induced membrane associated proteins with molecular masses of 36 and 21 kD which we also observed (lanes 1 and 2, Fig. 1B). Since the periplasmic carbonic anhydrase is cell wall associated, we fractionated a wild-type strain and a cell wall deficient strain of *C. reinhardtii* to determine whether the membrane associated 36 kD polypeptide was different than the periplasmic carbonic anhydrase. Carbonic anhydrase activity measurements indicated that the periplasmic carbonic anhydrase was largely in the soluble protein fraction while the ³⁵S labeled 36 kD protein was primarily in the low speed pellet. The low CO₂-induced 36 kD protein was also shown to be distinct from the 37 kD periplasmic carbonic anhydrase by immunoblots analysis of fractions from the CC-400 cells (Fig. 1C).

We have also shown that this 36 kD polypeptide is located in intact chloroplasts (Fig. 2), while the other proteins induced by growth on low CO₂ are not. Intact chloroplasts from *C. reinhardtii* have been isolated and partially characterized by a number of laboratories (6, 9, 11, 12, 18). The chloroplast preparations reported here are free of intact cells, nuclei, and other internal organelles as judged by electron microscopy. An intact plasma membrane is not present since the chloroplasts are permeable to P_i and DHAP. We cannot rule out the possibility of some contamination with plasma membrane fragments. In addition, these chloroplasts do retain the stromal enzymes rubisco (9, 11, 12) and carbonic anhydrase




Figure 2. ³⁵S-Labeled polypeptides of intact chloroplasts fractionated on Percoll gradients. Intact chloroplasts (100 μg protein) isolated from ³⁵S-labeled cells were subjected to SDS-PAGE and autoradiography. Lane 1, intact chloroplasts isolated from low-CO₂ grown cells; lane 2, intact chloroplasts isolated from high-CO₂ grown cells.

(8), and the envelope enzyme galactosyl transferase (11). The fact that the stromal enzymes are retained while the low CO₂ induced soluble polypeptides are lost supports the contention that those soluble proteins are not chloroplast localized.

The role of the 36 kD polypeptide is still unclear although the observation that its biosynthesis is induced along with the periplasmic carbonic anhydrase implies that it is important in CO₂ acquisition. In addition, this protein is not synthesized by the high-CO₂-requiring mutant *CIA-5* (13). Moroney *et al.* (12) and Sültemeyer *et al.* (18) have demonstrated that chloroplasts isolated from low CO₂-grown cells of *C. reinhardtii* have the ability to accumulate C_i to higher levels than chloroplasts from high CO₂-grown cells. Goyal and Tolbert (7) have made the same observation with chloroplasts from *Dunaliella*. The 36 kD polypeptide is the first report of a low-CO₂-inducible chloroplast protein. This protein may, in part, be responsible for the physiological differences in C_i uptake observed between chloroplasts isolated from low and high-CO₂-grown cells.

LITERATURE CITED

1. Aizawa K, Miyachi S (1986) Carbonic anhydrase and CO₂ concentrating mechanisms in microalgae and cyanobacteria. *FEMS Micro Rev* **39**: 215–233
2. Badger MR, Kaplan A, Berry JA (1980) Internal inorganic carbon pool of *Chlamydomonas reinhardtii*: Evidence for a carbon dioxide concentrating mechanism. *Plant Physiol* **66**: 407–413
3. Badour SS, Kim WK (1988) Detection of specific polypeptides in *Chlamydomonas segnis* adapted to atmospheric concentrations of CO₂, using a zwitterionic detergent. *Can J Bot* **66**: 1750–1754
4. Bailly J, Coleman JR (1988) Effect of CO₂ concentration on protein biosynthesis and carbonic anhydrase expression in *Chlamydomonas reinhardtii*. *Plant Physiol* **87**: 833–840
5. Coleman JR, Berry JA, Togasaki RK, Grossman AR (1984) Identification of extracellular carbonic anhydrase of *Chlamydomonas reinhardtii*. *Plant Physiol* **76**: 472–477
6. Goldschmidt-Clermont M, Malnoë P, Rochaix JD (1989) Preparation of *Chlamydomonas* chloroplasts for the *In vitro* impact of polypeptide precursors. *Plant Physiol* **89**: 15–18
7. Goyal A, Tolbert NE (1989) Uptake of inorganic carbon by isolated chloroplasts from air-adapted *Dunaliella*. *Plant Physiol* **89**: 1264–1269
8. Husic HD, Kitayama M, Togasaki RK, Moroney JV, Morris KL, Tolbert NE (1989) Identification of intracellular carbonic anhydrase in *Chlamydomonas reinhardtii* which is distinct from the periplasmic form of the enzyme. *Plant Physiol* **89**: 904–909
9. Klein U, Chen C, Gibbs M (1983) Photosynthetic properties of chloroplasts from *Chlamydomonas reinhardtii*. *Plant Physiol* **72**: 488–491
10. Manuel LJ, Moroney JV (1988) Inorganic carbon accumulation in *Chlamydomonas reinhardtii*: new proteins are made during adaptation to low CO₂. *Plant Physiol* **88**: 491–496
11. Mendiola-Morgenthaler L, Leu S, Boschetti A (1985) Isolation of biochemically active chloroplasts from *Chlamydomonas*. *Plant Sci* **38**: 33–39
12. Moroney JV, Kitayama M, Togasaki RK, Tolbert NE (1987) Evidence for inorganic carbon transport by intact chloroplasts of *Chlamydomonas reinhardtii*. *Plant Physiol* **83**: 460–463
13. Moroney JV, Husic HD, Tolbert NE, Kitayama M, Manuel LJ, Togasaki RK (1989) Isolation and characterization of a mutant of *Chlamydomonas reinhardtii* deficient in the CO₂ concentrating mechanism. *Plant Physiol* **89**: 897–903
14. Price CA, Reardon EM (1982) Isolation of chloroplasts for protein synthesis from spinach and *Euglena gracilis* by centrifugation in silica sols. In M Edelman, RB Hallick, NA Chua, eds, *Methods in Chloroplast Molecular Biology*. Elsevier, Amsterdam, pp 189–209
15. Spalding MH, Jeffrey M (1989) Membrane-associated polypeptides induced in *Chlamydomonas* by limiting CO₂ concentrations. *Plant Physiol* **89**: 133–137
16. Spalding MH, Spreitzer RJ, Ogren WL (1983) Carbonic anhydrase deficient mutant of *Chlamydomonas reinhardtii* requires elevated carbon dioxide concentrations for photoautotrophic growth. *Plant Physiol* **73**: 268–272
17. Sueoka N (1960) Mitotic replication of deoxyribonucleic acids in *Chlamydomonas reinhardtii*. *Proc Natl Acad Sci USA* **46**: 83–91
18. Sültemeyer DF, Klöck G, Kreuberg K, Fock H (1988) Photosynthesis and apparent affinity for dissolved inorganic carbon by cells and protoplasts of *Chlamydomonas reinhardtii* grown at high and low CO₂ concentrations. *Planta* **176**: 256–260
19. Toguri T, Muto S, Miyachi S (1986) Biosynthesis and intracellular processing of carbonic anhydrase in *Chlamydomonas reinhardtii*. *Eur J Biochem* **158**: 443–450