

9-11-2009

Search for electron antineutrino appearance at the $\Delta m^2 \square 1 \text{eV}^2$ Scale

A. A. Aguilar-Arevalo
Universidad Nacional Autónoma de México

C. E. Anderson
Yale University

S. J. Brice
Fermi National Accelerator Laboratory

B. C. Brown
Fermi National Accelerator Laboratory

L. Bugel
Columbia University

See next page for additional authors

Follow this and additional works at: https://repository.lsu.edu/physics_astronomy_pubs

Recommended Citation

Aguilar-Arevalo, A., Anderson, C., Brice, S., Brown, B., Bugel, L., Conrad, J., Djurcic, Z., Fleming, B., Ford, R., Garcia, F., Garvey, G., Gonzales, J., Grange, J., Green, C., Green, J., Imlay, R., Johnson, R., Karagiorgi, G., Katori, T., Kobilarcik, T., Linden, S., Louis, W., Mahn, K., Marsh, W., Mauger, C., McGary, V., Metcalf, W., Mills, G., Moore, C., Mousseau, J., Nelson, R., Nienaber, P., & Nowak, J. (2009). Search for electron antineutrino appearance at the $\Delta m^2 \square 1 \text{eV}^2$ Scale. *Physical Review Letters*, 103 (11) <https://doi.org/10.1103/PhysRevLett.103.111801>

This Article is brought to you for free and open access by the Department of Physics & Astronomy at LSU Scholarly Repository. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Scholarly Repository. For more information, please contact ir@lsu.edu.

Authors

A. A. Aguilar-Arevalo, C. E. Anderson, S. J. Brice, B. C. Brown, L. Bugel, J. M. Conrad, Z. Djurcic, B. T. Fleming, R. Ford, F. G. Garcia, G. T. Garvey, J. Gonzales, J. Grange, C. Green, J. A. Green, R. Imlay, R. A. Johnson, G. Karagiorgi, T. Katori, T. Kobilarcik, S. K. Linden, W. C. Louis, K. B.M. Mahn, W. Marsh, C. Mauger, V. T. McGary, W. Metcalf, G. B. Mills, C. D. Moore, J. Mousseau, R. H. Nelson, P. Nienaber, and J. A. Nowak



MIT Open Access Articles

Search for Electron Antineutrino Appearance at the Delta $m^2 \sim 1 \text{ eV}^2$ Scale

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation	MiniBooNE Collaboration et al. "Search for Electron Antineutrino Appearance at the Delta $m^2 \sim 1 \text{ eV}^2$ Scale." Physical Review Letters 103.11 (2009): 111801. © 2009 American Physical Society.
As Published	http://dx.doi.org/10.1103/PhysRevLett.103.111801
Publisher	American Physical Society
Version	Final published version
Citable link	http://hdl.handle.net/1721.1/54785
Terms of Use	Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.

Search for Electron Antineutrino Appearance at the $\Delta m^2 \sim 1 \text{ eV}^2$ Scale

A. A. Aguilar-Arevalo,¹² C. E. Anderson,¹⁵ S. J. Brice,⁵ B. C. Brown,⁵ L. Bugel,⁴ J. M. Conrad,¹¹ Z. Djurcic,⁴ B. T. Fleming,¹⁵ R. Ford,⁵ F. G. Garcia,⁵ G. T. Garvey,⁹ J. Gonzales,⁹ J. Grange,⁶ C. Green,^{5,9} J. A. Green,^{8,9} R. Imlay,¹⁰ R. A. Johnson,² G. Karagiorgi,¹¹ T. Katori,^{8,11} T. Kobilarcik,⁵ S. K. Linden,¹⁵ W. C. Louis,⁹ K. B. M. Mahn,⁴ W. Marsh,⁵ C. Mauger,⁹ V. T. McGary,¹¹ W. Metcalf,¹⁰ G. B. Mills,⁹ C. D. Moore,⁵ J. Mousseau,⁶ R. H. Nelson,³ P. Nienaber,¹⁴ J. A. Nowak,¹⁰ B. Osmanov,⁶ Z. Pavlovic,⁹ D. Perevalov,¹ C. C. Polly,^{7,8} H. Ray,^{6,9} B. P. Roe,¹³ A. D. Russell,⁵ M. H. Shaevitz,⁴ M. Sorel,^{4,*} J. Spitz,¹⁵ I. Stancu,¹ R. J. Stefanski,⁵ R. Tayloe,⁸ M. Tzanov,³ R. G. Van de Water,⁹ M. O. Wascko,^{10,†} D. H. White,⁹ M. J. Wilking,³ G. P. Zeller,^{4,9} and E. D. Zimmerman³

(MiniBooNE Collaboration)

¹University of Alabama, Tuscaloosa, Alabama 35487, USA

²University of Cincinnati, Cincinnati, Ohio 45221, USA

³University of Colorado, Boulder, Colorado 80309, USA

⁴Columbia University, New York, New York 10027, USA

⁵Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA

⁶University of Florida, Gainesville, Florida 32611, USA

⁷University of Illinois, Urbana, Illinois 61801, USA

⁸Indiana University, Bloomington, Indiana 47405, USA

⁹Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA

¹⁰Louisiana State University, Baton Rouge, Louisiana 70803, USA

¹¹Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

¹²Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Distrito Federal 04510, México

¹³University of Michigan, Ann Arbor, Michigan 48109, USA

¹⁴Saint Mary's University of Minnesota, Winona, Minnesota 55987, USA

¹⁵Yale University, New Haven, Connecticut 06520, USA

(Received 13 April 2009; published 11 September 2009)

The MiniBooNE Collaboration reports initial results from a search for $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillations. A signal-blind analysis was performed using a data sample corresponding to 3.39×10^{20} protons on target. The data are consistent with background prediction across the full range of neutrino energy reconstructed assuming quasielastic scattering, $200 < E_\nu^{\text{QE}} < 3000$ MeV: 144 electronlike events have been observed in this energy range, compared to an expectation of 139.2 ± 17.6 events. No significant excess of events has been observed, both at low energy, 200–475 MeV, and at high energy, 475–1250 MeV. The data are inconclusive with respect to antineutrino oscillations suggested by data from the Liquid Scintillator Neutrino Detector at Los Alamos National Laboratory.

DOI: 10.1103/PhysRevLett.103.111801

PACS numbers: 14.60.Lm, 14.60.Pq, 14.60.St

Motivated by the LSND observation of an excess of $\bar{\nu}_e$ events in a $\bar{\nu}_\mu$ beam [1], the MiniBooNE Collaboration has previously performed a search for $\nu_\mu \rightarrow \nu_e$ oscillations, the results of which showed no evidence of an excess of ν_e events for neutrino energies above 475 MeV [2,3]. Assuming no *CPT* or *CP* violation, the results exclude the LSND excess interpreted as two-neutrino oscillations at $\Delta m^2 \sim 0.1\text{--}100 \text{ eV}^2$ at 98% C.L. Similarly, the KARMEN experiment [4] has performed a direct search for $\bar{\nu}_e$ appearance, and has placed a limit independent of any *CPT* or *CP* violation assumption. However, a joint analysis of KARMEN and LSND results shows high compatibility [5,6]. A corresponding $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillation search has been performed at MiniBooNE and is presented in this Letter. This search serves as another direct test of

LSND and provides complementary information to that of KARMEN, having sensitivity to the lower Δm^2 oscillations allowed by the joint KARMEN-LSND analysis [7]. It should be noted that, in a simple two-neutrino oscillation model where *CPT* conservation is imposed, oscillation probabilities (mixing amplitudes and mass-squared differences) for neutrinos and antineutrinos cannot be different. Therefore, the oscillation search presented here is a direct search for existence of nonstandard oscillations where *CPT* is violated, or effectively violated.

Despite having observed no evidence for oscillations above 475 MeV, the MiniBooNE $\nu_\mu \rightarrow \nu_e$ search observed a 3.0σ excess of electronlike events at low energy, between 200–475 MeV [3]. Although the excess is incompatible with LSND-type oscillations, several hypotheses [8–13],

including sterile neutrino oscillations with CP violation, anomaly-mediated neutrino-photon coupling, and others, have been proposed that provide a possible explanation for the excess itself, and, in some cases, offer the possibility of reconciling the MiniBooNE ν_e excess with the LSND $\bar{\nu}_e$ excess. These phenomenological interpretations have provided additional motivation for an antineutrino appearance search at MiniBooNE.

The analysis presented in this Letter mirrors the blind search performed in neutrino mode [2]. It employs a two-neutrino oscillation model, where only $\bar{\nu}_\mu$ present in the MiniBooNE beam are allowed to oscillate into $\bar{\nu}_e$, at $\Delta m^2 \sim 0.1\text{--}100\text{ eV}^2$. Given that no evidence of ν_μ oscillations was observed in high-purity, high-statistics searches in neutrino mode [2,14], the analysis further assumes no $\bar{\nu}_\mu$ disappearance and no ν_μ oscillations. In addition, no contribution from the observed neutrino mode low energy excess has been accounted for in the antineutrino prediction.

The antineutrino flux [15] is produced by 8 GeV protons incident on a beryllium target. Negatively charged mesons produced in p -Be interactions are focused in the forward direction with the use of a toroidal magnetic field, and subsequently decay primarily into $\bar{\nu}_\mu$. In antineutrino mode, a large neutrino contamination (ν_μ and ν_e) of 15.9% is expected in the flux viewed by the detector, compared to 5.9% in neutrino mode. The intrinsic $\bar{\nu}_e$ and ν_e content is only 0.4% and 0.2%, respectively, coming primarily from $\pi \rightarrow \mu$ and K decays. The $\bar{\nu}_\mu$ flux peaks at $\sim 400\text{ MeV}$ and has a mean energy of $\sim 600\text{ MeV}$. See [15] for more details.

A detailed description of the MiniBooNE detector is available in [16]. The detector location was chosen to satisfy $L[\text{m}]/E[\text{MeV}] \sim 1$, similar to that of LSND, thus maximizing sensitivity to oscillations at $\Delta m^2 \sim 1\text{ eV}^2$. The detector is filled with pure mineral oil (CH_2). Neutrino interactions in the detector produce final state electrons or muons, which produce scintillation and Cherenkov light detected by photomultiplier tubes (PMTs) that line the interior of the detector. The simulation of light incident on the PMTs takes into account decays and strong and electromagnetic reinteractions in the detector, and includes processes that were added in the final ν_e appearance analysis [3]. The M_A^{QE} appearing in the nucleon axial vector form factor, and the Pauli blocking parameter κ used to parametrize neutrino quasielastic scattering on carbon, were adjusted by fits to MiniBooNE data, as were the coherent pion cross sections [17,18]. The M_A^{QE} and κ values of $1.23 \pm 0.08\text{ GeV}$ and 1.022 ± 0.021 , respectively, were used in this analysis. Two additional parameters, $M_A^{\text{QE},H} = 1.13 \pm 0.10\text{ GeV}$ and $M_A^{1\pi,H} = 1.10 \pm 0.10\text{ GeV}$, were introduced in the analysis to parametrize antineutrino quasielastic scattering on hydrogen and single pion production on hydrogen. These processes have a non-negligible contribution in antineutrino running mode, where roughly 25% of the

antineutrino quasielastic scatters are on hydrogen rather than carbon.

The detector cannot differentiate (on an event-by-event basis) a ν_μ from a $\bar{\nu}_\mu$ interaction, or a ν_e from a $\bar{\nu}_e$ interaction. Therefore, the reconstruction and selection requirements for $\bar{\nu}_e$ -induced charged-current quasielastic (CCQE) events, which is the characteristic signature of any possible signal from $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillations, are identical to those of the final neutrino mode analysis [3].

To provide a constraint on $\bar{\nu}_e$ candidate events, a $\bar{\nu}_\mu$ CCQE sample is also formed by looking for events with a muonlike Cherenkov ring and a cluster of delayed PMT hits from the decay of the muon into an electron. The first cluster of PMT hits (muon subevent) is required to have more than 200 inner detector PMT hits, and no more than six outer (veto) PMT hits. A maximum of 200 inner detector and six veto PMT hits are required for the second subevent (decay electron), and a minimum time cut of 1000 ns between the first and second subevents is required to ensure PMT stability for proper charge response. After reconstruction, the first subevent vertex and the track end point under the muon hypothesis are required to occur within the fiducial volume. The neutrino energy reconstructed from the outgoing muon energy and angle, E_ν^{QE} , is required to satisfy $E_\nu^{\text{QE}} > 150\text{ MeV}$. A cut on the separation distance between the muon and decay electron vertices as a function of reconstructed energy of the muon is also applied to provide rejection against backgrounds, mostly from charged-current (CC) π^+ interactions. For more details on the reconstruction method, see [19].

The oscillation parameters are extracted from a combined fit to $\bar{\nu}_e$ CCQE and $\bar{\nu}_\mu$ CCQE event distributions, following [3]. This fit method takes advantage of strong flux and cross-section correlations among the $\bar{\nu}_e$ CCQE and $\bar{\nu}_\mu$ CCQE event samples, since any possible $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ signal, as well as some $\bar{\nu}_e$ backgrounds, interact through the same process as $\bar{\nu}_\mu$ CCQE events, and are related to $\bar{\nu}_\mu$ CCQE events through the same π^+ or π^- decay chain at production. These correlations enter through the off-diagonal elements of the covariance matrix used in the χ^2 calculation, relating the contents of the bins of the $\bar{\nu}_e$ CCQE and $\bar{\nu}_\mu$ CCQE distributions. This procedure maximizes the sensitivity to $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillations when systematic uncertainties are included [20].

A sample of 14 107 data events passing $\bar{\nu}_\mu$ CCQE selection requirements is used in the analysis. This sample is compared to a Monte Carlo prediction which has been corrected to match the observed $\bar{\nu}_\mu$ CCQE data through a normalization factor of 1.22 applied to events from π^- decays in the beam, and 0.93 applied to events from π^+ decays in the beam. These normalization factors are extracted from a fit to the angular distributions of the outgoing μ^+ and μ^- in $\bar{\nu}_\mu$ and ν_μ CCQE interactions [7]. These two factors result in an overall 15% normalization correction which is covered by flux and cross-section un-

certainties. The same normalization correction is also applied to all possible signal events which share the same parent (π^-) as $\bar{\nu}_\mu$ CCQE events. The normalization correction is accounted for in the oscillation fit by a reduction in the quoted effective degrees of freedom (d.o.f.) by one unit. After correction, the sample contains 95% $\bar{\nu}_\mu$ and ν_μ produced in pion decays and 2.4% $\bar{\nu}_\mu$ and ν_μ produced in kaon decays. The neutrino content of the sample is 22%. The majority of events (71%) are true CCQE interactions, with CC π^\pm interactions being the dominant source of background (20%). This sample is included in the $\bar{\nu}_e$ appearance fits as a function of 8 bins of reconstructed neutrino energy, E_ν^{QE} , ranging from 0 to 1900 MeV.

Table I shows the number of predicted $\bar{\nu}_e$ CCQE background events for different ranges of E_ν^{QE} . The background estimates include both antineutrino and neutrino events, the latter representing $\sim 44\%$ of the total. The predicted backgrounds to the $\bar{\nu}_e$ CCQE sample are constrained by internal measurements at MiniBooNE. These measurements use event samples from regions in reconstructed kinematic variables where any possible signal from $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ is negligible, in order to preserve blindness. The constrained backgrounds include NC π^0 events, $\Delta \rightarrow N\gamma$ radiative events, and events from interactions outside the detector. The NC π^0 background events are adjusted in bins of π^0 momentum according to a direct π^0 rate measurement in antineutrino mode, following [17], which uses events reconstructed near the π^0 mass peak. The size of the applied correction to the total NC π^0 rate is less than 10%. The $\Delta \rightarrow N\gamma$ rate is indirectly constrained, being related to the measured π^0 rate through a branching fraction and final state interaction correction. The rate of backgrounds from external interactions is constrained through a direct measurement at MiniBooNE, using a separate event sample where the rate of external interaction events is enhanced.

TABLE I. The expected number of events for different E_ν^{QE} ranges (in MeV) from all of the backgrounds in the $\bar{\nu}_e$ appearance analysis and for the LSND central expectation (0.26% oscillation probability) of $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillations, for 3.39×10^{20} POT.

Process	200–300	300–475	475–1250
$\bar{\nu}_\mu$ CCQE	1.3	1.6	1.2
NC π^0	14.4	10.2	7.2
NC $\Delta \rightarrow N\gamma$	1.7	4.9	2.0
External events	2.2	2.5	1.9
Other $\bar{\nu}_\mu$	2.0	1.8	2.2
$\bar{\nu}_e$ from μ^\pm decay	2.3	5.9	17.1
$\bar{\nu}_e$ from K^\pm decay	1.4	3.8	11.7
$\bar{\nu}_e$ from K_L^0 decay	0.8	2.4	13.1
Other $\bar{\nu}_e$	0.5	0.6	1.21
Total background	26.7	33.6	57.8
0.26% $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$	0.6	3.7	12.6

Other backgrounds from misidentified ν_μ or $\bar{\nu}_\mu$ receive the $\bar{\nu}_\mu$ CCQE normalization correction according to their parentage at production (π^+ or π^-). Intrinsic ν_e and $\bar{\nu}_e$ events from the $\pi \rightarrow \mu$ decay chain also receive this normalization.

Systematic uncertainties are determined by considering the effects on the $\bar{\nu}_\mu$ and $\bar{\nu}_e$ CCQE rate prediction of variations of fundamental parameters within their associated uncertainty. These include uncertainties on the flux estimate, including beam modeling and hadron production at the target, uncertainties on neutrino cross sections, most of which are determined by *in situ* cross-section measurements at MiniBooNE or other experimental or theoretical sources, and uncertainties on detector modeling and reconstruction. By considering the variation from each source of systematic uncertainty on the $\bar{\nu}_e$ CCQE signal, background, and $\bar{\nu}_\mu$ CCQE prediction as a function of E_ν^{QE} , a covariance matrix in bins of E_ν^{QE} is constructed, which includes correlations between $\bar{\nu}_e$ CCQE (signal and background) and $\bar{\nu}_\mu$ CCQE. This covariance matrix is used in the χ^2 calculation of the oscillation fit.

Figure 1 (top) shows the E_ν^{QE} distribution for $\bar{\nu}_e$ CCQE observed data and background. A total of 144 events pass the $\bar{\nu}_e$ event selection requirements with $200 < E_\nu^{\text{QE}} < 3000$ MeV. The data agree with the background prediction within systematic and statistical uncertainties. Figure 1 (bottom) shows the event excess as a function of E_ν^{QE} . Also shown are expectations from the best $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillation parameters returned by the fit and from two other sets of neutrino oscillation parameters from the LSND allowed region [1]. The best oscillation fit for $200 < E_\nu^{\text{QE}} < 3000$ MeV corresponds to $(\Delta m^2, \sin^2 2\theta) = (4.42 \text{ eV}^2, 0.004)$, and has a χ^2 of 18.2 for 16 d.o.f., cor-




FIG. 1 (color online). Top: The E_ν^{QE} distribution for $\bar{\nu}_e$ CCQE data (points with statistical errors) and background (histogram with unconstrained systematic errors). Bottom: The event excess as a function of E_ν^{QE} . Also shown are the expectations from the best oscillation fit and from neutrino oscillation parameters in the LSND allowed region. The error bars include both statistical and systematic errors.

TABLE II. The number of data, background, and excess events in the $\bar{\nu}_e$ analysis for different E_ν^{QE} ranges. The corresponding numbers from the ν_e analysis [3] are on the right. The uncertainties include both statistical and constrained systematic errors.

Event sample	$\bar{\nu}_e$ analysis (3.39×10^{20} POT)	ν_e analysis [3] (6.46×10^{20} POT)
200–475 MeV		
Data	61	544
Background	61.5 ± 11.7	415.2 ± 43.4
Excess	-0.5 ± 11.7 (-0.04σ)	128.8 ± 43.4 (3.0σ)
475–1250 MeV		
Data	61	408
Background	57.8 ± 10.0	385.9 ± 35.7
Excess	3.2 ± 10.0 (0.3σ)	22.1 ± 35.7 (0.6σ)

responding to a χ^2 probability of 31%. The null fit yields $\chi^2/\text{d.o.f.} = 24.5/18$, with a χ^2 probability of 14%. A fit to $475 < E_\nu^{\text{QE}} < 3000$ MeV returns similar best-fit oscillation parameters, $(\Delta m^2, \sin^2 2\theta) = (4.42 \text{ eV}^2, 0.005)$, with $\chi^2/\text{d.o.f.} = 15.9/13$ and a χ^2 probability of 25%. The null fit to $475 < E_\nu^{\text{QE}} < 3000$ MeV yields $\chi^2/\text{d.o.f.} = 22.2/15$, with a χ^2 probability of 10%. The number of data, background, and excess events for different E_ν^{QE} ranges are summarized in Table II. No significant event excess is observed for $E_\nu^{\text{QE}} > 475$ MeV. Furthermore, no significant excess is observed for $E_\nu^{\text{QE}} < 475$ MeV, to be compared to a 3.0σ excess observed for $200 < E_\nu^{\text{QE}} < 475$ MeV in the ν_e appearance analysis [3].

The $\bar{\nu}_e$ data also exhibit reasonable agreement with predicted background in other reconstructed kinematic variables. Figure 2 shows the observed and predicted event distributions as functions of reconstructed Q^2 and $\cos(\theta)$ for $200 < E_\nu^{\text{QE}} < 3000$ MeV. Q^2 is determined from the energy of the outgoing lepton and its scattering angle with respect to the incident neutrino direction (θ) assuming CCQE scattering. Also shown in the figures are the predicted distributions from NC π^0 and $\Delta \rightarrow N\gamma$ backgrounds, which are events with a photon in the final state. The null χ^2 values from these comparisons are both acceptable, at $\chi^2/\text{d.o.f.} = 10.6/11$ and $\chi^2/\text{d.o.f.} = 8.4/11$ for Q^2 and $\cos(\theta)$, respectively.

The absence of a significant excess allows MiniBooNE to place a limit on $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillations as shown in Fig. 3. The bottom panel of the figure shows the MiniBooNE limits obtained from fits to events with $E_\nu^{\text{QE}} > 200$ MeV and $E_\nu^{\text{QE}} > 475$ MeV. Each 90% C.L. limit on $\sin^2 2\theta$ is obtained by a single-sided raster scan of the parameter space, where a $\Delta\chi^2 = \chi_{\text{limit}}^2 - \chi_{\text{best fit}}^2 < 1.64$ cut is applied for each slice in Δm^2 . The two limits are in agreement, with the one obtained for $E_\nu^{\text{QE}} > 200$ MeV placing a stronger bound for low Δm^2 oscillations, due to its slightly better sensitivity in that region (see top panel of Fig. 3). At higher Δm^2 values, both limits approach the corresponding sensitivities of the experiment, but at lower Δm^2 both limits are noticeably worse due to the observed data fluctuation between $475 < E_\nu^{\text{QE}} < 675$ MeV. The significance of that fluctuation in the $475 < E_\nu^{\text{QE}} < 675$ MeV range is 2.8σ (statistical \oplus constrained systematic).

Following [2], a secondary analysis based on boosted decision trees (BDT) has been performed and used as a cross-check for the oscillation analysis in the energy region $E_\nu^{\text{QE}} > 500$ MeV, where the BDT analysis is not dominated by systematic uncertainties. No significant excess of events is observed with the BDT analysis, yielding the limit shown in the top panel of Fig. 3. Although the limit from the BDT analysis is not as stringent as the main result discussed above, the two analyses are complementary and yield consistent results.

Following [2], a secondary analysis based on boosted decision trees (BDT) has been performed and used as a cross-check for the oscillation analysis in the energy region $E_\nu^{\text{QE}} > 500$ MeV, where the BDT analysis is not dominated by systematic uncertainties. No significant excess of events is observed with the BDT analysis, yielding the limit shown in the top panel of Fig. 3. Although the limit from the BDT analysis is not as stringent as the main result discussed above, the two analyses are complementary and yield consistent results.




FIG. 2 (color). The Q^2 (top panel) and $\cos(\theta)$ (bottom panel) distributions for data (points with statistical errors) and backgrounds (histogram with constrained systematic errors) for $E_\nu^{\text{QE}} > 200$ MeV. Also shown are the expected distributions from intrinsic $\bar{\nu}_e$ and ν_e , and NC π^0 and $\Delta \rightarrow N\gamma$ backgrounds.




FIG. 3 (color online). Top: MiniBooNE 90% C.L. limit (solid black line) and sensitivity (dashed black line) for events with $E_{\nu}^{\text{QE}} > 200$ MeV, within a two-neutrino $\bar{\nu}_{\mu} \rightarrow \bar{\nu}_e$ oscillation model. Also shown is the sensitivity for $E_{\nu}^{\text{QE}} > 475$ MeV (dashed gray line), and the limit from the BDT analysis (solid blue area) for $E_{\nu}^{\text{QE}} > 500$ MeV. Bottom: Limits from MiniBooNE for $E_{\nu}^{\text{QE}} > 200$ MeV and $E_{\nu}^{\text{QE}} > 475$ MeV, KARMEN [4], and Bugey [22]. The island contour in the bottom left corner is a lower $\sin^2 2\theta$ limit from a fit to $E_{\nu}^{\text{QE}} > 475$ MeV, excluding the points left of the line at 90% C.L. The MiniBooNE and Bugey curves are one-sided limits for $\sin^2 2\theta$ corresponding to $\Delta\chi^2 = 1.64$, while the KARMEN curve is a “unified approach” 2D contour. The shaded areas show the 90% and 99% C.L. LSND allowed regions.

In summary, MiniBooNE observes no significant excess of $\bar{\nu}_e$ events in the energy region $E_{\nu}^{\text{QE}} > 200$ MeV, for a data sample corresponding to 3.39×10^{20} protons on target (POT). Thus, with current statistics, MiniBooNE places a limit on two-neutrino $\bar{\nu}_{\mu} \rightarrow \bar{\nu}_e$ oscillations shown by the black line in Fig. 3. The result is inconclusive with respect to small amplitude mixing suggested by the LSND data, but more antineutrino data, which are currently being collected, will provide additional information. More constraints may also be provided by the off-axis NuMI beam data collected in MiniBooNE [21]. Interestingly, MiniBooNE observes no significant excess of $\bar{\nu}_e$ events

in the low energy region $200 < E_{\nu}^{\text{QE}} < 475$ MeV. The absence of an excess at low energy in antineutrino mode should help distinguish between several hypotheses suggested as explanations for the low energy excess observed in neutrino mode.

We acknowledge the support of Fermilab, the Department of Energy, and the National Science Foundation, and we acknowledge Los Alamos National Laboratory for LDRD funding. We also acknowledge the use of CONDOR software for the analysis of the data.

*Present address: IFIC, Universidad de Valencia and CSIC, Valencia 46071, Spain.

†Present address: Imperial College; London SW7 2AZ, United Kingdom.

- [1] C. Athanassopoulos *et al.*, Phys. Rev. Lett. **75**, 2650 (1995); **77**, 3082 (1996); **81**, 1774 (1998); A. Aguilar *et al.*, Phys. Rev. D **64**, 112007 (2001).
- [2] A. Aguilar-Arevalo *et al.*, Phys. Rev. Lett. **98**, 231801 (2007).
- [3] A. A. Aguilar-Arevalo *et al.*, Phys. Rev. Lett. **102**, 101802 (2009).
- [4] B. Armbruster *et al.*, Phys. Rev. D **65**, 112001 (2002).
- [5] E. D. Church *et al.*, Phys. Rev. D **66**, 013001 (2002).
- [6] A. A. Aguilar-Arevalo *et al.*, Phys. Rev. D **78**, 012007 (2008).
- [7] A. A. Aguilar-Arevalo *et al.*, Addendum to the MiniBooNE Run Plan: MiniBooNE Physics in 2006, <http://www-boone.fnal.gov/publicpages/loi.ps.gz>.
- [8] M. Sorel *et al.*, Phys. Rev. D **70**, 073004 (2004); G. Karagiorgi *et al.*, Phys. Rev. D **75**, 013011 (2007); A. Melchiorri *et al.*, J. Cosmol. Astropart. Phys. 01 (2009) 036; M. Maltoni and T. Schwetz, Phys. Rev. D **76**, 093005 (2007).
- [9] J. A. Harvey *et al.*, Phys. Rev. Lett. **99**, 261601 (2007); Phys. Rev. D **77**, 085017 (2008).
- [10] H. Pas *et al.*, Phys. Rev. D **72**, 095017 (2005).
- [11] T. Goldman *et al.*, Phys. Rev. D **75**, 091301 (2007).
- [12] A. E. Nelson and J. Walsh, Phys. Rev. D **77**, 033001 (2008).
- [13] V. A. Kostelecky and M. Mewes, Phys. Rev. D **69**, 016005 (2004); T. Katori *et al.*, Phys. Rev. D **74**, 105009 (2006).
- [14] A. A. Aguilar-Arevalo *et al.*, Phys. Rev. Lett. **103**, 061802 (2009).
- [15] A. A. Aguilar-Arevalo *et al.*, Phys. Rev. D **79**, 072002 (2009).
- [16] A. A. Aguilar-Arevalo *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **599**, 28 (2009).
- [17] A. A. Aguilar-Arevalo *et al.*, Phys. Lett. B **664**, 41 (2008).
- [18] A. A. Aguilar-Arevalo *et al.*, Phys. Rev. Lett. **100**, 032301 (2008).
- [19] R. B. Patterson *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **608**, 206 (2009).
- [20] D. W. Schmitz, Report No. FERMILAB-THESIS-2008-26.
- [21] P. Adamson *et al.*, Phys. Rev. Lett. **102**, 211801 (2009).
- [22] B. Achkar *et al.*, Nucl. Phys. **B434**, 503 (1995).