

1-22-2021

## Covariance of the extended holonomy

Rodolfo Gambini  
*Facultad de Ciencias*

Jorge Pullin  
*Louisiana State University*

Aureliano Skirzewski  
*Facultad de Ciencias*

Follow this and additional works at: [https://repository.lsu.edu/physics\\_astronomy\\_pubs](https://repository.lsu.edu/physics_astronomy_pubs)

---

### Recommended Citation

Gambini, R., Pullin, J., & Skirzewski, A. (2021). Covariance of the extended holonomy. *Classical and Quantum Gravity*, 38 (2) <https://doi.org/10.1088/1361-6382/abc667>

This Article is brought to you for free and open access by the Department of Physics & Astronomy at LSU Scholarly Repository. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Scholarly Repository. For more information, please contact [ir@lsu.edu](mailto:ir@lsu.edu).

# Covariance of the extended holonomy

Rodolfo Gambini<sup>1</sup>, Jorge Pullin<sup>2</sup>, Aureliano Skirzewski<sup>1</sup>

*1. Instituto de Física, Facultad de Ciencias,  
Iguá 4225, esq. Mataojo, 11400 Montevideo, Uruguay.*

*2. Department of Physics and Astronomy,  
Louisiana State University, Baton Rouge, LA 70803-4001*

It has been pointed out that the holonomy of generic extended loops is not gauge covariant. We show how to define a family of extended loops for which previous criticism does not apply. We also give sufficient conditions that extended loops must satisfy in order to yield covariant holonomies. This makes a quantum representation for Yang–Mills theories and gravity based on extended loops viable.

## I. INTRODUCTION

The use of loop based variables to study gauge theories can be traced all the way back to Faraday. In the context of Yang–Mills theory holonomies have been widely used to analyze the quantization, both in the continuum and in the lattice in gauge invariant terms. They have also been used to base a complete quantum representation in terms of loops, the loop representation, again both for Yang–Mills theories and gravity (see [1] and references therein). Broadly viewed, the holonomies can be seen as a way of providing test functions against which to smear the fields of a theory. Because loops are one dimensional objects, the resulting smearings tend to be distributional in nature. This creates regularization problems when operators act on the holonomies. In the context of gravity for instance, where one wishes to have quantum states that are invariant under diffeomorphisms, this has led to considering functions of thickened or framed loops [2]. In the Yang–Mills context, the definition of the inner product in the loop representation would require summations over families of loops that are not well defined. To deal with these issues the concept of extended loops and the ensuing extended holonomies was introduced [3, 4]. The idea is to construct smearing functions that share some of the properties of the smearings provided by loops, but that are more general and have three-dimensional support. Expansions formally similar to the ones used to define the non-Abelian holonomy can be constructed in terms of those smeared functions to construct an extended holonomy.

It was shown [5], however, that in spite of the formal analogy with the case of loops, some convergence issues appeared in the expansion which rendered the extended holonomy to be non-gauge covariant. A potential solution was suggested [6] to this problem by considering certain subsets of extended holonomies. However, the solution was not entirely satisfactory since the proposed subsets were ad-hoc in nature. In spite of these difficulties the techniques attracted some attention in the mathematical and particle physics literature [7]. Here we would like to overcome those limitations by providing a generic definition of the families of extended loops that yield properly covariant holonomies.

The structure of this paper is as follows. In the next section we will review the concept of multitanents in ordinary holonomies. In section 3 we discuss extended holonomies. In section 4 we will propose the construction of extended loops that yield covariant extended holonomies. Section 5 proposes an explicit construction of extended loops leading to covari-

ant holonomies. We end with a discussion.

## II. ORDINARY HOLOMOMIES AND MULTITANGENTS

The holonomy (whose trace is the *Wilson loop*) of a connection  $A_a$  is given by its path ordered exponential along a loop<sup>1</sup>. It can be rewritten as,

$$U_A(\gamma) = 1 + \sum_{n=1}^{\infty} (-i)^n \int dx_1^3 \cdots dx_n^3 A_{a_1}(x_1) \cdots A_{a_n}(x_n) T^{a_1 \cdots a_n}(x_1, \dots, x_n, \gamma) \quad (2.1)$$

where  $\gamma$  is a loop with a base point  $o$  which we take as its origin and the loop dependent multitangents  $T$  are given by,

$$\begin{aligned} T^{a_1 \cdots a_n}(x_1, \dots, x_n, \gamma) &= \int_{\gamma} dy_n^{a_n} \int_0^{y_n} dy_{n-1}^{a_{n-1}} \cdots \int_0^{y_2} dy_1^{a_1} \delta(x_n - y_n) \cdots \delta(x_1 - y_1) \\ &= \oint_{\gamma} dy_n^{a_n} \cdots \oint_{\gamma} dy_1^{a_1} \delta(x_n - y_n) \cdots \delta(x_1 - y_1) \Theta_{\gamma}(0, y_1, \dots, y_n) \end{aligned} \quad (2.2)$$

with the multi Heaviside function  $\Theta_{\gamma}(0, y_1, \dots, y_n)$  ordering the points along the loop starting at the origin and the Dirac deltas are three dimensional ones. These relations define the multitangents of “rank”  $n$ . It will be convenient to introduce the notation

$$T^{\mu_1 \cdots \mu_n}(\gamma) = T^{a_1 x_1 \cdots a_n x_n}(\gamma) = T^{a_1 \cdots a_n}(x_1, \dots, x_n, \gamma), \quad (2.3)$$

with  $\mu_i \equiv (a_i x_i)$ , which suggests better the role played by the  $x$  variables under diffeomorphisms [3, 4].

The multitangents satisfy a set of *algebraic identities*, which follow directly from properties of the Heaviside function,

$$T^{\underline{\mu_1 \cdots \mu_k} \mu_{k+1} \cdots \mu_n} \equiv \sum_{P_k} T^{P_k(\mu_1 \cdots \mu_n)} = T^{\mu_1 \cdots \mu_k} T^{\mu_{k+1} \cdots \mu_n}, \quad (2.4)$$

with the summation over all the permutations  $P_k$  of the first  $k$  of the  $\mu$  variables which preserve the ordering of the  $\mu_1, \dots, \mu_k$  and the  $\mu_{k+1}, \dots, \mu_n$  among themselves.

They also satisfy a *differential constraint*,

$$\frac{\partial}{\partial x_i^{a_i}} T^{a_1 x_1 \cdots a_i x_i \cdots a_n x_n} = (\delta(x_i - x_{i-1}) - \delta(x_i - x_{i+1})) T^{a_1 x_1 \cdots a_{i-1} x_{i-1} \ a_{i+1} x_{i+1} \cdots a_n x_n}, \quad (2.5)$$

with  $x_0$  and  $x_{n+1}$  equal to the origin of the loop  $o$ .

The holonomy is gauge covariant, that is, if we consider a gauge transformation  $U$ ,

$$A_a \rightarrow A'_a = U A_a U^\dagger - (\partial_a U) U^\dagger, \quad (2.6)$$

---

<sup>1</sup> It should be noted that in this context a “loop” is an equivalence class of curves that differ by retracings called “trees”. All curves in the class yield the same holonomy and have the same multitangents.

we have that,

$$U_{A'}(\gamma) = U_o U_A(\gamma) U_o^\dagger \quad (2.7)$$

Notice that the gauge transformation is a function of point and  $U_o$  is the gauge transformation evaluated at the origin of the loop. The covariance of the holonomy follows from the path ordered nature along the loop of the holonomy. It is customary to take the loop origin at infinity and the small gauge transformations (the ones connected to the identity) as the identity at infinity. However, we will not make assumptions about this in this paper.

A gauge field can be viewed as stemming from a representation of the group of loops in a Lie group  $G$ . Every representation defines a connection up to gauge transformations that leads to expansions for the holonomy that are convergent (for a detailed discussion see [8]).

The multitangents transform as multivector densities under the subgroup of coordinate transformations that leaves the base point  $o$  fixed. That is, if one has a transformation,

$$x^a \longrightarrow x'^a = D^a(x) \quad (2.8)$$

then

$$T^{a_1 x'_1 \dots a_n x'_n}(DL) = \frac{\partial x'^{a_1}}{\partial x_1^{b_1}} \dots \frac{\partial x'^{a_n}}{\partial x_n^{b_n}} \frac{1}{J(x_1)} \dots \frac{1}{J(x_n)} T^{b_1 x_1 \dots b_n x_n}(L). \quad (2.9)$$

where  $J$  is the Jacobian of the transformation. We will call objects whose components transform in this way *multitensors*.

### III. EXTENDED LOOPS AND HOLONOMIES

Given a multitensor  $E = (E^0, E^{\mu_1}, \dots, E^{\mu_1 \dots \mu_n}, \dots)$ , where  $E^0$  is a real constant that in what follows we take equal to one, we can define an extended holonomy,

$$U_A(E) = 1 + \sum_{n=1}^{\infty} \int (-i)^n dx_1 \dots dx_n A_{\mu_1} \dots A_{\mu_n} E^{\mu_1 \dots \mu_n}. \quad (3.1)$$

with  $E$  satisfying the differential and algebraic constraints. The multitensors  $E$  are a generalization of the multitangents. They have a product,

$$(E_1 \times E_2)^{\mu_1 \dots \mu_n} = \sum_{i=0}^n E_1^{\mu_1 \dots \mu_i} E_2^{\mu_{i+1} \dots \mu_n}, \quad (3.2)$$

that is related to the product of loops, which form a group [1],

$$T(\gamma_1 \circ \gamma_2) = T(\gamma_1) \times T(\gamma_2). \quad (3.3)$$

The product is associative and satisfies the differential constraint.

It is convenient to rewrite the generalized holonomy as,

$$U_A(E) = \sum_{n=0}^{\infty} A_{\mu_1} \dots A_{\mu_n} E^{\mu_1 \dots \mu_n}, \quad (3.4)$$

where from now on  $A_a(x) = -iA_a(x)$  and implicit in the sum are the integrals along space, with the term with  $n = 0$  ( $E$  with zero components, which we will call  $E^0$ ) equal to one. It is also useful to define the term containing  $n$  powers of the connection as,

$$U_A^{(n)}(E) = A_{\mu_1} \dots A_{\mu_n} E^{\mu_1 \dots \mu_n}. \quad (3.5)$$

Let us address the issue of gauge invariance. Consider an infinitesimal gauge transformation,

$$A^\Lambda = A + d\Lambda + [A, \Lambda] \quad (3.6)$$

the terms in the sum transform as,

$$U_{A^\Lambda}^{(n)} = U_A^{(n)} + \left[ U_A^{(n-1)}(E), \Lambda_o \right] + f^{(n)}_{(A,\Lambda)}(E) - f^{(n-1)}_{(A,\Lambda)}(E), \quad (3.7)$$

with  $\Lambda_o$  the gauge transformation parameter evaluated at the origin of the loop and,

$$f_{(A,\Lambda)}^{(n)}(E) = \sum_k X^{\mu_1 \dots \mu_n} A_{\mu_1} \dots A_{\mu_{k-1}} [A, \Lambda]_{\mu_k} A_{\mu_{k+1}} \dots A_{\mu_n}, \quad (3.8)$$

and therefore,

$$\sum_{n=0}^N U_{A^\Lambda}^{(n)}(E) = \sum_{n=0}^N U_A^{(n)}(E) + \left[ \sum_{n=0}^N U_A^{(n)}(E), \Lambda_o \right] - \left[ U_A^{(N)}(E), \Lambda_o \right] + f_{(A,\Lambda)}^{(N)}(E). \quad (3.9)$$

If  $U_A(E)$  converges, the next to last terms vanishes and the extended holonomy is invariant if and only if  $f_{(A,\Lambda)}^{(N)}(E) = 0$  for  $N \rightarrow \infty$ . It is not obvious that this holds for all  $E$  and  $A$ . In a nutshell, this was one of the points of the criticism in [5]. The possibility of having extended loops for which  $f(A, \Lambda)$  is non-vanishing allows in principle [5] to find counterexamples of extended loops that do not lead to gauge covariant holonomies. In what follows we will show how to construct explicitly extended loops that lead to gauge covariant holonomies and we will establish sufficient conditions that ensure the covariance.

#### IV. CONSTRUCTING EXTENDED INVARIANTS

Let us see that one can define extended loops that lead to gauge covariant extended holonomies. The starting point of the construction is the invariance for the case of loops, equation (2.7), and the observation made in [6]: one needs to restrict the type of extended loops considered, as we shall see in detail. We will confine the discussion to  $SU(N)$  but it can be extended to other Lie groups. Let us define, as we did with the extended loop a multitangent  $T(\gamma)$  with  $T^0(\gamma) = 1$  and introduce its product with a ‘‘multiconnection’’ (a product of connections),

$$T(\gamma) \cdot A \equiv U_A(\gamma) = \sum_{n=0}^{\infty} T^{\mu_1 \dots \mu_n} A_{\mu_1} \dots A_{\mu_n}, \quad (4.1)$$

and

$$T(\gamma) \cdot A_g = U_{A_g}(\gamma) = U_g(T(\gamma) \cdot A) U_g^\dagger, \quad (4.2)$$

where  $A_g$  is the gauge transformed  $A$  connection. Let us define

$$T(a, \gamma) \equiv (1 - a)\mathcal{I} + aT(\gamma), \quad (4.3)$$

with  $\mathcal{I}$  a multitensor that has as only non-vanishing component  $\mathcal{I}^0 = 1$ , that is,  $\mathcal{I}^{\mu_1 \dots \mu_n} = 0$  for  $n \neq 0$ . To put it another way,  $T^0(a, \gamma) = 1$  and  $T^{\mu_1 \dots \mu_n}(a, \gamma) = aT^{\mu_1 \dots \mu_n}(\gamma)$ . We therefore have that,

$$T(a, \gamma) \cdot A = (1 - a)I + aU_A(\gamma) \equiv U_A(a, \gamma), \quad (4.4)$$

with  $I$  the identity in the group, and for  $a = 1$  we have that  $T(1, \gamma) \cdot A = U_A(1, \gamma) = U_A(\gamma)$ . Notice however, that  $U(a, \gamma)$  is not an element of the group if  $a \neq 1$ .

In order to define the family of extended loops that will lead to a covariant holonomy, let us consider the expansion of the logarithm of  $T$ ,

$$F(a, \gamma) \equiv \ln(T(a, \gamma)) = \sum_{i=1}^{\infty} \frac{(-1)^i}{i} (\mathcal{I} - T(a, \gamma))^i, \quad (4.5)$$

(the  $i$ -th power is computed with the multitensor product introduced before) and therefore,

$$F(a, \gamma) \cdot A = \sum_{i=1}^{\infty} \frac{(-1)^i}{i} (I - U_A(a, \gamma))^i, \quad (4.6)$$

and we shall see that the series converges for  $a$  sufficiently small. Let  $\|M\|$  be the Frobenius norm of the matrix  $M$  defined as,

$$\|M\| \equiv \sqrt{\text{Tr}(M^\dagger M)} = \sqrt{\sum_i |\lambda_i(M)|^2} \quad (4.7)$$

with  $\lambda_i$  the eigenvalues, for diagonalizable matrices. An important property of the Frobenius norm is that it is submultiplicative  $\|AB\| \leq \|A\| \|B\|$ .

Let us consider the norm of  $I - U_A(a, \gamma)$ . If we have that

$$\|I - U_A(a, \gamma)\| < 1, \quad (4.8)$$

given the submultiplicative property of the norm, this immediately implies the expansion of the logarithm converges and therefore  $F(a, \gamma)$  is well defined. Evaluating,

$$\begin{aligned} \|I - U_A(a, \gamma)\| &= \|aI - aU_A(\gamma)\| = \sqrt{\text{Tr} \left( \left( aI - aU_A^\dagger(\gamma) \right) \left( aI - aU_A(\gamma) \right) \right)} \\ &= \sqrt{\text{Tr} \left( 2a^2 I - a^2 \left( U_A^\dagger(\gamma) + U_A(\gamma) \right) \right)} \leq 2a\sqrt{N}, \end{aligned} \quad (4.9)$$

which can be seen given the generic form of a unitary transformation in  $SU(N)$ . Therefore  $F(a, \gamma) \cdot A$  is well defined if  $a < 1/(2\sqrt{N})$ .

Let us proceed to verify that the extended holonomy transforms appropriately. Given that  $\ln U_A(a, \gamma) = F(a, \gamma) \cdot A$  and taking into account that,

$$U_{A_g}(a, \gamma) = U_o(1 - a)U_o^\dagger + aU_o U_A(\gamma)U_o^\dagger = U_o U_A(a, \gamma)U_o^\dagger, \quad (4.10)$$

where  $U_o$  is the gauge transformation at the loop origin, and that,

$$U_A(a, \gamma) = \exp(F(a, \gamma) \cdot A), \quad (4.11)$$

it follows from (4.6) that

$$F(a, \gamma) \cdot A_g = U_o F(a, \gamma) \cdot A U_o^\dagger. \quad (4.12)$$

Equality (4.11) holds for  $a < 1/(2\sqrt{N})$  and can be extended analytically to the value  $a = 1$ . To see this recall that the expansion of the exponential of the logarithm of  $x$  is

well defined  $\forall x$  even though the expansion of the logarithm is valid only for  $|x| < 1$ . The resulting series converges to  $\ln(|x|) + i\text{Arg}(x)$  with  $\text{Arg}(x)$  is the phase of  $x$  in the interval  $[-\pi, \pi]$ .

Evaluating at  $a = 1$  we have,

$$U_A(1, \gamma) = U_A(\gamma), \quad (4.13)$$

and

$$T(\gamma) = T(1, \gamma) = \exp(F(a, \gamma))|_{a=1}. \quad (4.14)$$

Note that  $F(a, \gamma)|_{a=1} \cdot A$  belongs in the algebra (see appendix), its exponential is unitary and coincides with the principal logarithm of  $U(a, \gamma)|_{a=1}$ .

The analytic extension allows various generalizations of the concept of loop that constitute extended loops for which the holonomy transforms covariantly.

For instance, starting from (4.11) and using (4.12) we can define a gauge covariant real power of a holonomy,

$$U_{A_g}(a, \gamma)^\lambda = \exp(\lambda F(a, \gamma) \cdot A_g) = \exp(U_o \lambda F(a, \gamma) \cdot AU_o^\dagger) = U_o U_A^\lambda(a, \gamma) U_o^\dagger, \quad (4.15)$$

and  $U_A(a = 1, \gamma)^\lambda$  is the principal branch of the real  $\lambda$ -th power of a holonomy associated with,

$$T^\lambda(\gamma) = \exp(\lambda F(a, \gamma))|_{a=1}, \quad (4.16)$$

which is an example of an extended loop that leads to a covariant holonomy. There are many examples of covariant extensions of loops. The technique presented obviously includes ordinary loops. Real powers of loops are clearly invertible and form a group, the associated holonomies are unitary and gauge covariant.

The covariant extensions stem from observing that the analytic extension,

$$F_A(\gamma) \equiv F(a, \gamma) \cdot A|_{a=1}, \quad (4.17)$$

belongs to the  $SU(N)$  algebra (see appendix) and is gauge covariant for all loops  $\gamma$ . That implies that,

$$F_A(\gamma_1) + F_A(\gamma_2), \lambda F_A(\gamma), [F_A(\gamma_1), F_A(\gamma_2)], \quad (4.18)$$

lead, through exponentiation, to elements of the group that are gauge invariant and define an extended loop algebra with their corresponding extended holonomies.

The idea that this algebra allows to define smoothed loops can be confirmed considering a bi-parametric family  $\gamma(\alpha, \beta)$  of loops. The quantity,

$$\exp\left(\int d\alpha d\beta \lambda(\alpha, \beta) [F(a, \gamma(\alpha, \beta)) \cdot A]\right)\Big|_{a=1} = U_A(\gamma(\alpha, \beta), \lambda(\alpha, \beta)) \quad (4.19)$$

where  $\lambda(\alpha, \beta)$  a suitable functional coefficient for each member of the family is an example of smoothed loop. Summarizing, starting from ordinary loops one can construct a large family of extended loops that lead to covariant holonomies. The key observation is that although  $F(a, \gamma)|_{a=1}$ , given by (4.5) in general is not well defined, since the convergence radius of the expansion of the exponential is infinite,  $[\exp(F(a, \gamma))]|_{a=1}$  is.

Summarizing, given  $T(\gamma) = \exp(F(a, \gamma))|_{a=1}$ , we can define a family of extended loops  $E(\lambda F_\gamma)$ ,

$$T^\lambda(\gamma) = \exp(\lambda F(a, \gamma))|_{a=1} \equiv E(\lambda F_\gamma), \quad (4.20)$$

and construct,

$$E(\lambda F_{\gamma_1} + \mu F_{\gamma_2}) = \exp(\lambda F(a, \gamma_1) + \mu F(a, \gamma_2)) \Big|_{a=1}, \quad (4.21)$$

or  $E(\int d\alpha d\beta \lambda(\alpha, \beta) F_{\gamma(\alpha, \beta)})$ , or  $E([F_{\gamma_1}, F_{\gamma_2}])$  and  $E$  of multiple commutators defined analogously. All of them take the form,

$$E = (E^0 = 1, E^{\mu_1}, E^{\mu_1, \mu_2}, \dots, E^{\mu_1 \dots \mu_n}), \quad (4.22)$$

but the  $E^{\mu_1 \dots \mu_n}$  satisfy additional conditions to the differential and algebraic constraints: they are exponentials of  $F$ 's, as constructed above. The components of  $E$  lead to series that converge to unitary and gauge covariant transformations and are examples of extensions that satisfy equations (3.3) and (3.4) of [5].

Notice that the  $F$ 's satisfy the differential constraint and a simpler version of the algebraic constraint given by  $F(a, \gamma)^{\underline{\mu_1 \dots \mu_k \mu_{k+1} \dots \mu_n}} \Big|_{a=1} = 0$ . This constraint is a key ingredient in the construction of elements of the Lie group algebra [4].

To prove that the constraint is satisfied, let us consider the continuous binomial expansion

$$T^\lambda(a, \alpha) = (\mathcal{I}(1-a) + aT(\alpha))^\lambda = \sum_{m=0}^{\infty} \binom{\lambda}{m} (1-a)^m a^{\lambda-m} T^{\lambda-m}(\alpha), \quad (4.23)$$

with  $\alpha$  a loop. Given that the multitangents satisfy the algebraic constraint

$$T^{\lambda-m}(\alpha)^{\underline{\mu_1 \dots \mu_k \mu_{k+1} \dots \mu_n}} = T^{\lambda-m}(\alpha)^{\mu_1 \dots \mu_k} T^{\lambda-m}(\alpha)^{\mu_{k+1} \dots \mu_n}, \quad (4.24)$$

differentiating the product in  $T^\lambda(a, \alpha)^{\underline{\mu_1 \dots \mu_k \mu_{k+1} \dots \mu_n}}$ ,

$$\begin{aligned} \frac{d}{d\lambda} T^\lambda(a, \alpha)^{\underline{\mu_1 \dots \mu_k \mu_{k+1} \dots \mu_n}} &= \sum_{m=0}^{\infty} (1-a)^m \frac{d}{d\lambda} \left( \binom{\lambda}{m} a^{\lambda-m} \right) T^{\lambda-m}(\alpha)^{\mu_1 \dots \mu_k} T^{\lambda-m}(\alpha)^{\mu_{k+1} \dots \mu_n} \\ &+ \sum_{m=0}^{\infty} \binom{\lambda}{m} (1-a)^m a^{\lambda-m} \frac{d}{d\lambda} T^{\lambda-m}(\alpha)^{\mu_1 \dots \mu_k} T^{\lambda-m}(\alpha)^{\mu_{k+1} \dots \mu_n} \\ &+ \sum_{m=0}^{\infty} \binom{\lambda}{m} (1-a)^m a^{\lambda-m} T^{\lambda-m}(\alpha)^{\mu_1 \dots \mu_k} \frac{d}{d\lambda} T^{\lambda-m}(\alpha)^{\mu_{k+1} \dots \mu_n}, \end{aligned} \quad (4.25)$$

where  $T^{-m}(\alpha) \equiv T^m(\bar{\alpha})$  [4]. Evaluating the derivative at  $\lambda = 0$ , we obtain the algebraic constraint of the generators of  $T^\lambda(a, \alpha)$ ,

$$\begin{aligned} F(a, \alpha)^{\underline{\mu_1 \dots \mu_k \mu_{k+1} \dots \mu_n}} &= \frac{d}{d\lambda} T^\lambda(a, \alpha)^{\underline{\mu_1 \dots \mu_k \mu_{k+1} \dots \mu_n}} \Big|_{\lambda=0} \\ &= \sum_{m=1}^{\infty} \left( \frac{1}{a} - 1 \right)^m \frac{(-1)^{m-1}}{m} (1 + \gamma(m-1)) T^m(\bar{\alpha})^{\mu_1 \dots \mu_k} T^m(\bar{\alpha})^{\mu_{k+1} \dots \mu_n} \\ &+ F(\alpha)^{\mu_1 \dots \mu_k} \mathcal{I}^{\mu_{k+1} \dots \mu_n} + \mathcal{I}^{\mu_1 \dots \mu_k} F(\alpha)^{\mu_{k+1} \dots \mu_n}, \end{aligned} \quad (4.26)$$

where  $\gamma$  is the Euler–Mascheroni constant. Finally, for  $1 < k < n$  and  $a \rightarrow 1$ ,  $F(a, \gamma)^{\underline{\mu_1 \dots \mu_k \mu_{k+1} \dots \mu_n}} \Big|_{a=1} = 0$  since  $\mathcal{I}^{\mu_1 \dots \mu_n} = 0$  for  $n \geq 1$ . This ensures that the product  $F \cdot A$  is in the algebra (see appendix). One can also demonstrate that the exponential of any quantity satisfying the homogeneous algebraic constraint produces an object that satisfies the algebraic constraint.



## V. AN EXPLICIT CHARACTERIZATION OF EXTENDED LOOPS LEADING TO COVARIANT HOLONOMIES

Up to now we have followed a constructive process to identify extended loops, either considering real powers of a loop or more general constructions, always starting from ordinary loops. However, it is convenient to have a notion of extended loops that lead to covariant holonomies, independent of their construction procedure .

The non covariance of the extended holonomy discussed in [5] was based on a two fold argument. One of them can be solved by regularization: gauge transformations of extended holonomies based on real powers of loop holonomies may appear to be non invariant due to the appearance of different branches of the exponentiation of a function by a real parameter. The complex power function is a multi-valued function The principal branch of the function is obtained by replacing  $\ln(z)$  with the principal branch of the logarithm. If one adds to this observation that ordinary loops lead to gauge invariant holonomies then this leads to extensions that also yield gauge invariant holonomies as we showed in previous sections.

The second problem regards the difficulty to prove that the gauge transformation of the local connection transforms correctly in the limit where the holonomy includes an infinite number of nodes (gauge field insertions) in the extended loop. That is, the last term in (4.6) vanishes. It is known that holonomies constructed in terms of loops transform correctly and one can prove it as in [9], by partitioning the loop  $\gamma$  into a collection of infinitesimal straight segments  $S_{z_i}^{z_{i+1}}$  that form a polygonal that, as you increase the number of segments approaches the curve . For a large number  $N + 1$  of segments

$$U_A(\gamma) = \mathcal{P} \exp \left\{ \int_{\gamma} A_a(x) dx^a \right\} \simeq \left( 1 + (x_{N+1}^{a_N} - x_N^{a_N}) A_{a_N}(x_N) + O((x_N - x_{N-1})^2) \right) \dots \\ \times \left( 1 + (x_1^{a_0} - x_0^{a_0}) A_{a_0}(x_0) + O((x_1 - x_0)^2) \right), \quad (5.1)$$

and we note that in the limit  $N \rightarrow \infty$  the right hand side reproduces the left hand side. We can identify the loop holonomy as an ordered product of infinitesimal open path parallel transports

$$U_A(S_{x_i}^{x_{i+1}}) = 1 + (x_{i+1}^{a_i} - x_i^{a_i}) A_{a_i}(x_i) + O((x_{i+1} - x_i)^2) \quad (5.2)$$

where it has been assumed that the distance between one point and the next is infinitesimal. It can easily be proven that arbitrary gauge transformations act as

$$U_{A^g}[S_{x_i}^{x_{i+1}}] = U_g(x_i) U_A[S_{x_i}^{x_{i+1}}] U_g^\dagger(x_{i+1}) + O((x_{i+1} - x_i)^2),$$

with  $g$  the element of the group associated with the gauge transformation, and we immediately get in the limit where the infinitesimal intervals go to zero

$$U_{A^g}[\gamma] = U_g(x_0) U_A[\gamma] U_g^\dagger(x_0),$$

where we have used  $x_{N+1} = x_0$  and  $U_g(x_0) = U_o$ .

To understand how these results extend to the case of extended loops, let us rewrite the above expression in terms of multitangents. To this aim, it is convenient to partition space into cubes and consider the set of cubes that are intersected by the path. We consider a cubic lattice characterized by a lattice size  $l$ , and substitute the curve inside each cell by a straight line, entering through  $y_i$  and exiting through  $y_{i+1}$ , and composing the straight lines

into an  $N + 1$  sided polygonal. Substituting the tangent vector  $dy^a \rightarrow y_{i+1}^a - y_i^a = \epsilon_i \hat{u}_i^a$ , with  $\epsilon_i$  of order  $l$  and  $\hat{u}_i^a$  unit vectors, we get,

$$\begin{aligned} T^{a_1 \dots a_n}(x_1, \dots, x_n, \gamma) &= \int_{\gamma} dy_n^{a_n} \int_0^{y_n} dy_{n-1}^{a_{n-1}} \dots \int_0^{y_2} dy_1^{a_1} \delta(x_n - y_n) \dots \delta(x_1 - y_1) \\ \rightarrow T_N^{a_1 x_1 \dots a_n x_n}(\gamma) &= \sum_{\substack{i_1 < i_2 \dots < i_n \\ 0 \leq i_j \leq N}} \epsilon_{i_1} \dots \epsilon_{i_n} \hat{u}_{i_1}^{a_1} \dots \hat{u}_{i_n}^{a_n} \delta(y_{i_1} - x_1) \dots \delta(y_{i_n} - x_n), \end{aligned} \quad (5.3)$$

where  $y_{i_j}^a = \sum_{k=0}^{i_j-1} \epsilon_k \hat{u}_k^a + y_0^a$ . Notice that this expression is valid for  $n \leq N + 1$  since the inequalities in the sum cannot be satisfied if  $n > N + 1$ , in that case  $T_N^{a_1 x_1 \dots a_n x_n} = 0$ .

Taking into account (5.1) we have that,

$$U_A(\gamma) = \lim_{N \rightarrow \infty} \left( 1 + T_N^{\mu_1} A_{\mu_1} + \dots + T_N^{\mu_1 \dots \mu_p} A_{\mu_1} \dots A_{\mu_p} + \dots + T_N^{\mu_1 \dots \mu_{N+1}} A_{\mu_1} \dots A_{\mu_{N+1}} \right) \quad (5.4)$$

and the term,

$$T_N^{\mu_1 \dots \mu_N} A_{\mu_1} \dots A_{\mu_N} = O\left(\left(\frac{1}{N}\right)^N\right), \quad (5.5)$$

and this is true for any finite connection  $A$ . In particular for both  $A$  and its gauge transformed  $A_g$ . Let us note that the polygonal multitangents  $T_N$  satisfy the algebraic constraint and the differential constraint up to higher order terms,

$$\begin{aligned} \partial_{x_i^a} T_N^{a_1 x_1, \dots, a_i x_i, \dots, a_n x_n} &= \Theta(N - n) [\delta(x_i - x_{i+1}) - \delta(x_i - x_{i-1})] T_N^{a_1 x_1, \dots, a_{i-1} x_{i-1}, a_{i+1} x_{i+1}, \dots, a_n x_n} \\ &+ O\left((x_{i+1} - x_{i-1})^2\right). \end{aligned} \quad (5.6)$$

Since expression (5.4) is an explicit form of (5.1) which is manifestly gauge invariant in the limit  $N \rightarrow \infty$ , (5.4) is too. This can be verified directly observing that the remainder term of  $f_{(A,\Lambda)}^{(N)}(E)$  of (3.9) vanishes due to (5.5) in what concerns the algebraic constraint

$$T^{\mu_1 \dots \mu_k \mu_{k+1} \dots \mu_n} = T^{\mu_1 \dots \mu_k} T^{\mu_{k+1} \dots \mu_n}, \quad (5.7)$$

Notice, however, that the polygonal multitangents do not form a group because  $T_N \times T_M = T_{N+M+1}$ . However the inverse of  $T_N$  is a  $T_N$  and the polygonal loops for arbitrary  $N$  do form a group.

So the multitangent,

$$T = \lim_{N \rightarrow \infty} T_N, \quad (5.8)$$

where  $T$  are the vectors associated with the multitangent in the notation of (4.22) and,

$$\lim_{N \rightarrow \infty} T_N^{\mu_1 \dots \mu_N} = 0. \quad (5.9)$$

These properties of the multitangents can be directly generalized to extended loops giving a sufficient criterion for the latter to lead to a covariant holonomy: the extended loop must be given by a limit  $E = \lim_{N \rightarrow \infty} E_N$  with the  $E_N$ 's satisfying a differential constraint (5.6) and a condition like (5.5) that implies limit like (5.9).

It is immediate to show that the explicit constructions presented in previous sections satisfy these conditions for a suitable definition of the limits involved. Let us see this

explicitly for the case of a real power  $\lambda$  of a loop. Let us define a polynomial approximation to the algebra element  $F(a, \gamma)$ ,

$$F(m, a, \gamma) = \sum_{i=1}^m \frac{-1}{i} (\mathcal{I} - T_{m^4}(a, \gamma))^i, \quad (5.10)$$

with the extended loop,

$$E_{m^4}(a, \gamma^\lambda) = \sum_{j=0}^m \frac{(\lambda F(m, a, \gamma))^j}{j!}. \quad (5.11)$$

It is clear that this expression satisfies the previous conditions. Noticing that the highest order term comes from  $F(m, a, \gamma)^m$  and it takes the form,

$$T_m(a, \gamma)^{\mu_1 \dots \mu_m} \times T_m(a, \gamma)^{\mu_{m+1} \dots \mu_{2m}} \times T_m(a, \gamma)^{\mu_{2m+1} \dots \mu_{3m}} \times T_m(a, \gamma)^{\mu_{3m+1} \dots \mu_{4m}} \quad (5.12)$$

which obviously goes to zero as  $T_m^{\mu_1 \dots \mu_m}$  goes to zero.

## VI. CONCLUSIONS

We have shown how to generate large families of extended loops that yield covariant holonomies. They include ordinary loops and real powers of them, among others. The real powers of loops constitute a group with associated holonomies that are unitary and gauge invariant. The center of the idea is to construct extended loops using the expansion of the logarithm of multitensors. Through an analytic extension they can be shown to yield covariant holonomies. We have also given sufficient conditions for extended loops that lead to covariant holonomies and showed that the extended loops obtained from the previous construction satisfy them.

This opens the possibility of using the ensuing extended loops to create extended loop representations of interest for the non-perturbative quantization of Yang–Mills theories and potentially gravity. In the case of Yang–Mills theories the use of extended loops could have advantages over the use of ordinary loops when one wishes to define the inner product and the closure relations, and for the renormalization of non-perturbative Schrödinger-like equations.

## VII. ACKNOWLEDGMENTS

This work was supported in part by Grants NSF-PHY-1603630, NSF-PHY-1903799, funds of the Hearne Institute for Theoretical Physics, CCT-LSU, Pedeciba and Fondo Clemente Estable FCE\_1\_2014\_1\_103803.

## Appendix

Let us show explicitly that  $F(a, \gamma) \cdot A$  with  $F$  satisfying the homogeneous algebraic identity  $F^{\mu_1 \dots \mu_k \mu_{k+1} \dots \mu_n} = 0$  is in the algebra. To do that we use a technique developed in [3, 4]. We define the matrix of delta functions,

$$\delta^{\mu_1 \dots \mu_n}_{\nu_1 \dots \nu_m} = \delta_{n,m} \delta_{\nu_1}^{\mu_1} \dots \delta_{\nu_n}^{\mu_n} \quad (7.1)$$

and the vector (using the same notation as in (4.22)),

$$\delta_{\nu_1 \dots \nu_i} = (0, \delta_{\nu_1 \dots \nu_i}^{\mu_1}, \dots, \delta_{\nu_1 \dots \nu_i}^{\mu_1 \dots \mu_n}, \dots) . \quad (7.2)$$

We define then a projector from generic multitangents to those satisfying the homogeneous algebraic constraint, given by

$$\Omega^{\mu_1 \dots \mu_n}_{\nu_1 \dots \nu_m} \equiv \frac{\delta_{n,m}}{m} [[\dots [\delta_{\nu_1}, \delta_{\nu_2}], \dots], \delta_{\nu_n}]^{\mu_1 \dots \mu_n} \theta(m-1) + \delta_{m,1} \delta_{\nu_1}^{\mu_1} \quad (7.3)$$

where

$$[\delta_{\nu_1}, \delta_{\nu_2}]^{\mu_1 \mu_2} = (\delta_{\nu_1} \delta_{\nu_2} - \delta_{\nu_2} \delta_{\nu_1})^{\mu_1 \mu_2} = \delta_{\nu_1}^{\mu_1} \delta_{\nu_2}^{\mu_2} - \delta_{\nu_2}^{\mu_1} \delta_{\nu_1}^{\mu_2}, \quad (7.4)$$

and the  $\theta$  function means that the terms is non-vanishing for  $m > 1$ . From the above definition,

$$\begin{aligned} \Omega^{\underline{\mu_1 \dots \mu_k} \mu_{k+1} \dots \mu_{n+1}}_{\nu_1 \dots \nu_{n+1}} &= \frac{n}{n+1} \left( \Omega^{\underline{\mu_1 \dots \mu_k} \mu_{k+1} \dots \mu_n}_{\nu_1 \dots \nu_n} \delta_{\nu_{n+1}}^{\mu_{n+1}} \right. \\ &+ \Omega^{\underline{\mu_1 \dots \mu_{k-1}} \mu_k \mu_{k+1} \dots \mu_{n+1}}_{\nu_1 \dots \nu_n} \delta_{\nu_{n+1}}^{\mu_k} \\ &- \delta_{\nu_{n+1}}^{\mu_1} \Omega^{\underline{\mu_2 \dots \mu_k} \mu_{k+1} \dots \mu_{n+1}}_{\nu_1 \dots \nu_n} \\ &\left. - \delta_{\nu_{n+1}}^{\mu_{k+1}} \Omega^{\underline{\mu_1 \dots \mu_k} \mu_{k+2} \dots \mu_{n+1}}_{\nu_1 \dots \nu_n} \right), \end{aligned} \quad (7.5)$$

we immediately have that if,

$$\Omega^{\underline{\mu_1 \dots \mu_l} \mu_{l+1} \dots \mu_n}_{\nu_1 \dots \nu_n} = 0, \quad \forall l/1 \leq l < n, \quad (7.6)$$

then,

$$\Omega^{\underline{\mu_1 \dots \mu_k} \mu_{k+1} \dots \mu_{n+1}}_{\nu_1 \dots \nu_{n+1}} = 0, \quad \forall k/1 \leq k < n+1. \quad (7.7)$$

Given that  $\Omega$  is a projector, one has that

$$F \cdot A = (\Omega \cdot F) \cdot A = F \cdot (\Omega \cdot A), \quad (7.8)$$

and  $\Omega \cdot A$  is in the algebra and  $F \cdot A$  is too. This can be seen in the following way,

$$(\Omega_{\nu_1 \dots \nu_n} \cdot A)_{\nu_1 \dots \nu_n} = \Omega^{\mu_1 \dots \mu_n}_{\nu_1 \dots \nu_n} A_{\mu_1} \dots A_{\mu_n} \quad (7.9)$$

$$\begin{aligned} &= \frac{1}{n} \left( \Omega^{\mu_1 \dots \mu_{n-1}}_{\nu_1 \dots \nu_{n-1}} A_{\mu_1} \dots A_{\mu_{n-1}} \dots A_{\mu_{n-1}} A_{\nu_n} \right. \\ &\quad \left. - A_{\nu_n} \Omega^{\mu_2 \dots \mu_n}_{\nu_1 \dots \nu_{n-1}} A_{\mu_2} \dots A_{\mu_n} \right) \end{aligned} \quad (7.10)$$

$$= \frac{1}{n} [\Omega_{\nu_1 \dots \nu_{n-1}} \cdot A, A_{\nu_n}] \quad (7.11)$$

$\vdots$

$$= \frac{1}{n!} [\dots [A_{\nu_1}, A_{\nu_2}], \dots, A_{\nu_n}], \quad (7.12)$$

as can be seen inductively and therefore if  $A_\nu$  is in the algebra, so is  $\Omega_{\nu_1 \dots \nu_n} \cdot A$ .

---

[1] R. Gambini, J. Pullin, "Loops, knots, gauge theories and quantum gravity", Cambridge University Press, Cambridge, UK (1996).

- [2] R. Gambini, J. Pullin, “The Gauss linking number in quantum gravity”, in “Knots and quantum gravity”, J. Baez (editor), Oxford University Press, Oxford, Uk (1994) [gr-qc/9310025].
- [3] C. Di Bartolo, R. Gambini, J. Griego and J. Pullin, Phys. Rev. Lett. **72**, 3638 (1994) doi:10.1103/PhysRevLett.72.3638 [gr-qc/9312029]; C. Di Bartolo, R. Gambini and J. Griego, Phys. Rev. D **51**, 502 (1995) doi:10.1103/PhysRevD.51.502 [gr-qc/9406039]
- [4] C. Di Bartolo, R. Gambini and J. Griego, Commun. Math. Phys. **158**, 217 (1993) doi:10.1007/BF02108073 [gr-qc/9303010];
- [5] T. A. Schilling, J. Math. Phys. **37**, 4041 (1996) doi:10.1063/1.531615 [gr-qc/9503064].
- [6] C. Di Bartolo, R. Gambini, J. Griego and J. Pullin, J. Math. Phys. **36**, 6510 (1995) doi:10.1063/1.531254 [gr-qc/9503059].
- [7] J. N. Tavares, J. Geom Phys. 26, 311 (1998); I. O. Cherednikov, T. Mertens and F. F. Van der Veken, De Gruyter Stud. Math. Phys. **24** (2014).
- [8] R. Gambini and A. Trias, R. Gambini and A. Trias, Phys. Rev. D **23**, 553 (1981).
- [9] Giles, R. (1981). *Phys. Rev.* D24, 2160.