

5-10-2021

Study of Parity Sheaves Arising from Graded Lie Algebras

Tamanna Chatterjee

Follow this and additional works at: https://repository.lsu.edu/gradschool_dissertations



Part of the [Physical Sciences and Mathematics Commons](#)

Recommended Citation

Chatterjee, Tamanna, "Study of Parity Sheaves Arising from Graded Lie Algebras" (2021). *LSU Doctoral Dissertations*. 5539.

https://repository.lsu.edu/gradschool_dissertations/5539

This Dissertation is brought to you for free and open access by the Graduate School at LSU Scholarly Repository. It has been accepted for inclusion in LSU Doctoral Dissertations by an authorized graduate school editor of LSU Scholarly Repository. For more information, please contact gradetd@lsu.edu.

STUDY OF PARITY SHEAVES ARISING FROM GRADED LIE ALGEBRAS

A Dissertation

Submitted to the Graduate Faculty of the
Louisiana State University and
Agricultural and Mechanical College
in partial fulfillment of the
requirements for the degree of
Doctor of Philosophy

in

The Department of Mathematics

by

Tamanna Chatterjee

B.S., University of Burdwan, India, 2013

M.S., RKMVERI, India, 2016

August 2021

This thesis is dedicated to my father Kishan Chatterjee.

Acknowledgments

This dissertation would not ever exist without the enormous effort and encouragement from my advisor Pramod N. Achar. No words can be sufficient for all the insightful tips, criticism, patience and unconditional support that I got from you. Thank you Pramod for being my advisor, for suggesting the problem and for always being there through all my rough times both mathematically and non-mathematically.

Thanks to the department of mathematics, all the faculties, graduate students and officials. It's been one year I have not met most of you because of the pandemic, but I will always carry the memory of cosiness that I have shared with you all. Thanks to professor Oliver Dasbach and Shea Vela-Vick for agreeing to be in my committee. Thank you professor Oxley for teaching me how to teach and always being there with so many useful tips and encouragement. Thanks Nurdin for being such a great office-mate and helping me out with all my stupid computer issues. Thank you Ryan for all those quite essential late night chit-chats that kept me alive through the confinement. Thanks to Kent, Joey, Prerna, Alisina, Shashika, Ruchira, Rima for coloring this journey with gorgeous vibes.

A big thanks to my father, Kishan, who is not in this world to hear it but without your existence in my life I can not imagine myself being this 'me'. Loosing you in the last year of my PhD and in the time of confinement due to the pandemic was the hardest battle that I fought till now. But it's you and your stories that kept me moving towards the light while being in the tunnel of deepest darkness. Thanks to my mom, Krishnapriya for all the struggles that you had to go through just to keep me strong enough throughout this journey. Nothing can describe the effort and patience that you had to make to raise me since my childhood to a strong, independent lady. After all the roughness that we went

through last year I hope this little piece of paper will bring a smile on your face, which will last long.

Thanks to all my father's friends Zinna, Nowbat, Tapas and many others all over the world, without you I would not be able collect this strength of moving forward. Thanks to Asha Murphy, my therapist, for listening to me for hours and days and giving me the space to burst into tears. Thanks to my uncle, Kshitish without whom I would never think of exploring this magnificent subject, Mathematics. Thanks for keeping faith on me in all my failures. Thank you my cousin, Pralay for all the academic tips but more than that for being my brother whom I can trust. Thank you chotomama, my maternal uncle Amalesh for being so supportive to me and my mom. Thank you Sukhamay kaku for always being my guardian angel here in Baton Rouge from the day of my arrival till today.

Lastly thanks to my significant other, Sudipta Ghosh for all the supports, jokes, foods, and understanding me in my toughest time. Without your presence this could have been an unfinished journey.

Table of Contents

Acknowledgments	iii
List of Tables	vi
Abstract	vii
Chapter 1. Introduction	1
Chapter 2. Graded Lie Algebras	4
2.1. Grading	5
2.2. Cuspidal pairs	7
2.3. Lusztig's work in characteristic 0.	10
Chapter 3. Parity Sheaves	13
3.1. Parity sheaves	14
3.2. Torsion primes and pretty good primes	15
3.3. Existence of parity for nilpotent cone	17
3.4. Existence of parity on \mathfrak{g}_n	19
Chapter 4. Lemmas On Varieties With \mathbb{C}^\times -Action	22
4.1. \mathbb{C}^\times -action on varieties	22
Chapter 5. Induction and Restriction	31
5.1. Induction and restriction	32
5.2. Lusztig's original definition	34
5.3. Cleanness for cuspidal pairs	35
Chapter 6. Induction Diagram For Cuspidal Pairs	42
6.1. Induction diagram for cuspidal pairs	42
6.2. Parity preserved for cuspidal pairs	43
Chapter 7. Existence Of Parity Sheaves	48
7.1. n -rigidity	49
7.2. Existence of parity sheaves	50
7.3. Modular reduction	54
7.4. Normal complexes	59
Chapter 8. Examples	61
8.1. \mathfrak{sp}_4-case:	61
8.2. \mathfrak{sl}_4-case:	69
8.3. Construction of \mathfrak{p} , \mathfrak{n} , and \mathfrak{l} described in ??	74
Bibliography	79

Vita 81

List of Tables

8.1. Orbits in \mathfrak{sp}_4	66
8.2. Stalks of $\text{Ind}_P^G \mathcal{IC}(\mathcal{O}_{prin}, \mathcal{L})$	69
8.3. Orbits in \mathfrak{sl}_4	72
8.4. Stalks of $\text{Ind}_P^G \mathcal{IC}(\mathcal{O}_{prin} \times \mathcal{O}_{prin}, \mathcal{L} \boxtimes \mathcal{L})$	74
8.5. Table for the Levis	78

Abstract

Let G be a complex, connected, reductive, algebraic group, and $\chi : \mathbb{C}^\times \rightarrow G$ be a fixed cocharacter that defines a grading on \mathfrak{g} , the Lie Algebra of G . Let G_0 be the centralizer of $\chi(\mathbb{C}^\times)$. In this dissertation, we study G_0 -equivariant parity sheaves on \mathfrak{g}_n , under some assumptions on the field \mathbb{k} and the group G . The assumption on G holds for GL_n and for any G , it recovers results of Lusztig[Lu] in characteristic 0. The main result is that every parity sheaf occurs as a direct summand of the parabolic induction of some cuspidal pair.

Chapter 1. Introduction

Let G be a complex, connected, reductive, algebraic group and $\chi : \mathbb{C}^\times \rightarrow G$ be a fixed cocharacter. Let G_0 be the centralizer of $\chi(\mathbb{C}^\times)$ and $\mathfrak{g}_n \subset \mathfrak{g}$ be the subspace such that $Ad(\chi(t))$ acts on it by t^n times identity. We are particularly interested in studying the derived category of G_0 -equivariant perverse sheaves on \mathfrak{g}_n , denoted by $D_{G_0}^b(\mathfrak{g}_n, \mathbb{k})$. Here \mathbb{k} is a field of positive characteristic. The simple perverse sheaves on \mathfrak{g}_n are indexed by $(\mathcal{O}, \mathcal{L})$, where \mathcal{O} is a G_0 -orbit contained in \mathfrak{g}_n and \mathcal{L} is an irreducible G_0 -equivariant \mathbb{k} -local system on \mathcal{O} . We denote this set of pairs by $\mathcal{S}(\mathfrak{g}_n, \mathbb{k})$. We define in section ??, the subset of all cuspidal pairs, $\mathcal{S}(\mathfrak{g}_n, \mathbb{k})^{\text{cusp}} \subset \mathcal{S}(\mathfrak{g}_n, \mathbb{k})$. We denote the simple perverse sheaf associated to $(\mathcal{O}, \mathcal{L})$ by $\mathcal{IC}(\mathcal{O}, \mathcal{L})$. Motivated by applications to affine Hecke Algebras, Lusztig has worked in $\mathbb{k} = \mathbb{C}$ and has proved in [Lu] that every simple perverse sheaf is a direct summand of the parabolic induction of some cuspidal pair. But in positive characteristics, this result is not true.

Following the pattern from other works in modular representation theory, often the appropriate replacement for “semisimple complex” is “parity complex”. Parity sheaves are classified as the class of constructible complexes on some stratified varieties, where the strata satisfies some cohomology vanishing properties [JMW]. We denote the parity sheaf associated to the pair $(\mathcal{O}, \mathcal{L})$ by $\mathcal{E}(\mathcal{O}, \mathcal{L})$. So the most fundamental question that arises is if they exist on \mathfrak{g}_n . Before going into that question we make some assumptions on the field coefficient. We assume that the characteristic l of \mathbb{k} is “pretty good” and the field is “big-enough” for G . Both the definitions are given in subsection 3.2. A pair $(\mathcal{O}, \mathcal{L}) \in \mathcal{S}(\mathfrak{g}_n, \mathbb{k})$ is said to be clean if $\mathcal{IC}(\mathcal{O}, \mathcal{L})$ has vanishing stalks on $\bar{\mathcal{O}} - \mathcal{O}$.

Under the above assumptions, we assume Mautner’s cleanness conjecture (conjec-

ture 2.2.7) is true, this plays an important role in the proofs of the main theorems of this paper. Mautner's conjecture already holds when the characteristic l does not divide the order of the Weyl group of the group G or if every irreducible factor of the root system of G is either of type A, B_4, C_3, D_5 or of exceptional types. We make another conjecture (conjecture 3.2.9), at the end of section ?? that on the nilpotent cone, parabolic induction preserves the parity of any cuspidal pair on a Levi subgroup. This conjecture is known to be true for GL_n and in characteristic 0 it is true for any group G . In section 8 we will show that this conjecture is also true for Sp_4 and SL_4 . The following are the main results of this paper.

1. For any cuspidal pair, $(\mathcal{O}, \mathcal{L}) \in \mathcal{S}(\mathfrak{g}_n, \mathbb{k})^{\text{cusp}}$, $\mathcal{IC}(\mathcal{O}, \mathcal{L})$ is clean and so $\mathcal{IC}(\mathcal{O}, \mathcal{L}) = \mathcal{E}(\mathcal{O}, \mathcal{L})$.
2. Parabolic induction takes parity complexes to parity complexes.
3. For any pair $(\mathcal{O}, \mathcal{L}) \in \mathcal{S}(\mathfrak{g}_n, \mathbb{k})$, $\mathcal{E}(\mathcal{O}, \mathcal{L})$ exists and is a direct summand of the parabolic induction of some cuspidal pair.

The proof of the existence of parity sheaves for the space of quiver representation of type A, D, E is given in [Ma], using some other methodology. For some of these cases the space coming from the quiver representation is the same space that we study here. So for these cases existence of parity sheaf has already been proved for \mathfrak{g}_n .

Outline

In chapter 2 we build the necessary conception about grading. In chapter 3 we talk about background, assumptions and notations needed for parity sheaves to make sense. Chapter 4 contains the lemmas on the varieties having \mathbb{C}^\times action on it. In Chapter 5, we define $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}}$ and $\text{Res}_{\mathfrak{p}}^{\mathfrak{g}}$ in the graded setting and prove the existence of parity sheaves for

cuspidal pairs. In chapter 6, we redefine both $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}}$ and Ind_P^G for cuspidal pairs and in theorem 6.2.1, we prove that the parity condition is preserved for cuspidal pairs. In chapter 7, we prove that the parity sheaf exists for a general pair $(\mathcal{O}, \mathcal{L}) \in \mathcal{S}(\mathfrak{g}_n, \mathbb{k})$ and in 7.4.1, we prove that the parabolic induction preserves parity for a general pair. In chapter 8, we compute some examples.

Chapter 2. Graded Lie Algebras

Graded setting

In section 2.1, we define the grading on the Lie Algebra \mathfrak{g} of a complex, connected, algebraic group G and also a subspace $G_0 \subset G$. G_0 acts on \mathfrak{g}_n by the adjoint action. Our aim is to study $D_{G_0}^b(\mathfrak{g}_n)$ in positive characteristic, which is the G_0 -equivariant derived category on \mathfrak{g}_n . There are finitely many G_0 -orbits in \mathfrak{g}_n , so they provides the required stratification to talk about constructible complexes on \mathfrak{g}_n . We define $\mathcal{S}(G)$ and $\mathcal{S}(\mathfrak{g}_n)$, where both are collection of pairs containing a orbit and a local system on that orbit.

Cuspidal pairs and cleanness

In section 2.2, we define cuspidal pairs on nilpotent cone, which are the simple objects in the equivariant derived category not being induced from some smaller Levi. 0-cuspidal is the one in $\mathcal{S}(G, \mathbb{k})$ which is in the image of the modular reduction map defined in 2.2.1. We define cuspidal pair in graded setting. Cleanness is a vanishing property of simple objects that we can define both for nilpotent cone and on \mathfrak{g}_n . In characteristic 0 on nilpotent cone, \mathcal{IC} 's are clean. Lusztig [Lu2] has proved, in characteristic 0 a cuspidal pair in $\mathcal{S}(\mathfrak{g}_n)$ is clean. We assume that cleanness conjecture holds in our setting.

Study of perverse sheaves on graded Lie Algebras

In this section we talk about the work already done in characteristic 0 in graded setting. The aim of this thesis is to extend the results proved by Lusztig in [Lu] and [Lu1] in positive characteristics with some more restrictions on the field coefficients (Assumption 3.2.4).

2.1. Grading

Let \mathbb{k} be a field of characteristic $l > 0$. We consider sheaves with coefficients in \mathbb{k} . The varieties we work on will be over \mathbb{C} . Let G be a connected, reductive, algebraic group over \mathbb{C} and \mathfrak{g} be the Lie Algebra of G . If H is an algebraic group acting on X , we denote by $D_H^b(X, \mathbb{k})$ or $D_H^b(X)$, the derived category of H -equivariant constructible sheaves, which is defined in [BL], and $\text{Perv}_H(X, \mathbb{k})$, its full subcategory of H -equivariant perverse \mathbb{k} -sheaves. The constant sheaf on X with value \mathbb{k} is denoted by $\underline{\mathbb{k}}_X$ or more simply $\underline{\mathbb{k}}$.

We fix a cocharacter map, $\chi : \mathbb{C}^\times \rightarrow G$ and define,

$$G_0 = \{g \in G \mid g\chi(t) = \chi(t)g, \forall t \in \mathbb{C}^\times\}.$$

For $n \in \mathbb{Z}$, define,

$$\mathfrak{g}_n = \{x \in \mathfrak{g} \mid \text{Ad}(\chi(t))x = t^n x, \forall t \in \mathbb{C}^\times\}.$$

This defines a grading on \mathfrak{g} ,

$$\mathfrak{g} = \bigoplus_{n \in \mathbb{Z}} \mathfrak{g}_n.$$

Clearly, $\mathfrak{g}_0 = \text{Lie}(G_0)$ and G_0 acts on \mathfrak{g}_n . We have the following lemma from [Lu, pp. 158],

Lemma 2.1.1. *For $n \neq 0$, G_0 acts on \mathfrak{g}_n with only finitely many orbits.*

Recall that \mathfrak{sl}_2 is the Lie Algebra of SL_2 generated by,

$$e = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, h = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, f = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

Let $J_n = \{\phi : \mathfrak{sl}_2 \rightarrow \mathfrak{g} \mid \phi(e) \in \mathfrak{g}_n, \phi(f) \in \mathfrak{g}_{-n}, \phi(h) \in \mathfrak{g}_0\}$. We have an action of G_0 on J_n by $(g, \phi) \rightarrow \text{Ad}(g) \circ \phi$. It is easy to check that this action is well-defined.

Theorem 2.1.2. *The map from the set of G_0 -orbits on J_n to the set of G_0 -orbits on \mathfrak{g}_n defined by $\phi \rightarrow \phi(e)$ is a bijection.*

Proof. The proof follows from [Lu, Prop 3.3]. □

2.1.1. The set $\mathcal{S}(G, \mathbb{k})$ and $\mathcal{S}(\mathfrak{g}_n, \mathbb{k})$

Let \mathcal{N}_G be the nilpotent cone of G . Recall that G acts on \mathcal{N}_G and has finitely many orbits.

The set $\mathcal{S}(G, \mathbb{k})$ is the set of pairs (C, \mathcal{E}) satisfying the condition that $C \subset \mathcal{N}_G$ is a nilpotent G -orbit in \mathfrak{g} and \mathcal{E} is an irreducible G -equivariant \mathbb{k} -local system on C (up to isomorphism). G -equivariant local systems on C are in one-to-one correspondence with the irreducible representations of the component group $A_G(x) := G_x/G_x^o$ on \mathbb{k} -vector spaces, where x is in C . Hence, it follows that the set $\mathcal{S}(G, \mathbb{k})$ is finite. Sometimes when there is no confusion about the field of coefficients then we will just use $\mathcal{S}(G)$.

Let $\mathcal{S}(\mathfrak{g}_n, \mathbb{k})$ or $\mathcal{S}(\mathfrak{g}_n)$ be the set of all pairs $(\mathcal{O}, \mathcal{L})$ where \mathcal{O} is a G_0 -orbit in \mathfrak{g}_n and \mathcal{L} is an irreducible, G_0 -equivariant \mathbb{k} -local system on \mathcal{O} (upto isomorphism). For fixed \mathcal{O} , G_0 -equivariant local systems on \mathcal{O} are in one to one correspondence with the irreducible representation of $A_{G_0}(x) := (G_0)_x/(G_0)_x^o$ for $x \in \mathcal{O}$. Hence by Lemma 2.1.1, $\mathcal{S}(\mathfrak{g}_n)$ is finite.

Recall G_0 acts on \mathfrak{g}_n by the adjoint action. Now we have $\mathbb{C}^\times \times G_0$ action on \mathfrak{g}_n by $(t, g) \rightarrow t^{-n} Ad(g)$.

Lemma 2.1.3. *The $\mathbb{C}^\times \times G_0$ -orbits and G_0 -orbits coincide and each G_0 -equivariant local system is also $\mathbb{C}^\times \times G_0$ -equivariant and hence \mathbb{C}^\times -equivariant.*

Proof. Since there are finitely many G_0 -orbits in \mathfrak{g}_n , we can choose a \mathbb{C}^\times -line L in \mathfrak{g}_n and

G_0 -orbit \mathcal{O} so that $\mathcal{O} \cap L$ is dense in L . Now we can choose $x \in \mathcal{O} \cap L$. Therefore $L = \mathbb{C}^\times \cdot x$. Let $y \in \mathbb{C}^\times \cdot x - \mathcal{O}$ and let \mathcal{O}' be the G_0 -orbit of y . As y is in the closure of $\mathcal{O} \cap \mathbb{C}^\times \cdot x$, which is a subset of $\bar{\mathcal{O}}$. Therefore, $\mathcal{O}' \subset \bar{\mathcal{O}}$. Hence $\dim \mathcal{O}' < \dim \mathcal{O}$. Now as \mathbb{C}^\times action commutes with G_0 action, so $G_0^x = G_0^y$. Hence $\dim \mathcal{O}' = \dim \mathcal{O}$, which is a contradiction. So we can conclude that $\mathbb{C}^\times \cdot x - \mathcal{O}$ is empty.

For the second part, it is quite easy to show that

$$(\mathbb{C}^\times \times G_0)^x / (\mathbb{C}^\times \times G_0)^{x,\circ} \cong G_0^x / (G_0^x)^\circ.$$

Hence, $\text{Loc}_{f,G_0}(\mathfrak{g}_n) \cong \text{Loc}_{f,\mathbb{C}^\times \times G_0}(\mathfrak{g}_n)$. As \mathbb{C}^\times is sitting inside $\mathbb{C}^\times \times G_0$, any $\mathbb{C}^\times \times G_0$ -equivariant sheaf is \mathbb{C}^\times -equivariant. Hence we can conclude the G_0 -equivariant local system is $\mathbb{C}^\times \times G_0$ -equivariant and hence \mathbb{C}^\times -equivariant. \square

2.2. Cuspidal pairs

The simple objects in $\text{Perv}_G(\mathcal{N}_G, \mathbb{k})$ are of the form $\mathcal{IC}(C, \mathcal{E})$, where $(C, \mathcal{E}) \in \mathcal{S}(G)$. Let P be a parabolic subgroup of G with unipotent radical U_P and let $L \subset P$ be a Levi factor of P . One can identify L with P/U_P through the natural morphism, $L \hookrightarrow P \rightarrow P/U_P$. We consider a diagram,

$$\mathcal{N}_L \xleftarrow{\pi_P} \mathcal{N}_L + \mathfrak{u}_P \xrightarrow{e_P} G \times^P (\mathcal{N}_L + \mathfrak{u}_P) \xrightarrow{\mu_P} \mathcal{N}_G$$

where $\mathfrak{u}_P = \text{Lie}(U_P)$, π_P, e_P are the obvious maps and $\mu_P(g, x) = \text{Ad}(g)x$. Let

$$i_P = \mu_P \circ e_P : \mathcal{N}_L + \mathfrak{u}_P \rightarrow \mathcal{N}_G$$

The parabolic restriction functor denoted by,

$$\text{Res}_P^G : D_G^b(\mathcal{N}_G, \mathbb{k}) \rightarrow D_L^b(\mathcal{N}_L, \mathbb{k})$$

is defined by $\text{Res}_P^G(\mathcal{F}) = \pi_{P!} e_P^* \mu_P^* \text{For}_L^G(\mathcal{F}) = \pi_{P!} i_P^* \text{For}_L^G(\mathcal{F})$. Here

$$\text{For}_L^G : D_G^b(\mathcal{N}_G, \mathbb{k}) \rightarrow D_L^b(\mathcal{N}_G, \mathbb{k})$$

is the forgetful functor. The parabolic induction comes from the same diagram above.

$$\text{Ind}_P^G : D_L^b(\mathcal{N}_L, \mathbb{k}) \rightarrow D_G^b(\mathcal{N}_G, \mathbb{k})$$

and is defined by, $\text{Ind}_P^G(\mathcal{F}) := \mu_{P!}(e_P^* \text{For}_P^G)^{-1} \pi_P^*(\mathcal{F})$. Here again For_P^G denotes the forgetful functor and, $e_P^* \text{For}_P^G : D_G^b(G \times^P (\mathcal{N}_L + \mathfrak{u}_P)) \rightarrow D_P^b(\mathcal{N}_L + \mathfrak{u}_P)$ is the induction equivalence map.

Definition 2.2.1. 1. A simple object \mathcal{F} in $\text{Perv}_G(\mathcal{N}_G, \mathbb{k})$ is called *cuspidal* if $\text{Res}_P^G(\mathcal{F}) = 0$, for any proper parabolic P and Levi factor $L \subset P$.

2. A pair $(C, \mathcal{E}) \in \mathcal{S}(G)$, is called *cuspidal* if the corresponding simple perverse sheaf $\mathcal{IC}(C, \mathcal{E})$ is cuspidal.

Remark 2.2.2. Notice that the set of cuspidal pairs depends on the characteristic l of the field of coefficients \mathbb{k} ; so we sometime call it *l-cuspidal*. We will denote the subset of cuspidal pairs in $\mathcal{S}(G)$ or $\mathcal{S}(G, \mathbb{k})$ by $\mathcal{S}(G)^{\text{cusp}}$ or $\mathcal{S}(G, \mathbb{k})^{\text{cusp}}$.

Remark 2.2.3. From [AJHR2, Remark 2.3(1)], if $\mathcal{IC}(C, \mathcal{E})$ is cuspidal then so is

$\mathbb{D}\mathcal{IC}(C, \mathcal{E}) = \mathcal{IC}(C, \mathcal{E}^\vee)$, where \mathbb{D} is the Verdier duality functor and \mathcal{E}^\vee is the dual local system of \mathcal{E} .

2.2.1. Modular reduction

Let \mathbb{K} be a finite extension of \mathbb{Q}_l with ring of integers \mathbb{O} and residue field \mathbb{k} . Then $(\mathbb{K}, \mathbb{O}, \mathbb{k})$ constitutes an l -modular system and we can talk about the modular reduction map. Let $E \in (\mathbb{K}, \mathbb{O}, \mathbb{k})$, and $K_{G_0}(\mathfrak{g}_n, E)$ be the Grothendieck group of $D_{G_0}^b(\mathfrak{g}_n, E)$. Then the modular reduction map,

$$d : K_{G_0}(\mathfrak{g}_n, \mathbb{K}) \rightarrow K_{G_0}(\mathfrak{g}_n, \mathbb{k})$$

is defined by $d[\mathcal{IC}(\mathcal{O}, \mathcal{L})] = [\mathbb{k} \otimes_{\mathcal{O}}^L \mathcal{IC}(\mathcal{O}, \mathcal{L}_{\mathcal{O}})]$, where $\mathcal{L}_{\mathcal{O}}$ is a torsion-free part \mathcal{O} -local system. In the same way we can define the modular reduction on the nilpotent cone,

$$K_G(\mathcal{N}_G, \mathbb{K}) \rightarrow K_G(\mathcal{N}_G, \mathbb{k}).$$

If the characteristic l of \mathbb{k} is rather good for G (see Definition 2.2.6), this modular reduction induces a bijection by [AJHR],

$$\mathcal{S}(G, \mathbb{K}) \xrightarrow{\cong} \mathcal{S}(G, \mathbb{k}).$$

We will discuss the modular reduction in more detail in 7.3. The pair $(C, \mathcal{E}) \in \mathcal{S}(G, \mathbb{k})$ will be called 0-cuspidal if it is in the image of some cuspidal pair under d , and we will denote the set of 0-cuspidal pairs by $\mathcal{S}(G, \mathbb{k})^{0\text{-cusp}}$ or $\mathcal{S}(G)^{0\text{-cusp}}$.

Definition 2.2.4. $(\mathcal{O}, \mathcal{L}) \in \mathcal{S}(\mathfrak{g}_n, \mathbb{k})$ will be called cuspidal if there exists a pair $(C, \mathcal{E}) \in \mathcal{S}(G)^{0\text{-cusp}}$, such that $C \cap \mathfrak{g}_n = \mathcal{O}$ and $\mathcal{L} = \mathcal{E}|_{\mathcal{O}}$. We will denote the set of all cuspidal pairs on \mathfrak{g}_n by, $\mathcal{S}(\mathfrak{g}_n)^{\text{cusp}}$.

Remark 2.2.5. Notice that the definition of cuspidal on \mathfrak{g}_n is not coming from the restriction functor as for the nilpotent cone and there is no l version in definition of cuspidal pairs on \mathfrak{g}_n .

2.2.2. Cleanness

A pair $(C, \mathcal{E}) \in \mathcal{S}(G)$ is called l -clean if the corresponding $\mathcal{IC}(C, \mathcal{E})$ has vanishing stalks on $\bar{C} - C$. Similarly, a pair $(\mathcal{O}, \mathcal{L}) \in \mathcal{S}(\mathfrak{g}_n)$ is called l -clean if the corresponding $\mathcal{IC}(\mathcal{O}, \mathcal{L})$ has vanishing stalks on $\bar{\mathcal{O}} - \mathcal{O}$.

Definition 2.2.6. A prime number l is said to be a rather good prime for a group G , if it is a good prime for G and does not divide $|Z(G)/Z(G)^\circ|$.

By [AJHR, Lemma 2.1], a prime is rather good if and only if l does not divide $|A_G(x)|$ for any $x \in \mathcal{N}_G$. The following is a part of a series of (unpublished) conjectures by C. Mautner.

Conjecture 2.2.7. *(Mautner's cleanness conjecture) If the characteristic l of \mathbb{k} is a rather good prime for G , then every 0-cuspidal pair $(C, \mathcal{E}) \in \mathcal{I}(G)$ is l -clean.*

Remark 2.2.8. *This conjecture has been already proved when the characteristic l does not divide the order of the Weyl group of G , or if every irreducible factor of the root system of G is either of type A, B_4, C_3, D_5 , or of exceptional type [AJHR].*

Remark 2.2.9. *The cleanness conjecture is not true if we replace 0-cuspidal by l -cuspidal. A counter example is when $G = GL(2)$, and $l = 2$ then the unique 2-cuspidal pair $(O_{(2)}, \underline{\mathbb{k}})$ is not 2-clean. The proof is explained in [AJHR2, Remark 2.5].*

In this paper we assume this conjecture is true.

2.3. Lusztig's work in characteristic 0.

Like induction and restriction on nilpotent cone, we can define these functors in the graded setting. For this we use the following induction diagram,

$$\mathfrak{l}_n \xleftarrow{\pi} \mathfrak{p}_n \xrightarrow{e} G_0 \times^{P_0} \mathfrak{p}_n \xrightarrow{\mu} \mathfrak{g}_n.$$

Define, $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} : D_{L_0}^b(\mathfrak{l}_n) \rightarrow D_{G_0}^b(\mathfrak{g}_n)$ and $\text{Res}_{\mathfrak{p}}^{\mathfrak{g}} : D_{G_0}^b(\mathfrak{g}_n) \rightarrow D_{L_0}^b(\mathfrak{l}_n)$. We will talk about these functors in details in section 5. In his paper [Lu], Lusztig has studied semisimple complexes on graded pieces in char 0, where the following theorems hold,

Theorem 2.3.1. *For any cuspidal pair $(\mathcal{O}, \mathcal{L}) \in \mathcal{I}(\mathfrak{g}_n)$, $\mathcal{IC}(\mathcal{O}, \mathcal{L})$ is clean.*

Theorem 2.3.2. *For $(\mathcal{O}, \mathcal{L}) \in \mathcal{I}(\mathfrak{g}_n)$, there exist a parabolic subgroup P with Levi L and $(\mathcal{O}', \mathcal{L}') \in \mathcal{I}(\mathfrak{l}_n)^{\text{cusp}}$, so that $\mathcal{IC}(\mathcal{O}, \mathcal{L})$ occurs as a direct summand of $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} \mathcal{IC}(\mathcal{O}', \mathcal{L}')$.*

Theorem 2.3.3. $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}}$ sends semisimple complexes to semisimple complexes.

In proving Theorem 2.3.1, the cleanness of \mathcal{IC} 's for cuspidal pairs on nilpotent cone and the Lemma 2 from [Lu1] plays an important role. In this thesis we prove the cleanness of cuspidal pairs on the graded pieces in positive characteristic (Theorem 5.3.1). For cleanness on nilpotent cone in positive characteristic we assume that Mautner's cleanness conjecture (Conjecture 2.2.7) holds in our context. To prove a analogous Lemma for Lemma 2, we made some effort in chapter 4. In characteristic 0, for any Levi L contained in a parabolic P and for any cuspidal pair $(C, \mathcal{F}) \in \mathcal{S}(L)$,

$$\text{Ind}_P^G \mathcal{IC}(C, \mathcal{F}) \cong \bigoplus_{j=1}^r \mathcal{F}_j[2s_j], \text{ by [Lu, 2.6(b)]}, \quad (3.1)$$

where s_j 's are integers and \mathcal{F}_j 's are G -equivariant local systems. This statement plays an important role in the proof of Theorem 2.3.2. But this statement is not true in positive characteristic. So in this thesis we replace the above statement by Conjecture 3.2.9. To prove Theorem 2.3.2, an intermediate step was to fix an element $x \in \mathcal{O}$ and construct a parabolic P with Levi L associated to x with $\mathcal{O}_L = \mathcal{O} \cap \mathfrak{l}_n$ and $\mathcal{L}' = \mathcal{L}|_{\mathcal{O}_L}$. This (L, χ) is n -rigid (for definition see 7.1). The Lemma that helps to prove Theorem 2.3.2, is the following [Lu, Lemma 6.8],

Lemma 2.3.4. *Let $(\mathcal{O}, \mathcal{L}) \in \mathcal{S}(\mathfrak{g}_n)$ and \mathfrak{l}_n and L as constructed above. Then*

- *The support of $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} \mathcal{IC}(\mathcal{O}_L, \mathcal{L})$ is \mathcal{O} ,*
- *and $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} \mathcal{IC}(\mathcal{O}_L, \mathcal{L})|_{\mathcal{O}} = \mathcal{L}[\dim \mathcal{O}_L]$.*

We prove a analogous Theorem in positive characteristic (Theorem 7.2.2) which involves parity sheaves. This Lemma together with the above equation (3.1) helps to prove [Lu, Prop. 7.3], whose direct consequence is Lemma 2.3.2. Equation (3.1) is not true in positive characteristic, simply because decomposition theorem does not hold. So to prove

a analogous result as [Lu, Prop. 7.3], we take the help of modular reduction map (section 7.3) and prove Prop. 7.3.4.

The theorems above helps to establish a connection between the affine hecke Algebra and semisimple complexes. But all of them are false in positive characteristics. So in this thesis we tried to prove some analogous result associating parity sheaves instead of semi-simple complexes.

Chapter 3. Parity Sheaves

Introduction to parity sheaves

Parity sheaves were first introduced by Juteau, Mautner and Williamson in their paper ‘Parity sheaves’ [JMW]. They are constructible complexes with some cohomology vanishing property. In characteristic 0 they coincide with intersection cohomology complexes but in positive characteristic they are new objects. As we mentioned before, Theorem 2.3.1, Theorem 2.3.2 and Theorem 2.3.3 are not true in positive characteristic. One aim will be to replace these theorems by something similar associated to parity sheaves.

In section 3.1, we define parity sheaves in the general context, where an algebraic group H acts on the algebraic variety X . To talk about parity sheaves we need some cohomology vanishing condition that is mentioned in (1.1). Finally the definition of parity comes from the Theorem 3.1.2. That cohomology vanishing condition to hold in our context we need to pose more restriction on the field \mathbb{k} that comes from section 3.2.

Restriction on the characteristic of \mathbb{k}

The condition (1.1) in our context transforms in to the vanishing of $H_{G_0}^*(\mathcal{L})$ in odd degrees. In [JMW], they have introduced the notion of ‘rather good’ prime in terms of torsion prime. Here we need the characteristic of the field coefficients to be ‘pretty good’. Also we assume that the field \mathbb{k} is big enough for G . This serves for the irreducibility after applying the modular reduction map defined in 2.2.1. Theorem 3.2.7 sets up the proper environment to talk about parity sheaves on nilpotent cone. At the end of this section we make another conjecture (Conjecture 3.2.9).

Existence on nilpotent cone

As parity sheaves do not exist automatically as intersection cohomology complexes do, it needs some work to prove the existence. In their paper [JMW], Juteau-Mautner-williamson has dealt with this question in the context of nilpotent cone. Construction of even and semismall resolution serves the purpose in several cases as by [JMW, Prop. 2.34], the pushforward of such morphism takes parity to parity.

Existence on \mathfrak{g}_n

The main goal of this thesis is to prove the existence of parity sheaves on \mathfrak{g}_n in characteristic positive. By Theorem 3.2.8, there exist atmost one parity sheaf for a pair $(\mathcal{O}, \mathcal{L}) \in \mathcal{I}(\mathfrak{g}_n)$. But still the existence is a question. In this section we have tried to explain the proof of Maksimao [Ma] that works for us in case of $G = GL_n$.

3.1. Parity sheaves

Let H be a linear algebraic group and X be a H -variety. We fix a stratification

$$X = \coprod_{\lambda \in \Lambda} X_\lambda$$

of X into smooth connected locally closed (H -stable) subsets. For each $\lambda \in \Lambda$, $i_\lambda : X_\lambda \hookrightarrow X$ denotes the inclusion map and let d_λ be the dimension of X_λ . For each $\lambda \in \Lambda$, let $\text{Loc}_{f,H}(X_\lambda, \mathbb{k})$ or $\text{Loc}_{f,H}(X_\lambda)$ denote the category of H -equivariant \mathbb{k} -local systems of finite rank on X_λ .

According to [JMW], to talk about parity sheaves on X we need the condition below,

$$H_H^*(\mathcal{L}) = 0 \text{ for odd degrees,} \tag{1.1}$$

for any local system $\mathcal{L} \in \text{Loc}_{f,H}(X_\lambda)$ and for any $\lambda \in \Lambda$.

- Definition 3.1.1.** 1. A complex $\mathcal{F} \in D_H^b(X)$ is called **-even* if for each $\lambda \in \Lambda$ and $n \in \mathbb{Z}$, $H^n(i_\lambda^* \mathcal{F})$ belongs to $\text{Loc}_{f,H}(X_\lambda)$ and vanishes for n odd. A complex $\mathcal{F} \in D_H^b(X)$ is called **-odd* if for each $\lambda \in \Lambda$ and $n \in \mathbb{Z}$, $H^n(i_\lambda^* \mathcal{F})$ belongs to $\text{Loc}_{f,H}(X_\lambda)$ and vanishes for n even. Similarly, we can define *!-even* and *!-odd* complexes.
2. A complex \mathcal{F} is called *even* if it is both **-even* and *!-even*. A complex \mathcal{F} is called *odd* if it is both **-odd* and *!-odd*.
3. A complex \mathcal{F} is called *parity* if it splits as the direct sum of an even complex and an odd complex.

Like \mathcal{IC} sheaves, the definition of parity sheaves comes from a theorem. The following theorem requires the assumption (1.1) on $\text{Loc}_{f,H}(X_\lambda)$.

Theorem 3.1.2. *Let \mathcal{F} be an indecomposable parity complex. Then*

1. *The support of \mathcal{F} is irreducible and hence of the form \bar{X}_λ , for some $\lambda \in \Lambda$.*
2. *$\mathcal{F}|_{X_\lambda}$ is isomorphic to $\mathcal{L}[m]$ for some indecomposable object \mathcal{L} in $\text{Loc}_{f,H}(X_\lambda)$ and some integer m .*
3. *Any indecomposable parity complex supported on \bar{X}_λ and extending $\mathcal{L}[m]$ is isomorphic to \mathcal{F} .*

The proof of the theorem is given in [JMW, 2.12].

Definition 3.1.3. *A parity sheaf is an indecomposable parity complex with support \bar{X}_λ and extending $\mathcal{L}[m]$ for some indecomposable $\mathcal{L} \in \text{Loc}_{f,H}(X_\lambda)$ and for some $m \in \mathbb{Z}$. When such a complex exists we denote it by $\mathcal{E}(X_\lambda, \mathcal{L})$ or $\mathcal{E}(\lambda, \mathcal{L})$ and this has the property that $\mathcal{E}(\lambda, \mathcal{L})|_{X_\lambda} = \mathcal{L}[\dim X_\lambda]$. This is the unique parity sheaf associated with (λ, \mathcal{L}) up to shift.*

Remark 3.1.4. 1. *If \mathcal{L} is not indecomposable then $\mathcal{E}(\lambda, \mathcal{L})$ denotes the direct sum of parity complexes coming from the direct summand of \mathcal{L} .*

2. *If $\mathcal{L} = \mathbb{k}_{X_\lambda}$, then we may write $\mathcal{E}(\lambda, \mathcal{L})$ as $\mathcal{E}(\lambda)$.*

3.2. Torsion primes and pretty good primes

Let G be a reductive group with the root datum $(\mathbf{X}, \Phi, \mathbf{Y}, \Phi^\vee)$. A reductive subgroup of G is called regular if it contains a maximal torus. If the group G is complex re-

ductive then all the regular reductive subgroups are in bijection with \mathbb{Z} -closed subsystems of Φ , that is $\Phi_1 \subset \Phi$.

Definition 3.2.1. *A prime p is called a torsion prime for G if for some regular reductive subgroup H of G , $\pi_1(H)$ has p -torsion.*

Definition 3.2.2. *A prime p is called pretty good for G if for all subsets $\Phi_1 \subset \Phi$, $\mathbf{X}/\mathbb{Z}\Phi_1$ and $\mathbf{Y}/\mathbb{Z}\Phi_1^\vee$ have no p -torsion.*

The properties of reductive groups for which p is a pretty good prime have been discussed in [Her, Remark 5.4]. From those properties and using the tables of centralisers from [Car], we have the following lemma.

Lemma 3.2.3. *A prime p is pretty good for G if and only if for all $x \in \mathcal{N}$, p is not a torsion prime for C_x , where C_x is the maximal reductive quotient of $(G^x)^\circ$, and the order of $A_G(x)$ is invertible in \mathbb{k} .*

This is the right time to make some assumption on the characteristic of \mathbb{k} .

Assumption 3.2.4. *1. The characteristic l of \mathbb{k} is a pretty good prime for G .*

2. The field \mathbb{k} is big enough for G ; i.e, for every Levi subgroup L of G and pair $(C_L, \mathcal{E}_L) \in \mathcal{S}(L)$, the irreducible L -equivariant \mathbb{k} -local system \mathcal{E}_L is absolutely irreducible.

Remark 3.2.5. *Note that pretty good implies that $|A_G(x)|$ is invertible in \mathbb{k} . Hence pretty good implies rather good. So if conjecture 2.2.7 holds then it in particular holds for pretty good primes.*

Remark 3.2.6. *From [AJHR, Lemma 2.2(1)], if a prime l is rather good for G then it is rather good for all the Levi subgroups. Hence $|A_L(x)|$ is still invertible in \mathbb{k} for any Levi subgroup $L \subset G$.*

Recall that nilpotent orbits are even dimensional[CM, 1.4]. As a direct conse-

quence of the above lemma, we have the next theorem.

Theorem 3.2.7. *Let C be a nilpotent orbit in \mathfrak{g} and $\mathcal{L} \in \text{Loc}_{f,G}(C, \mathbb{k})$, then $H_G^*(\mathcal{L})$ vanishes in odd degrees.*

Proof. The proof is given in [JMW, Lemma 4.17] □

Theorem 3.2.8. *Let \mathcal{O} be a G_0 -orbit in \mathfrak{g}_n and $\mathcal{L} \in \text{Loc}_{f,G_0}(\mathcal{O}, \mathbb{k})$, then $H_{G_0}^*(\mathcal{L})$ vanishes in odd degrees.*

The proof of this theorem will be given in section 6.

Now, talking about parity sheaves makes sense both in G - and G_0 -equivariant settings as we know (1.1) is true for \mathcal{N}_G and \mathfrak{g}_n .

Conjecture 3.2.9. *Let P be a parabolic subgroup of G and L be its Levi subgroup. For a pair $(C, \mathcal{E}) \in \mathcal{I}(L)^{0\text{-cusp}}$, $\text{Ind}_P^G \mathcal{IC}(C, \mathcal{E})$ is a parity complex.*

In characteristic 0, the proof follows from the decomposition theorem and [Lu2, 24.8]. In positive characteristic, the result is still unknown. Throughout this paper we will assume this result is true. In the last section we will give some example where the conjecture holds.

3.3. Existence of parity for nilpotent cone

It is known that in characteristic 0, intersection cohomology complexes on nilpotent cone coincide with the parity sheaves, thus the existence of parity sheaves are assured.

However, in positive characteristic with the assumption 3.2.4 on the field coefficients, Theorem 3.2.7 assures that there exists atmost one parity sheaf for each pair $(\mathcal{O}, \mathcal{L}) \in \mathcal{I}(G)$ and here we will discuss this existence. By [JMW, prop. 2.34], a even morphism sends

parity complexes to parity complexes. By [DCLP], the Springer resolution,

$$\pi : \tilde{\mathcal{N}} = G \times^P \mathfrak{u} \rightarrow \bar{\mathcal{O}}_{reg} = \mathcal{N}$$

is semismall and even. Therefore $\mathcal{E}(\mathcal{O}_{reg}, \mathbb{k})$ exists. Also $\pi_* \mathbb{k}_{\tilde{\mathcal{N}}}[\dim \mathcal{N}]$ is perverse and parity. So each pair $(\mathcal{O}, \mathcal{L})$ for which $\mathcal{E}(\mathcal{O}, \mathcal{L})$ occurs as direct summand of $\pi_* \mathbb{k}_{\tilde{\mathcal{N}}}[\dim \mathcal{N}]$ with non-zero multiplicity, exists. With the condition that $|W|$, the order of the Weyl group, is invertible in \mathbb{k} , these pairs will be the same with those occurring in characteristic 0. For proving the existence of $\mathcal{E}(\mathcal{O}, \mathbb{k})$ for any nilpotent orbit \mathcal{O} , we need a resolution like above. In case of GL_n , for each orbit \mathcal{O} , it has been proved in [BO], that there exists an GL_n -equivariant semi-small resolution with each fiber obtaining a affine paving. Which provides the requirement.

3.3.1. Classical groups

In case of classical groups we need some more works to do, to find the resolution needed. We fix an orbit \mathcal{O} and an element $x \in \mathcal{O} \cap \mathfrak{u}$. By Jacobson-Morozov theorem [CM], we find a \mathfrak{sl}_2 -triple (x, h, y) in \mathfrak{g} . The element h induces a grading on \mathfrak{g} , which we have discussed in details in the later sections. Under this grading x lies in \mathfrak{g}_2 and $y \in \mathfrak{g}_{-2}$. P be the standard parabolic associated to the simple roots lying in \mathfrak{g}_0 . We consider the natural map,

$$\pi_{\mathcal{O}} : G \times^P \mathfrak{g}_{\geq 2} \rightarrow \bar{\mathcal{O}}.$$

This map is a proper morphism. The existence of $\mathcal{E}(\mathcal{O}, \mathbb{k})$ boils down to proving the above map is even. In [Fr], Fresse has proved the following theorem,

Theorem 3.3.1. *For a classical group G with parabolic subgroup P and i a P -stable ideal*

in the Lie Algebra \mathfrak{p} , the following morphism is even,

$$\pi_{P,i} : G \times^P i \rightarrow \mathfrak{g}.$$

This theorem serves our purpose with $i = \mathfrak{g}_{\geq 0}$, which is clearly a P -stable ideal. So for each orbit \mathcal{O} , the resolution $\pi_{\mathcal{O}}$ provides the existence of parity for a larger collection of pairs but still not for all.

3.4. Existence of parity on \mathfrak{g}_n .

In [JMW], Juteau-Mautner-Williamson gave the following definition,

Definition 3.4.1. *Let $X = \coprod_{\lambda \in \Lambda} X_\lambda$ and $Y = \coprod_{\mu \in \Delta} Y_\mu$ be two stratified H -spaces. A morphism $f : X \rightarrow Y$ is called **stratified** if $f^{-1}(Y_\mu)$ is union of strata and the restriction of f , $f|_{X_\lambda} : X_\lambda \rightarrow Y_\mu$ is a submersion with smooth fibers. The map f is called **even** if for any $y \in Y$ and \mathcal{L} a local system on $f^{-1}(y)$, we have,*

$$H_H^*(f^{-1}(y), \mathcal{L}) = 0 \text{ for odd degrees.}$$

The following theorem is proved in [JMW],

Theorem 3.4.2. *f be a proper and even morphism and \mathcal{E} be a parity complex then $f_*\mathcal{F}$ is again a parity complex.*

A direct consequence of the above theorem is that for X_λ , if we can show that there exist a even and proper resolution

$$\pi : \tilde{X}_\lambda \rightarrow \bar{X}_\lambda$$

and a parity sheaf $\tilde{\mathcal{F}}$ on \tilde{X}_λ with the property $\tilde{\mathcal{F}}|_{\pi^{-1}(X_\lambda) \cong X_\lambda} \cong \mathcal{L}[d_\lambda]$, where \mathcal{L} is a local system on X_λ and d_λ is the dimension of X_λ , then we can say $\mathcal{E}(X_\lambda, \mathcal{L})$ exists. So we can reduce the question of existence of parity to the existence of a proper, even morphism.

For G a complex algebraic group and X and Y , complex algebraic varieties and f be a G -equivariant proper map, the fibers of f has a specific geometric description: they are the flag versions of quiver Grassmannians. Let $\Gamma = (I, J)$ be a quiver with I being the vertex set and J being the *arrows*. Fix an increasing series of dimension vectors $v = (v_1, \dots, v_d)$. Let $v = v_d$ and V be a Γ -representation with $\dim v$. A flag version of quiver Grassmannian is defined as,

$$\mathcal{F}_v(V) = \{V_{v_1} \subset V_{v_2} \subset \dots \subset V_{v_d} = V\}.$$

The fiber $f^{-1}(y)$ as discussed above is always of the form $\mathcal{F}_v(V)$. In his paper [Ma], Maksimau has proved,

Theorem 3.4.3. Γ be a Dynkin quiver of type A, D and E . Then,

$$H^*(\mathcal{F}_v(V), \mathbb{Z}) = 0 \text{ for odd degrees}$$

and is free.

3.4.1. Our setting

In our set-up, we want to study $D_{G_0}^b(\mathfrak{g}_n, \mathbb{k})$, where G_0 is the centralizer of a fixed co-character χ and \mathfrak{g}_n is the n -th graded piece of the action of G_0 on \mathfrak{g} , coming from adjoint action. Here \mathbb{k} is a field of characteristic > 0 . For $G = GL_n$, let the co-character is of the following form,

$$\begin{pmatrix} t^{a_1} & & & \\ & \ddots & & \\ & & & t^{a_r} \end{pmatrix}$$

where $a_1 \geq a_2 \dots \geq a_k$. We can rewrite the a_i 's in blocks of equal entries like following,

$$\underbrace{b_1 \dots b_1}_{m_1 \text{ times}} > \dots > \underbrace{b_k, \dots, b_k}_{m_k \text{ times}}$$

where, $n = m_1 + \dots + m_k$. In this case the above defined G_0 is $GL_{m_1} \times \dots \times GL_{m_k}$ and \mathfrak{g}_n is the collection of matrices with nonzero entries in all (i, j) -th position if $b_i - b_j = n$. For detailed example, see 8.3. We can split \mathbb{C}^n in the collection of vector spaces V_1, \dots, V_k of dimension m_1, \dots, m_k respectively. We can write \mathfrak{g}_n as the collection of maps $\mathbb{C}^n \rightarrow \mathbb{C}^n$ with V_j being sent to V_i if $b_i - b_j = n$. Let's make a quiver with k -vertices as follows,

$$\bullet_{v_1} \quad \bullet_{v_2} \quad \dots \quad \bullet_{v_k} \cdot$$

We will draw an edge between $V_j \rightarrow V_i$ if $b_i - b_j = n$. So what we will get is a union of type A dynkin diagrams. \mathfrak{g}_n is the collection of representations of the above quiver with dimension vectors (m_1, \dots, m_k) . Therefore we can apply Theorem 3.4.3 to show the existence of parity for any pair $(\mathcal{O}, \mathcal{L}) \in \mathcal{I}(\mathfrak{g}_n)$, with $G = GL_m$.

Chapter 4. Lemmas On Varieties With \mathbb{C}^\times -Action

Let X be a variety defined over \mathbb{C} and G be a connected linear algebraic group which acts on X . In this section all the sheaf coefficients will be considered over \mathbb{k} whose characteristic satisfies Assumption 3.2.4.

4.1. \mathbb{C}^\times -action on varieties

Lemma 4.1.1. *If $\mathcal{F} \in D_G^b(X)$ and $\overline{\mathcal{F}} = \text{For}(\mathcal{F}) \in D_c^b(X)$, then $H_c^*(\overline{\mathcal{F}}) = 0$ implies $H_{G,c}^*(\mathcal{F}) = 0$. Equivalently, $R\Gamma_c(\overline{\mathcal{F}}) = 0$ implies $R\Gamma_c(\mathcal{F}) = 0$*

Proof. $R\Gamma_c(\overline{\mathcal{F}}) = \text{For}(R\Gamma_c\mathcal{F})$. Now the argument follows from the following statement, for X as defined above and $\mathcal{M} \in D_G^b(X)$, if $\text{For}(\mathcal{M}) = 0$ then $\mathcal{M} = 0$. Suppose $\mathcal{M} \neq 0$. We will proceed by induction on the i , such that $H^i(\mathcal{M}) \neq 0$. If there is unique i , then $\mathcal{M} \cong H^i(\mathcal{M})$, which up to a shift, belongs to $\text{Perv}_G(X)$. But we know that $\text{For} : \text{Perv}_G(X) \rightarrow \text{Perv}(X)$ is faithful. Hence $\text{For}(\mathcal{M}) \neq 0$.

If there are more than one i , then we use the distinguished triangle,

$$\tau^{\leq k}(\mathcal{M}) \rightarrow \mathcal{M} \rightarrow \tau^{\geq k+1}(\mathcal{M}) \rightarrow . \quad (1.1)$$

So using the forgetful functor we get,

$$\text{For}(\tau^{\leq k}(\mathcal{M})) \rightarrow \text{For}(\mathcal{M}) \rightarrow \text{For}(\tau^{\geq k+1}(\mathcal{M})) \rightarrow . \quad (1.2)$$

As $\mathcal{M} \neq 0$, either the first or the third term in the long exact sequence in (1.1) should be nonzero. By induction in (1.2), either the first or the third term is nonzero. Hence $\text{For}(\mathcal{M}) \neq 0$. □

Lemma 4.1.2. *Let V be a finite-dimensional vector space with a nontrivial linear \mathbb{C}^\times -action on it. Then there exists a nonzero vector with a stabilizer of minimum size and*

there exists a nonzero vector with a stabilizer of maximum size among those with finite stabilizers.

Proof. As \mathbb{C}^\times acts on a vector space V , we have a grading on V given by,

$$V = \bigoplus_{\lambda \in \mathbb{Z}} V_\lambda,$$

where V_λ 's are eigenspaces of the \mathbb{C}^\times -action. Note that stabilizers are subgroups of \mathbb{C}^\times .

If they are finite then they are cyclic and of the form $\mathbb{Z}/n\mathbb{Z}$. So it is obvious that there exists an element of minimum-sized stabilizer. Now let $v \in V$ and $v \notin V_0$. Then we can write v as: $v = \bigoplus v_\lambda$, where $v_\lambda \in V_\lambda$. By definition $t.v_\lambda = t^\lambda v_\lambda$. Hence $t \in \text{Stab}(v_\lambda)$ if and only if t is a λ -root of unity. In other words, $|\text{Stab}(v_\lambda)| = |\lambda|$ for $\lambda \neq 0$. Now if t is in $\text{Stab}(v)$, then t must stabilize all the v_λ 's. Hence t must be c -th root of unity where c divides λ for all $\lambda \neq 0$. Hence $\max |\text{Stab}(v)| = \max\{|\lambda| \mid V_\lambda \neq 0\}$. \square

Lemma 4.1.3. *Let \mathbb{C}^\times acts on Y , a variety over \mathbb{C} , with finite stabilizers. Assume that all the stabilizers have order not divisible by l , where l is the characteristic of \mathbb{k} . Then for any object $\mathcal{F} \in D_{\mathbb{C}^\times}^b(Y, \mathbb{k})$, $\dim(\mathrm{H}_{\mathbb{C}^\times}^*(Y, \mathcal{F})) < \infty$.*

Proof. For a general variety Y , as all the subgroups of \mathbb{C}^\times are finite, there exists a stabilizer of minimum size, say n . Let $U = \{y \in Y, |\text{Stab}(y)| = n\}$. By the Sumihiro embedding theorem [Su], we can cover Y by \mathbb{C}^\times invariant subvarieties, each of which is equivariantly isomorphic to a \mathbb{C}^\times -invariant closed subvariety of \mathbb{A}^N for some N , on which \mathbb{C}^\times acts linearly. Now this action is not trivial as we have already assumed that \mathbb{C}^\times acts non-trivially on Y with finite stabilizers. Hence, by Lemma 4.1.2, these stabilizers will have maximum size. Now the claim is that U is open. Let $Z = U^c = \{y \in Y \mid |\text{Stab}(y)| > n\}$. As

the maximum sized stabilizer exists, we can choose a finite subgroup M of \mathbb{C}^\times which contains all the stabilizers. Let $m \in M$ which is not in the minimum sized stabilizers. Then $Z_m = \{y \in Y | m \in \text{Stab}(y)\}$ is closed. The collection $\{Z_m\}$ is finite as m is coming from a finite subgroup M . Also $Z = \cup_m Z_m$, finite union of closed sets, hence is closed. So U is open.

So we have the open and closed embeddings,

$$Z \xrightarrow{i} Y \xleftarrow{j} U$$

which give us the distinguished triangle,

$$i_* i^! \mathcal{F} \rightarrow \mathcal{F} \rightarrow j_* j^* \mathcal{F} \rightarrow . \quad (1.3)$$

Now we are at a place to use Noetherian induction on Y . The theorem is true for the empty set. So we can assume that it is true for all the proper closed subvarieties of Y , particularly for Z . Now for $u \in U$, let $K = \text{Stab}(u) = \mathbb{Z}/n\mathbb{Z}$. Let $H = \mathbb{C}^\times / K$, then H acts freely on U .

Let $\mathcal{F} \in D_{\mathbb{C}^\times}^b(U, \mathbb{k})$. The goal is to show $\dim H_{\mathbb{C}^\times}^*(\mathcal{F}) < \infty$. According to [BL, Sec 6], if G' and G are two topological groups acting on two varieties X and Y respectively, and $\phi : G' \rightarrow G$ be a homomorphism of topological groups with $f : X \rightarrow Y$, a ϕ -equivariant map, then there exist two functors,

$$Q_f^* : D_G^+(Y) \rightarrow D_{G'}^+(X)$$

and

$$Q_{f*} : D_{G'}^+(X) \rightarrow D_G^+(Y).$$

Here Q_f^* and Q_{f*} are adjoint to each other. In our case, we take both X and Y to be U and $G' = \mathbb{C}^\times$, $G = H$ and we take f to be the identity. So we have,

$$Q_{id*} : D_{\mathbb{C}^\times}^+(U) \rightarrow D_H^+(U).$$

Therefore if $\mathcal{F} \in D_{\mathbb{C}^\times}^b(U)$, then $\text{Hom}_{D_H^+(U)}^*(\mathbb{k}, Q_{id*}\mathcal{F}) \cong \text{Hom}_{D_{\mathbb{C}^\times}^+(U)}^*(\mathbb{k}, \mathcal{F})$. Now if we can show $Q_{id*}\mathcal{F}$ is in $D_H^b(U)$, then as we already know $D_H^b(U) \cong D_c^b(U/H)$ (non-equivariant) as H acts freely on U , so $\dim(\text{Hom}_{D_c^b(U/H)}^*(\mathbb{k}, Q_{id*}\mathcal{F})) < \infty$. The next following fact is from [BL, 7.3]. For $\mathcal{F} \in D_{\mathbb{C}^\times}^b(U) \subset D_{\mathbb{C}^\times}^+(U)$, we have the commutative diagram below,

$$\begin{array}{ccc} D_{\mathbb{C}^\times}^+(U) & \xrightarrow{Q_{id*}} & D_{\mathbb{C}^\times/K}^+(U) \\ \downarrow \text{For}_K^{\mathbb{C}^\times} & & \downarrow \text{For}^{\mathbb{C}^\times/K} \\ D_K^+(U) & \xrightarrow{Q_{id*}} & D^+(U) \end{array}$$

Therefore we have $\text{For}^{\mathbb{C}^\times/K} Q_{id*}\mathcal{F} \cong Q_{id*} \text{For}_K^{\mathbb{C}^\times} \mathcal{F}$. Let $\mathcal{G} = Q_{id*}\mathcal{F}$. So \mathcal{G} is bounded if and only if $\text{For}^H \mathcal{G}$ is bounded. As \mathcal{F} is from bounded derived category then so is $\text{For}_K^{\mathbb{C}^\times} \mathcal{F}$.

Therefore to show $\text{For}^H \mathcal{G}$ is bounded it is enough to show,

$$Q_{id*} : D_K^b(U) \rightarrow D^+(U)$$

takes values in $D^b(U)$. As K is finite hence discrete, so by [BL, Cor 8.4.2], $D_K^b(U) \cong D^b Sh_K(U)$. But we know $Sh_K(U, \mathbb{k}) \cong Sh(U, \mathbb{k}[K])$, where $\mathbb{k}[K]$ is a commutative semisimple ring as $l \nmid |K|$. By the same corollary, Q_{id*} corresponds to a exact functor. So we can conclude Q_{id*} takes $D_K^b(U)$ to $D^b(U)$. Now coming back to our actual proof, if we apply a_{X*} to (1.3), where $a_X : X \rightarrow \{pt\}$, we get,

$$H_{\mathbb{C}^\times}^*(i^! \mathcal{F}) \rightarrow H_{\mathbb{C}^\times}^*(\mathcal{F}) \rightarrow H_{\mathbb{C}^\times}^*(j^* \mathcal{F}) \rightarrow$$

As $\dim(H_{\mathbb{C}^\times}^*(i^! \mathcal{F})) < \infty$ by induction hypothesis and $\dim(H_{\mathbb{C}^\times}^*(j^* \mathcal{F})) < \infty$ by the above result. Hence $\dim(H_{\mathbb{C}^\times}^*(\mathcal{F})) < \infty$. □

Lemma 4.1.4. *Let X be a variety over \mathbb{C} and H be a connected linear algebraic group acting trivially on X . Then for $\mathcal{F} \in D_H^b(X, \mathbb{k})$,*

$$H_{H,c}^*(X, \mathcal{F}) \cong H_c^*(X, \mathcal{F}) \otimes H_H^*(pt, \mathbb{k}).$$

Proof. Let $U_n \rightarrow \{pt\}$ be the n -acyclic resolution for $\{pt\}$. Then $U_n \times X \rightarrow X$ is the n -acyclic resolution of X . By [BL, 2.1], if $\mathcal{F} \in D_H^b(X, \mathbb{k})$ this implies $\mathcal{F} \boxtimes \underline{\mathbb{k}}_{U_n/H} \in D_c^b(X \times U_n/H, \mathbb{k})$ such that for $i < n$,

$$\begin{aligned} H_{H,c}^i(X, \mathcal{F}) &\cong \text{Hom}_{D_H^b(pt, \mathbb{k})}(\underline{\mathbb{k}}_{pt}, a_{X!}\mathcal{F}[i]) \\ &\cong \text{Hom}_{D_c^b(U_n/H \times X)}(\underline{\mathbb{k}}, \underline{\mathbb{k}} \boxtimes a_{X!}\mathcal{F}[i]) \\ &\cong H^i(R\Gamma(\underline{\mathbb{k}}_{U_n/H} \boxtimes a_{X!}\mathcal{F})). \end{aligned}$$

As for constructible sheaves all sheaf functors commute with \boxtimes . Therefore we have for $i < n$,

$$\begin{aligned} H^i(R\Gamma(\underline{\mathbb{k}}_{U_n/H} \boxtimes \mathcal{F})) &\cong \bigoplus_{j+k=i} H^j(R\Gamma(\underline{\mathbb{k}}_{U_n/H})) \otimes H^k(R\Gamma(\mathcal{F})) \\ &\cong \bigoplus_{j+k=i} H_H^j(pt, \mathbb{k}) \otimes H^k(X, \mathcal{F}). \end{aligned}$$

So we are done. □

Lemma 4.1.5. *Let Y be an algebraic variety over \mathbb{C} and Y_0 be the fixed point set of this action. Assume that \mathbb{C}^\times acts on $Y - Y_0$ with finite stabilizers and all the stabilizers of $Y - Y_0$ have order not divisible by l . Let $\mathcal{F} \in D_{\mathbb{C}^\times}^b(Y, \mathbb{k})$. If $H_c^j(Y, \mathcal{F}) = 0$ for all j , then $H_c^j(Y_0, \mathcal{F}) = 0$ for all j .*

Proof. Let $\mathcal{F} \in D_{\mathbb{C}^\times}^b(Y)$ and $H_c^j(Y, \mathcal{F}) = 0$. By Lemma 4.1.1, $H_{\mathbb{C}^\times, c}^j(Y, \mathcal{F}) = 0$. Let $Y_1 = Y - Y_0$, then we have the open and closed embeddings,

$$Y_0 \xrightarrow{i} Y \xleftarrow{j} Y_1.$$

This gives us the long exact sequence of \mathbb{C}^\times equivariant cohomology

$$j_! j^* \mathcal{F} \rightarrow \mathcal{F} \rightarrow i_* i^* \mathcal{F} \rightarrow \dots$$

From that we get, $H_{\mathbb{C}^\times, c}^i(Y_0, \mathcal{F}) = H_{\mathbb{C}^\times, c}^{i+1}(Y_1, \mathcal{F})$, for all i . By Lemma 4.1.3, $\dim(H_{\mathbb{C}^\times, c}^*(Y_1, \mathcal{F}))$ is finite. Therefore we can conclude that $\dim(H_{\mathbb{C}^\times, c}^*(Y_0, \mathcal{F}))$ is also finite. By Lemma 4.1.4,

$$H_{\mathbb{C}^\times, c}^*(Y_0, \mathcal{F}) \cong H_c^*(Y_0, \mathcal{F}) \otimes H_{\mathbb{C}^\times}^*(pt, \mathbb{k}). \quad (1.4)$$

Recall, $H_{\mathbb{C}^\times}^*(pt, \mathbb{k}) \cong \text{Sym}(\mathbb{k})$, which is infinite dimensional. In equation (1.4), if

$H_c^*(Y_0, \mathcal{F}) \neq 0$ then LHS is finite dimensional and RHS is infinite dimensional, a contradiction. So $H_c^*(Y_0, \mathcal{F}) = 0$. □

Lemma 4.1.6. *Let $\mathcal{G} \in D_{\mathbb{C}^\times}^b(pt, \mathbb{k})$.*

1. *If $H^i(\text{For}^{\mathbb{C}^\times}(\mathcal{G})) = 0$, for all i odd, Then $\text{Hom}(\mathbb{k}_{pt}, \mathcal{G}[i]) = 0$, for all i odd.*
2. *If $\text{Hom}^*(\mathbb{k}, \mathcal{G})$ is free over $H_{\mathbb{C}^\times}^*(pt)$ and 0 for odd degrees, then $H^*(\text{For}^{\mathbb{C}^\times}(\mathcal{G})) = 0$ for odd degrees.*

Proof. 1. By Lemma 4.1.1, $H^i(\mathcal{G}) = 0$ for i odd. If $H^i(\mathcal{G})$ is nonzero for a unique i , then $\mathcal{G} = \mathbb{k}_{pt}[i]$, and clearly the statement is true. If there is more than one nonzero cohomology, then we will use induction on the number of nonzero cohomology sheaves and we will use truncation on \mathcal{G} to reduce to the case, $\mathcal{G} = \bigoplus \mathbb{k}_{pt}[2m]$. Hence,

$$\text{Hom}(\mathbb{k}_{pt}, \mathcal{G}[i]) = \bigoplus_m \text{Hom}(\mathbb{k}_{pt}, \mathbb{k}_{pt}[i + 2m]) = \bigoplus_m H_{\mathbb{C}^\times}^{i+2m}(pt),$$

which is zero when i is odd.

2. Note $H_{\mathbb{C}^\times}^*(\mathcal{G})$ is free over $H_{\mathbb{C}^\times}^*(pt)$. So, using the fact that $H_{\mathbb{C}^\times}^*(\mathcal{G})$ is 0 in odd degrees, we can choose basis elements, $\gamma_i \in H_{\mathbb{C}^\times}^{-2n_i}(\mathcal{G})$ for $i = 1, \dots, k$. Therefore γ_i is a

map from $\mathbb{k}_{pt}[2n_i]$ to \mathcal{G} . Hence we can define

$$\gamma : \mathbb{k}_{pt}[2n_1] \oplus \cdots \oplus \mathbb{k}_{pt}[2n_k] \rightarrow \mathcal{G}.$$

This map induces an isomorphism in equivariant cohomology. Let $\mathcal{F} = Cone(\gamma)$. Now the claim is that $\mathcal{F} \cong 0$. If $\mathcal{F} \neq 0$, then let k be the smallest integer such that $H^k(\mathcal{F}) \neq 0$. As $H^k(\mathcal{F}) \in Loc_{f, \mathbb{C}^\times}(pt, \mathbb{k})$ which is again equivalent to finite-dimensional \mathbb{k} -vector spaces, so there is a nonzero map $\mathbb{k}_{pt} \rightarrow H^k(\mathcal{F})$. Now $H^k(\mathcal{F})[-k] \cong \tau^{\leq k} \mathcal{F}$. Hence we have a nonzero map,

$$\mathbb{k}_{pt}[-k] \rightarrow \tau^{\leq k} \mathcal{F} \rightarrow \mathcal{F}.$$

In other words, $H_{\mathbb{C}^\times}^k(\mathcal{F}) \cong \text{Hom}(\mathbb{k}_{pt}[-k], \mathcal{F}) \neq 0$, which is a contradiction. Hence $cone(\gamma) = 0$ and γ is an isomorphism. Therefore,

$$H^j(\text{For}(\mathcal{G})) \cong \bigoplus_{i=1}^k H^{j+2n_i}(\mathbb{k}_{pt}),$$

which is 0 for j odd. □

Theorem 4.1.7. *Let X be a \mathbb{C} -variety with a \mathbb{C}^\times -action on it. Let $X^{\mathbb{C}^\times}$ be the fixed point set and $\mathcal{F} \in D_{\mathbb{C}^\times, c}^b(X, \mathbb{k})$, a local system. If $H_c^a(X, \mathcal{F}) = 0$ for a odd, then $H_c^a(X^{\mathbb{C}^\times}, \mathcal{F}) = 0$ for a odd, provided characteristic l of \mathbb{k} does not divide the order of the stabilizers on $X - X^{\mathbb{C}^\times}$.*

Proof. Let $Z = X^{\mathbb{C}^\times}$ and $U = X - Z$. Let i, j be the inclusion maps of Z and U respectively, and,

$$Z \xrightarrow{i} X \xleftarrow{j} U.$$

Also let $a : X \rightarrow \{pt\}$. Let $\mathcal{G} = a_!(\mathcal{F})$. We have the distinguished triangle below

$$j_! \mathcal{F}|_U \rightarrow \mathcal{F} \rightarrow i_! i^* \mathcal{F} \rightarrow \quad .$$

We can apply $a_!$ to this, and get,

$$\mathcal{G} \rightarrow a_! i^* \mathcal{F} \rightarrow a_!(\mathcal{F}|_U)[1] \rightarrow \quad .$$

Now,

$$H^i(\text{For}(\mathcal{G})) = H_c^i(X, \mathcal{F}) = 0 \text{ for } i \text{ odd.}$$

Hence by Lemma 4.1.6, $H_{\mathbb{C}^\times}^i(\mathcal{G}) = 0$ for i odd. This implies that the map,

$$H_{\mathbb{C}^\times}^i(a_! i^* \mathcal{F}) \rightarrow H_{\mathbb{C}^\times}^{i+1}(\mathcal{F}|_U)$$

is injective for i odd. As the characteristic of k does not divide the order of the stabilizers on $X - X^{\mathbb{C}^\times}$, by lemma 4.1.3, $\dim(H_{\mathbb{C}^\times}^*(\mathcal{F}|_U)) < \infty$. The claim is that $H_{\mathbb{C}^\times}^*(a_! i^* \mathcal{F})$ is free over $H_{\mathbb{C}^\times}^*(pt)$. $i^* \mathcal{F} \in D_{\mathbb{C}^\times}^b(Z)$. If \mathcal{F} is a local system then $i^* \mathcal{F}$ is also a local system. Let $\mathcal{E} = i^* \mathcal{F}$ is a local system on Z , where \mathbb{C}^\times acts on Z trivially.

Hence by Lemma 4.1.4,

$$H_{\mathbb{C}^\times}^*(a_! \mathcal{E}) \cong H_{\mathbb{C}^\times}^*(pt) \otimes a_! \mathcal{E}$$

and this is free over $H_{\mathbb{C}^\times}^*(pt)$. So in our context $H_{\mathbb{C}^\times}^{\text{odd}}(a_! \mathcal{E})$ is either is 0 or infinite-dimensional. If infinite-dimensional, then it is a contradiction because it has an injective map to a finite dimensional cohomology. Hence it must be 0 for odd degrees, that is $H_{\mathbb{C}^\times}^*(Z, \mathcal{F}) = 0$ for odd degrees. Now by Lemma 4.1.6(2), $H_c^*(Z, \mathcal{F}) = 0$ for odd degrees. □

Theorem 4.1.8. *Let M be an object in $D_{\mathbb{C}^\times}^b(pt)$. Assume that $H_{\mathbb{C}^\times}^*(M)$ is finite-dimensional, then the Euler characteristic of $H^*(M)$ (nonequivariant cohomology) is 0.*

Proof. From [BL, Th. 3.7.1],

$$\text{For}^{\mathbb{C}^\times} : D_{\mathbb{C}^\times}^b(pt) \rightarrow D^b(pt)$$

has a left adjoint $\text{Ind}_!$. Let $a : \mathbb{C}^\times \times pt \rightarrow pt$ be the projection on pt , which is the constant map in this case and $\nu : pt \rightarrow \mathbb{C}^\times \times pt$, the inclusion map. Here $\nu^* \text{For}^{\mathbb{C}^\times}[-2]$ is the induction equivalence map. Then the formula for $\text{Ind}_!$ is $a_! \mathbb{D}(\nu^* \text{For}^{\mathbb{C}^\times}[-2])^{-1} : D^b(pt) \rightarrow D_{\mathbb{C}^\times}^b(pt)$, where \mathbb{D} is the equivariant Verdier duality. Therefore $\text{Ind}_!(\underline{\mathbb{k}}_{pt}) = R\Gamma_c(\underline{\mathbb{k}}_{\mathbb{C}^\times}[2])$. So,

$$H^i(\text{Ind}_! \underline{\mathbb{k}}_{pt}) \begin{cases} \cong \mathbb{k} \text{ for } i = 0, -1 \\ \cong 0 \text{ otherwise.} \end{cases} \quad (1.5)$$

We have the distinguished triangle,

$$\tau^{\leq -1} \text{Ind}_!(\underline{\mathbb{k}}_{pt}) \rightarrow \text{Ind}_!(\underline{\mathbb{k}}_{pt}) \rightarrow \tau^{\geq 0} \text{Ind}_!(\underline{\mathbb{k}}_{pt}) \rightarrow \dots$$

Using (1.5) this distinguished triangle reduces to

$$\mathbb{k}[1] \rightarrow \text{Ind}_!(\underline{\mathbb{k}}_{pt}) \rightarrow \mathbb{k} \rightarrow \dots$$

Note that, $\text{Hom}(\text{Ind}_! \underline{\mathbb{k}}_{pt}, M) \cong \text{Hom}(\underline{\mathbb{k}}_{pt}, \text{For}^{\mathbb{C}^\times}(M)) \cong H^i(M)$ and $\text{Hom}(\underline{\mathbb{k}}_{pt}, M) \cong H_{\mathbb{C}^\times}^i(M)$. Now we apply $\text{Hom}(-, M)$ to the above distinguished triangle and get the long exact sequence,

$$\rightarrow H_{\mathbb{C}^\times}^{i-1}(M) \rightarrow H^i(M) \rightarrow H_{\mathbb{C}^\times}^i(M) \rightarrow \dots$$

From the assumption on M , this LES have finitely many terms. Therefore,

$$\chi(H_{\mathbb{C}^\times}^{i-1}(M)) + \chi(H_{\mathbb{C}^\times}^i(M)) = \chi(H^i(M)).$$

Here χ denotes the Euler characteristics. But $\chi(H_{\mathbb{C}^\times}^{i-1}(M)) = -\chi(H_{\mathbb{C}^\times}^i(M))$, so the left hand side is 0. So $\chi(H^i(M)) = 0$ and we are done.

□

Chapter 5. Induction and Restriction

Induction and restriction in graded set-up

We have already talked about the induction diagram for nilpotent cone in 2.2. Similarly we can define induction diagram in graded set-up. For that we need to choose a parabolic subgroup P containing $\chi(\mathbb{C}^\times)$ and a Levi, $L \subset P$. Using the diagram we can define $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}}$ and $\text{Res}_{\mathfrak{p}}^{\mathfrak{g}}$, these two functors. $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}}$ is going to play an important role throughout this entire thesis, which commutes with the Verdier duality. Another major property is that, like on nilpotent cone, restriction is left adjoint to induction. Transitivity of $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}}$ plays a key factor in the proof of Theorem 7.4.2, where we prove the existence of parity on graded pieces.

Lusztig's definition matches with us

In [Lu], Lusztig has defined the induction diagram and $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}}$ for the first time for graded set-up. The induction diagram and the functor $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}}$ that we defined is slightly different in this thesis. It has much more similarity with the already defined induction functor on nilpotent cone. So using several base change diagram we prove in Lemma 5.2.1 that Lusztig's $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}}$ matches with our $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}}$, which is much more convenient to use in this particular thesis.

For cuspidal, \mathcal{IC} is parity

In characteristic 0, it is a well-known fact that \mathcal{IC} coincides with parity for any pair on the nilpotent cone. Lusztig has proved that in characteristic 0, \mathcal{IC} coincides with parity on graded pieces. This result was still unknown in positive characteristic, both on nilpotent cone and graded pieces. In this thesis we have already assumed that Mautner's

cleanness conjecture (Conjecture 2.2.7) holds in our set-up. using this assumption we imitate the proof of Lusztig [Lu3] for cleanness of cuspidal pair in characteristic 0 and show that in positive characteristic, \mathcal{IC} 's are clean for cuspidal pair on graded pieces. Which eventually shows \mathcal{IC} and parity coincide on \mathfrak{g}_n . therefore parity exists for cuspidal pairs.

5.1. Induction and restriction

Let P be a parabolic subgroup of G containing $\chi(\mathbb{C}^\times)$. Let L and U be a Levi subgroup and the unipotent radical, respectively. We can choose L so that χ gets mapped in to L . Let $\mathfrak{p}, \mathfrak{l}, \mathfrak{n}$ be the Lie Algebras of P, L, U respectively. Then $\mathfrak{p}, \mathfrak{l}, \mathfrak{n}$ inherit grading from \mathfrak{g} :

$$\mathfrak{p} = \bigoplus_{n \in \mathbb{Z}} \mathfrak{p}_n, \mathfrak{l} = \bigoplus_{n \in \mathbb{Z}} \mathfrak{l}_n, \mathfrak{n} = \bigoplus_{n \in \mathbb{Z}} \mathfrak{n}_n,$$

where $\mathfrak{p}_n = \mathfrak{p} \cap \mathfrak{g}_n, \mathfrak{n}_n = \mathfrak{n} \cap \mathfrak{g}_n$ and $\mathfrak{l}_n = \mathfrak{p}_n / \mathfrak{n}_n$. From now on the composition of $\chi : \mathbb{C}^\times \rightarrow P$ and $P \twoheadrightarrow P/U = L$ will also be denoted by $\chi : \mathbb{C}^\times \rightarrow L$.

Let's recall the induction diagram from 2.2

$$\mathcal{N}_L \xleftarrow{\pi_P} \mathcal{N}_L + \mathfrak{u}_P \xrightarrow{e_P} G \times^P (\mathcal{N}_L + \mathfrak{u}_P) \xrightarrow{\mu_P} \mathcal{N}_G,$$

where $\mathfrak{u}_P = \text{Lie}(U_P)$, π_P, e_P are the obvious maps and $\mu_P(g, x) = \text{Ad}(g)x$. A slight modification of this diagram gives us the induction diagram in the graded setting.

$$\begin{array}{ccccc} \mathfrak{l}_n & \xleftarrow{\pi} & \mathfrak{p}_n & \xrightarrow{e} & G_0 \times^{P_0} \mathfrak{p}_n & \xrightarrow{\mu} & \mathfrak{g}_n \\ & & & \searrow & \text{---} & \nearrow & \\ & & & & i & & \end{array}$$

As before, π is projection, e, i are inclusions and $\mu(g, x) = \text{Ad}(g)x$. The induction functor is denoted by

$$\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} : D_{L_0}^b(\mathfrak{l}_n) \rightarrow D_{G_0}^b(\mathfrak{g}_n).$$

As $P_0 = L_0 \times U_0$ and U_0 acts on \mathfrak{l}_n trivially, we have equivalence of categories $D_{P_0}^b(\mathfrak{l}_n) \cong D_{L_0}^b(\mathfrak{l}_n)$. So instead of starting from $D_{L_0}^b(\mathfrak{l}_n)$ we can start from $D_{P_0}^b(\mathfrak{l}_n)$. So we define

$$\mathrm{Ind}_{\mathfrak{p}}^{\mathfrak{g}}(\mathcal{F}) := \mu_1(\underbrace{e^* \mathrm{For}_{P_0}^{G_0}}_{\text{Induction Equivalence}})^{-1} \pi^*(\mathcal{F}).$$

Here $e^* \mathrm{For}_{P_0}^{G_0} : D_{G_0}^b(G_0 \times^{P_0} \mathfrak{p}_n) \rightarrow D_{P_0}^b(\mathfrak{p}_n)$ is the induction equivalence map, hence its inverse makes sense. The definition of restriction also comes from the diagram above, $\mathrm{Res}_{\mathfrak{p}}^{\mathfrak{g}} : D_{G_0}^b(\mathfrak{g}_n) \rightarrow D_{L_0}^b(\mathfrak{l}_n)$ is defined by,

$$\mathrm{Res}_{\mathfrak{p}}^{\mathfrak{g}}(\mathcal{F}) := \pi_1 \iota^* \mathrm{For}_{L_0}^{G_0}(\mathcal{F}).$$

Theorem 5.1.1. *The functor $\mathrm{Ind}_{\mathfrak{p}}^{\mathfrak{g}}$ commutes with \mathbb{D} , the Verdier duality functor.*

Proof. The map μ is proper, therefore it commutes with \mathbb{D} . By [BL, prop. 7.6.2], $\mathbb{D}(e^* \mathrm{For}_{P_0}^{G_0}) = (e^* \mathrm{For}_{P_0}^{G_0})\mathbb{D}[-2 \dim G_0/P_0]$. The map π is smooth and has relative dimension of $\dim \mathfrak{p}_n - \dim \mathfrak{l}_n = 2 \dim G_0/P_0$. Therefore $\mathbb{D}\pi^* = \pi^!\mathbb{D} = \pi^*\mathbb{D}[2 \dim G_0/P_0]$. Combining all these facts we can see, $\mathbb{D}\mathrm{Ind}_{\mathfrak{p}}^{\mathfrak{g}} = \mathrm{Ind}_{\mathfrak{p}}^{\mathfrak{g}}\mathbb{D}$. □


5.1.1. Transitivity

Before going into the main result of this section we will talk about the transitivity of induction. Let P be a parabolic subgroup of G containing the Levi subgroup L which contains $\chi(\mathbb{C}^\times)$. Let R be a parabolic contained in P with Levi $M \subset L$, which again contains $\chi(\mathbb{C}^\times)$. Then $R \cap L$ is a parabolic subgroup of L with the Levi factor M . Let $\mathfrak{r}, \mathfrak{m}$ be the Lie Algebras of R, M respectively.

Theorem 5.1.2. *Let $R \subset P$ and $M \subset L$ as defined above. Then for $\mathcal{F} \in D_{M_0}^b(\mathfrak{m}_n)$,*

$$\mathrm{Ind}_{\mathfrak{p}}^{\mathfrak{g}} \mathrm{Ind}_{\mathfrak{r} \cap \mathfrak{m}}^{\mathfrak{l}}(\mathcal{F}) \cong \mathrm{Ind}_{\mathfrak{r}}^{\mathfrak{g}}(\mathcal{F}).$$

Proof. The proof is clear from the diagram below.



□

5.2. Lusztig's original definition

Lusztig's original definition of the restriction is same as we defined above. But for induction, he used a different diagram.

$$\mathfrak{l}_n \xleftarrow{p_1} E' \xrightarrow{p_2} E'' \xrightarrow{p_3} \mathfrak{g}_n$$

where $E' = G_0 \times^{U_0} \mathfrak{p}_n$ and $E'' = G_0 \times^{P_0} \mathfrak{p}_n$. Here $p_1(g, x) = \pi(x)$, p_2 is the obvious map and $p_3(g, x) = Ad(g)x$. Induction is defined by $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}}(\mathcal{F}) =$

$$p_{3!} \left(\underbrace{p_2^* \text{For}_{P_0}^{G_0}}_{\text{Induction Equivalence}} \right)^{-1} p_1^*(\mathcal{F}).$$

Lemma 5.2.1. *Lusztig's original definition of induction matches with the definition given here.*

Proof. It follows from the diagram below.

$$\begin{array}{ccccccc}
 \mathfrak{l}_n & \xleftarrow{\pi} & \mathfrak{p}_n & \xrightarrow{h} & G_0 \times \mathfrak{p}_n & \xrightarrow{q_P} & G_0 \times^{P_0} \mathfrak{p}_n & \xrightarrow{p_3=\mu} & \mathfrak{g}_n \\
 & & & & \downarrow q_U & & \uparrow p_2 & & \\
 & & & & G_0 \times^{U_0} \mathfrak{p}_n & & & &
 \end{array}$$

Clearly,

$$\begin{aligned}
p_{3!}(p_2^* \text{For}_{P_0}^{G_0})^{-1} p_1^*(\mathcal{F}) &= p_{3!} \underbrace{\text{For}_{P_0}^{G_0}{}^{-1} q_P{}^{*-1} q_U^*}_{(p_2^* \text{For}_{P_0}^{G_0})^{-1}} \underbrace{q_U{}^{*-1} h^{*-1} \pi^*}_{p_1^*} \\
&= p_{3!}((q_P \circ h)^* \text{For}_{P_0}^{G_0})^{-1} \pi^*(\mathcal{F}) \\
&= \mu_!(e^* \text{For}_{P_0}^{G_0})^{-1} \pi^*(\mathcal{F}) \\
&= \text{Ind}_{\mathfrak{p}}^{\mathfrak{g}}(\mathcal{F}).
\end{aligned}$$

□

So from now we can use any of the induction diagrams defined above.

5.3. Cleanness for cuspidal pairs

Theorem 5.3.1. $(\mathcal{O}, \mathcal{L}) \in \mathcal{I}(\mathfrak{g}_n)^{\text{cusp}}$ is clean.

Proof. Let $(\mathcal{O}, \mathcal{L}) \in \mathcal{I}(\mathfrak{g}_n)^{\text{cusp}}$ and $(C, \mathcal{E}) \in \mathcal{I}(G)^{0-\text{cusp}}$ so that $C \cap \mathfrak{g}_n = \mathcal{O}$ and $\mathcal{L} = \mathcal{E}|_{\mathcal{O}}$.

Note that \mathcal{E}^\vee is also cuspidal by Remark 2.2.3. Let X be another G_0 -orbit in \mathfrak{g}_n other than \mathcal{O} . We will show that $\mathcal{IC}(\mathcal{O}, \mathcal{L})|_X = 0$ and $\mathcal{IC}(\mathcal{O}, \mathcal{L}^\vee)|_X = 0$. For descending induction, assume it is true for orbits X' , where $\dim(X) < \dim(X') < \dim(\mathfrak{g}_n)$. Let $x \in X$.

By Theorem 2.1.2, we can find

$$\phi : \mathfrak{sl}_2 \rightarrow \mathfrak{g} \text{ such that } \phi(e) = x \in \mathfrak{g}_n, \phi(f) = x' \in \mathfrak{g}_{-n}, \phi(h) \in \mathfrak{g}_0$$

where e, f, h are defined in the background. Let $\tilde{\phi} : SL_2 \rightarrow G$ be such that $d\tilde{\phi} = \phi$. Define

$\chi' : \mathbb{C}^\times \rightarrow G$ by

$$\chi'(a) = \tilde{\phi} \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}.$$

Let $\mathfrak{g}^{x'}$ be the centralizer of x' in \mathfrak{g} . Let $\Sigma = x + \mathfrak{g}^{x'}$ and $\tilde{\Sigma} = \Sigma \cap \mathfrak{g}_n$. According to Slodowy [Sw, pp. 109],

$$\Sigma \text{ is transversal to the } G\text{-orbit of } x \text{ in } \mathfrak{g}. \quad (3.1)$$

Now \mathbb{C}^\times acts on G by conjugation via χ and on \mathfrak{g} by $a^{-n}Ad(\chi(a))$, call it ψ , which fixes x and preserves Σ as $x' \in \mathfrak{g}_n$. The action of G on \mathfrak{g} is \mathbb{C}^\times -equivariant. So we can restrict the action to the fixed point sets of \mathbb{C}^\times -actions and see that G_0 acts on \mathfrak{g}_n . Using (3.1) we deduce that

$$\tilde{\Sigma} \text{ is transverse to the } G_0\text{-orbit of } x \text{ in } \mathfrak{g}_n. \quad (3.2)$$

Now we define another action ψ' , \mathbb{C}^\times acts on Σ by $a \rightarrow a^{-2}Ad(\chi'(a))$. This action is well-defined; if $x + y \in \Sigma$, then $[y, x'] = 0$; so $[Ad(\chi'(a)y, Ad(\chi'(a))x'] = 0$. Let $c_{\chi'(a)}$ denote the conjugation by $\chi'(a)$. Now $Ad(\chi'(a))x' = d(c_{\chi'(a)})d\tilde{\phi}|_f = d(c_{\chi'(a)}\tilde{\phi})|_f = a^{-2}d\tilde{\phi}|_f = a^{-2}x'$. So we have $[Ad(\chi'(a))y, x'] = 0$. Also $Ad(\chi'(a))x = d(c_{\chi'(a)}) \circ d\tilde{\phi}|_e = d(c_{\chi'(a)} \circ \tilde{\phi})|_e = a^2d\tilde{\phi}|_e = a^2x$. Hence $a^{-2}Ad(\chi'(a))(x + y) \in \Sigma$. Now we will show that

$$\text{this action } \psi' \text{ stabilizes } \tilde{\Sigma} \text{ and } \mathcal{O} \cap \Sigma. \quad (3.3)$$

To show the first part it is enough to show that if $y \in \mathfrak{g}_n$, then $Ad(\chi'(a))y \in \mathfrak{g}_n$, because we already have shown that Σ is stable under this action. Now $\phi(h) \in \mathfrak{g}_0$, so the Lie subalgebra generated by $\phi(h)$ is in \mathfrak{g}_0 . Thus by [Hum, Theorem 13.1], $\tilde{\phi} \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \subset G_0$. Therefore χ' commutes with χ and we are done with the proof that $Ad(\chi'(a))y \in \mathfrak{g}_n$. If $y \in \mathcal{O}$, then $Ad(\chi'(a))y$ is also in \mathcal{O} as $Im(\chi') \subset G_0$ and \mathcal{O} is a G_0 -orbit.

Now we can consider a \mathfrak{sl}_2 action on \mathfrak{g} by $(s, v) \in \mathfrak{sl}_2 \times \mathfrak{g}$ goes to $[\phi(s), v]$. Via this action the Lie Algebra generated by $\phi(h)$ acts on $\mathfrak{g}^{x'}$. The unique lift of this action after

multiplying by t^{-2} , where $t \in \mathbb{C}^\times$, gives rise to ψ' , that we talked already. Now in the original action f acts on $\mathfrak{g}^{x'}$ gives 0. Therefore all the eigen values of the action of h on $\mathfrak{g}^{x'}$ will be negative.

$$\text{Hence the action } \psi' \text{ is a repelling action on } \Sigma \text{ to } x. \quad (3.4)$$

$$\text{by Conjecture 2.2.7, } \mathcal{IC}(C, \mathcal{E})|_{\bar{c}-C} = 0. \quad (3.5)$$

As \mathcal{E}^\vee is also cuspidal, the same result is true for \mathcal{E}^\vee .

Using the transversal property of (3.1) and the definition of transversal slice, the map $\mu : G \times \Sigma \rightarrow \mathfrak{g}$ is smooth of relative dimension $\dim G - \dim G \cdot x$.

Hence by [BBD, pp. 110], pullback with a shift takes \mathcal{IC} 's to \mathcal{IC} 's.

$$\begin{array}{ccc} G \times \Sigma & \xrightarrow{\mu} & \mathfrak{g} \\ \uparrow h & \nearrow & \\ \Sigma & & \end{array}$$

where μ is smooth and h induces the induction equivalence, $D^b(\Sigma) \cong D_G^b(G \times \Sigma)$. Hence from the above diagram, we can say that

$$\mathcal{IC}(C, \mathcal{E})|_\Sigma = \mathcal{IC}(C \cap \Sigma, \mathcal{E}|_{C \cap \Sigma})[m] \quad (3.6)$$

where $m = \dim C - \dim(C \cap \Sigma)$. Similarly, $\mathcal{IC}(C, \mathcal{E}^\vee)|_\Sigma = \mathcal{IC}(C \cap \Sigma, \mathcal{E}^\vee|_{C \cap \Sigma})[m]$. By (3.5) and (3.6),

$$\mathcal{IC}(C \cap \Sigma, \mathcal{E}|_{C \cap \Sigma})|_{(\bar{c}-C) \cap \Sigma} = 0 \text{ and } \mathcal{IC}(C \cap \Sigma, \mathcal{E}^\vee|_{C \cap \Sigma})|_{(\bar{c}-C) \cap \Sigma} = 0. \quad (3.7)$$

Using the repelling action from (3.4) and Lemma 2.1.3, [BR, Theorem 1], we get,

$$\begin{cases} \mathcal{IC}(C \cap \Sigma, \mathcal{E}|_{C \cap \Sigma})_x = R\Gamma(\mathcal{IC}(C \cap \Sigma, \mathcal{E}|_{C \cap \Sigma})), \text{ and} \\ \mathcal{IC}(C \cap \Sigma, \mathcal{E}^\vee|_{C \cap \Sigma})_x = R\Gamma(\mathcal{IC}(C \cap \Sigma, \mathcal{E}^\vee|_{C \cap \Sigma})). \end{cases} \quad (3.8)$$

Now $x \in (\bar{C} - C) \cap \Sigma$, so from (3.7),

$$\mathcal{IC}(C \cap \Sigma, \mathcal{E}|_{C \cap \Sigma})_x = 0. \text{ So by (3.8), } R\Gamma(\mathcal{IC}(C \cap \Sigma, \mathcal{E}|_{C \cap \Sigma})) = 0. \quad (3.9)$$

This implies,

$$\begin{cases} R\Gamma_c(\mathcal{IC}(C \cap \Sigma, \mathcal{E}^\vee|_{C \cap \Sigma})) &= R\Gamma_c(\mathbb{D}\mathcal{IC}(C \cap \Sigma, \mathcal{E}|_{C \cap \Sigma})) \\ &= \mathbb{D}R\Gamma(\mathcal{IC}(C \cap \Sigma, \mathcal{E}|_{C \cap \Sigma})) = 0. \end{cases} \quad (3.10)$$

Similarly,

$$R\Gamma(\mathcal{IC}(C \cap \Sigma, \mathcal{E}^\vee|_{C \cap \Sigma})) = 0 \text{ and } R\Gamma_c(\mathcal{IC}(C \cap \Sigma, \mathcal{E}|_{C \cap \Sigma})) = 0. \quad (3.11)$$

Now we claim that

$$R\Gamma_c(\mathcal{E}^\vee|_{C \cap \Sigma}) = 0. \quad (3.12)$$

From the open-closed embedding,

$$C \cap \Sigma \xrightarrow{j} \bar{C} \cap \Sigma \xleftarrow{i} (\bar{C} - C) \cap \Sigma$$

gives us the distinguished triangle,

$$j_!j^*\mathcal{IC}(C \cap \Sigma, \mathcal{E}^\vee|_{C \cap \Sigma}) \rightarrow \mathcal{IC}(C \cap \Sigma, \mathcal{E}^\vee|_{C \cap \Sigma}) \rightarrow i_*i^*\mathcal{IC}(C \cap \Sigma, \mathcal{E}^\vee|_{C \cap \Sigma}) \rightarrow \dots$$

We can apply $R\Gamma_c$ to get

$$R\Gamma_c(j_!j^*\mathcal{IC}(C \cap \Sigma, \mathcal{E}^\vee|_{C \cap \Sigma})) \rightarrow R\Gamma_c(\mathcal{IC}(C \cap \Sigma, \mathcal{E}^\vee|_{C \cap \Sigma})) \rightarrow R\Gamma_c(i_*i^*\mathcal{IC}(C \cap \Sigma, \mathcal{E}^\vee|_{C \cap \Sigma})) \rightarrow \dots$$

The first term in this distinguished triangle is $R\Gamma_c(\mathcal{E}^\vee|_{C \cap \Sigma})$ with a shift. The second term is 0 by (3.10) and third term is 0 by (3.7), hence (3.12) is proved.

For the action ψ of \mathbb{C}^\times on $C \cap \Sigma$ by $a \rightarrow a^{-n} Ad(\chi(a))$, the fixed point set is $\mathcal{O} \cap \tilde{\Sigma}$. So by Lemma 4.1.5,

$$R\Gamma_c(\mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}}) = 0. \quad (3.13)$$

By the transversal property (3.2), we have,

$$\mathcal{IC}(\mathcal{O}, \mathcal{L})|_{\tilde{\Sigma}} = \mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}|_{\mathcal{O} \cap \tilde{\Sigma}})[n], \quad (3.14)$$

and

$$\mathcal{IC}(\mathcal{O}, \mathcal{L}^\vee)|_{\tilde{\Sigma}} = \mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}})[n], \quad (3.15)$$

where $n = \dim \mathcal{O} - \dim \mathcal{O} \cap \tilde{\Sigma}$.

By repelling property (3.4) we have,

$$\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}|_{\mathcal{O} \cap \tilde{\Sigma}})_x = R\Gamma(\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}|_{\mathcal{O} \cap \tilde{\Sigma}})) \quad (3.16)$$

and

$$\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}})_x = R\Gamma(\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}})). \quad (3.17)$$

Here $\overline{\mathcal{O} \cap \tilde{\Sigma}} - \{x\}$ is the union of $V \cap \tilde{\Sigma}$, where each V is a G_0 orbit whose closure contains x , hence also X . So $\dim V > \dim X$. Also $\overline{\mathcal{O} \cap \tilde{\Sigma}} \cap X = \{x\}$. So we can use the induction hypothesis on $\overline{\mathcal{O} \cap \tilde{\Sigma}} - (\mathcal{O} \cap \tilde{\Sigma}) - \{x\}$ and (3.15), therefore,

$$\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}}) \text{ is } 0 \text{ on } \overline{\mathcal{O} \cap \tilde{\Sigma}} - (\mathcal{O} \cap \tilde{\Sigma}) - \{x\}. \quad (3.18)$$

Now we use the open and closed embeddings below for $\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}})|_{\overline{\mathcal{O} \cap \tilde{\Sigma}} - \{x\}}$,

$$\mathcal{O} \cap \tilde{\Sigma} \xrightarrow{j} \overline{\mathcal{O} \cap \tilde{\Sigma}} - \{x\} \xleftarrow{i} \overline{\mathcal{O} \cap \tilde{\Sigma}} - (\mathcal{O} \cap \tilde{\Sigma}) - \{x\}.$$

This gives us the distinguished triangle,

$$j_!j^*\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}})|_{\overline{\mathcal{O} \cap \tilde{\Sigma}} - \{x\}} \rightarrow \mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}})|_{\overline{\mathcal{O} \cap \tilde{\Sigma}} - \{x\}} \rightarrow i_*i^*\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}})|_{\overline{\mathcal{O} \cap \tilde{\Sigma}} - \{x\}} \rightarrow \cdot$$

We have,

$$R\Gamma_c(\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}})|_{\overline{\mathcal{O} \cap \tilde{\Sigma}} - \{x\}}) = 0 \quad (3.19)$$

as the first term in the distinguished triangle, $R\Gamma_c(\mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}})$ vanishes by (3.13) and the third term vanishes by (3.18).

Now from the open-closed embedding,

$$\{x\} \xrightarrow{i} \overline{\mathcal{O} \cap \tilde{\Sigma}} \xleftarrow{j} \overline{\mathcal{O} \cap \tilde{\Sigma}} - \{x\}$$

we get,

$$j_!j^*\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}}) \rightarrow \mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}}) \rightarrow i_*i^*\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}}).$$

By (3.19),

$$\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}})_x = R\Gamma_c\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}}). \quad (3.20)$$

From (3.16),

$$\mathbb{D}\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}|_{\mathcal{O} \cap \tilde{\Sigma}})_x = \mathbb{D}R\Gamma(\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}|_{\mathcal{O} \cap \tilde{\Sigma}})) = R\Gamma_c\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}})$$

Hence from (3.20), we get,

$$\mathbb{D}\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}|_{\mathcal{O} \cap \tilde{\Sigma}})_x = \mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}})_x.$$

Since $\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}})_x$ lives in degrees < 0 . Hence $\mathbb{D}\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}|_{\mathcal{O} \cap \tilde{\Sigma}})_x$ lives in degrees > 0 . But $\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}})_x$ again lives in degrees < 0 , which is a contradiction. So $\mathcal{IC}(\mathcal{O} \cap \tilde{\Sigma}, \mathcal{L}^\vee|_{\mathcal{O} \cap \tilde{\Sigma}})_x = 0$ and by (3.14), $\mathcal{IC}(\mathcal{O}, \mathcal{L})_x = 0$. Hence we are done. \square

Corollary 5.3.2. *For $(\mathcal{O}, \mathcal{L}) \in \mathcal{I}(\mathfrak{g}_n)^{\text{cusp}}$, the parity sheaf $\mathcal{E}(\mathcal{O}, \mathcal{L})$ exists and $\mathcal{IC}(\mathcal{O}, \mathcal{L}) = \mathcal{E}(\mathcal{O}, \mathcal{L})$.*

Proof. From the previous theorem $(\mathcal{O}, \mathcal{L})$ is clean, i.e $\mathcal{IC}(\mathcal{O}, \mathcal{L})$ restricted to $\bar{\mathcal{O}} - \mathcal{O}$ is 0. So $\mathcal{IC}(\mathcal{O}, \mathcal{L}) = j_! \mathcal{L}[\dim \mathcal{O}]$, where $j : \mathcal{O} \hookrightarrow g_n$, which obviously satisfies the parity condition.

Hence by uniqueness of an indecomposable parity complex, $\mathcal{IC}(\mathcal{O}, \mathcal{L}) = \mathcal{E}(\mathcal{O}, \mathcal{L})$. □

Chapter 6. Induction Diagram For Cuspidal Pairs

Special diagram for cuspidal pairs

Mautner's conjecture asserts that \mathcal{IC} 's are clean for cuspidal pairs on nilpotent cone. We have proved in Theorem 5.3.1 that \mathcal{IC} 's are clean, therefore they coincide with parity sheaves and they are the push-forward of the corresponding local system (with a shift) by the inclusion map of the corresponding orbit. Combining these facts and base change diagrams we can redefine the induction diagram both for nilpotent cone and in graded set-up.

$\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}}$ sends parity to parity for cuspidals

Here we prove that for a cuspidal pair, $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}}$ takes parity to parity. We have tried to imitate the same result proved by Lusztig [Lu] in characteristic 0. But because of the failure of decomposition theorem in > 0 characteristic, [Lu, 2.6(b)] fails here, which plays a key role in the proof there. Conjecture 3.2.9 serves as a replacement for the statement 2.6(c) here. Using the definition of cuspidal pair in graded setting, which asserts the existence of a cuspidal pair on nilpotent cone and also the redefined induction diagram in previous section allows us to show that cohomology of the $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}}$ applied to the parity sheaf (for cuspidal) is the direct summand of the cohomology of the stalk of Ind_P^G of the corresponding cuspidal pair on nilpotent cone. Then using the Conjecture 3.2.9 provides the desired result.

6.1. Induction diagram for cuspidal pairs

In this chapter, we first redefine Lusztig's induction diagram for cuspidal pairs. Let P be a parabolic subgroup of G and L, U be its Levi subgroup and the unipotent radical,

respectively. Let $(\mathcal{O}, \mathcal{L}) \in \mathcal{I}(\mathfrak{l}_n)^{\text{cusp}}$. We define the induction diagram to be,

$$\mathcal{O} \xleftarrow{p'_1} \mathcal{O} + \mathfrak{u}_n \xrightarrow{p'_2} G_0 \times^{P_0} (\mathcal{O} + \mathfrak{u}_n) \xrightarrow{p'_3} \mathfrak{g}_n. \quad (1.1)$$

We define $p'_3 : G_0 \times^{P_0} (\mathcal{O} + \mathfrak{u}_n) \rightarrow \mathfrak{g}_n$ to be $p'_3(g, z) = \text{Ad}(g)z$,

$p'_2 : \mathcal{O} + \mathfrak{u}_n \rightarrow G_0 \times^{P_0} (\mathcal{O} + \mathfrak{u}_n)$, p'_2 to be the obvious map and,

$p'_1 : \mathcal{O} + \mathfrak{u}_n \rightarrow \mathcal{O}$ to be $p'_1(z) = \pi(z)$, where $\pi : \mathfrak{p}_n \rightarrow \mathfrak{l}_n$. We start with $(\mathcal{O}, \mathcal{L}) \in \mathcal{I}(\mathfrak{l}_n)^{\text{cusp}}$

and redefine the induction diagram. We define

$$\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}}(\mathcal{I}\mathcal{C}(\mathcal{O}, \mathcal{L})) = p'_{3!}(p'_2{}^* \text{For}_{L_0}^{G_0})^{-1} p'_1{}^*(\mathcal{L}[\dim \mathcal{O}]).$$

Lemma 6.1.1. *This definition of induction for cuspidal pairs coincides with Lusztig's original definition.*

Proof. By Theorem 5.3.1, $(\mathcal{O}, \mathcal{L})$ is clean and it coincides with the Lusztig's definition of induction because of the following commutative diagram.

$$\begin{array}{ccccccc} \mathcal{O} & \xleftarrow{p'_1} & \mathcal{O} + \mathfrak{u}_n & \xrightarrow{p'_2} & G_0 \times^{P_0} (\mathcal{O} + \mathfrak{u}_n) & \xrightarrow{p'_3} & \mathfrak{g}_n \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \mathfrak{l}_n & \xleftarrow{p_1} & E' & \xrightarrow{p_2} & E'' & \xrightarrow{p_3} & \mathfrak{g}_n \end{array}$$

□

6.2. Parity preserved for cuspidal pairs

Theorem 6.2.1. *Let P be a parabolic subgroup of G and L be its Levi subgroup. If*

$(\mathcal{O}, \mathcal{L}) \in \mathcal{I}(\mathfrak{l}_n)^{\text{cusp}}$, then $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}}(\mathcal{E}(\mathcal{O}, \mathcal{L}))$ is parity.

Proof. By corollary 5.3.2, $\mathcal{E}(\mathcal{O}, \mathcal{L})$ exists. Let $(C, \mathcal{E}) \in \mathcal{I}(L)^{0\text{-cusp}}$ be such that, $C \cap \mathfrak{g}_n =$

\mathcal{O} and $\mathcal{E}|_{\mathcal{O}} = \mathcal{L}$. Let $y \in \mathfrak{g}_n$ and $c : G \times^P (C + \mathfrak{u}_P) \rightarrow \mathcal{N}_G$ be the map introduced in the

previous subsection. Let $Y_y = c^{-1}(y)$. Then we have an isomorphism $G/P \times (C + \mathfrak{u}_P) \rightarrow G \times^P (C + \mathfrak{u}_P)$ given by,

$$(gP, x) \mapsto (g, Ad(g^{-1})x).$$

It is easy to check that under this isomorphism the map c becomes,

$$(gP, x) \mapsto x,$$

and $Y_y = \{gP \in G/P | Ad(g^{-1})y \in \pi_P^{-1}(C)\}$. Recall that nilpotent G -orbits in \mathfrak{g} are all even dimensional. Then using the definition of induction from (??) and from the base change diagram below, and Conjecture 3.2.9, we get,

$$H_c^a(Y_y, (b^* \text{For}_P^G)^{-1} a^* \mathcal{E}[\dim C]|_{Y_y}) = 0 \text{ for } a \text{ odd.}$$

$$\begin{array}{ccc} G \times^P (C + \mathfrak{u}_P) & \xrightarrow{c} & \mathcal{N}_G \\ \uparrow & & \uparrow \\ Y_y & \xrightarrow{c} & y \end{array}$$

We define an action of \mathbb{C}^\times on Y_y by, $(t, gP) \rightarrow \chi(t)gP$. This is well defined as $y \in \mathfrak{g}_n$. Let $(Y_y)^{\mathbb{C}^\times}$ be the fixed point set. From Theorem 4.1.7,

$$H_c^a((Y_y)^{\mathbb{C}^\times}, (b^* \text{For}_P^G)^{-1} a^* \mathcal{E}[\dim C]|_{(Y_y)^{\mathbb{C}^\times}}) = 0, \text{ for } a \text{ odd.} \quad (2.2)$$

We will show that $(Y_y)^{\mathbb{C}^\times} = \sqcup_i Z^i$, where $P^i, i \in [1, b]$, is defined to be a set of representatives of G_0 -orbits of parabolic subgroups in G conjugate to P containing $\chi(\mathbb{C}^\times)$. Let L^i and U_{P^i} be the Levi and the unipotent radical of P^i respectively. An element of G conjugates P to P^i , conjugating C by the same element gives C^i contained in \mathfrak{l}^i . Let

$$Z^i = \{g(P^i)_0 \in G_0/(P^i)_0 | Ad(g^{-1})y \in (\pi^i)^{-1}(C^i)\},$$

where $\pi^i : \mathfrak{p}^i \rightarrow \mathfrak{l}^i$ and $(P^i)_0 = P^i \cap G_0$. We want to identify $g(P^i)_0 \in Z^i$ with $gg'P$ in $(Y_y)^{\mathbb{C}^\times}$, where $g' \in G$ is fixed and $g'Pg'^{-1} = P^i$. Note $g(P^i)_0 \in Z^i$, so $Ad(g^{-1})y \in (\pi^i)^{-1}(C^i)$, hence

$$Ad(gg')^{-1}y = Ad(g')^{-1}Ad(g^{-1})y \in Ad(g'^{-1})(\pi^i)^{-1}(C^i),$$

which is by definition $\pi_P^{-1}(C)$. Also,

$$(gg')^{-1}\chi(t)gg' = g'^{-1}g^{-1}\chi(t)gg' = g'^{-1}\chi(t)g',$$

and $\chi(t) \in P^i$. Therefore by definition of g' , $g'^{-1}\chi(t)g'$ belongs to P . By definition,

$$(Y_y)^{\mathbb{C}^\times} = \{gP \in G/P \mid Ad(g^{-1})y \in \pi_P^{-1}C, g^{-1}\chi(t)g \in P\}.$$

Hence $gg'P$ is in $(Y_y)^{\mathbb{C}^\times}$. Conversely, if $hP \in (Y_y)^{\mathbb{C}^\times}$, then $h^{-1}\chi(t)h \in P$. We can define $P^i = hPh^{-1}$ and $g' = h, g = e$, then by definition $gg' = h$ and $y = Ad(h)Ad(h^{-1})y \in Ad(h)\pi^{-1}C = (\pi^i)^{-1}C^i$, hence $e(P^i)_0 \in Z^i$ by identifying this with $hP \in (Y_y)^{\mathbb{C}^\times}$. In the definition of Z^i , the condition $Ad(g^{-1})y \in (\pi^i)^{-1}(C^i)$ can be redefined as below.

If $y \in \mathfrak{g}_n$ and $g \in G_0$, then this implies $Ad(g^{-1})y \in \mathfrak{g}_n$. Hence we can restate the condition $Ad(g^{-1})y \in (\pi^i)^{-1}(C^i)$ as,

$$Z^i = \{g(P^i)_0 \in G_0/(P^i)_0 \mid Ad(g^{-1})y \in \mathfrak{p}_n^i, \pi^i(Ad(g^{-1})y) \in \mathcal{O}^i\},$$

where $\mathcal{O}^i = C^i \cap \mathfrak{g}_n$. In the redefined induction diagram above, if we use the isomorphism

$$G_0/P_0 \times (\mathcal{O} + \mathfrak{u}_n) \xrightarrow{\cong} G_0 \times^{P_0} (\mathcal{O} + \mathfrak{u}_n)$$

we can see $Z^i = (p'_3)^{i-1}(y)$, where $(p'_3)^i$ is the map associated to (P^i, L^i) similar to

how we defined p'_3 . Hence from base change and the diagram below,

$$\left\{ \begin{aligned} \mathrm{H}_c^a(Z^i, (b^* \mathrm{For}_P^G)^{-1} a^* \mathcal{E}[\dim C]|_{Z^i}) &= \mathrm{H}^a(p'_3!(p'_2^* \mathrm{For}_{P_0}^{G_0})^{-1} p'_1^*(i! \mathcal{E}[\dim C])|_{\mathcal{O}})_y \\ &= \mathrm{H}^a(\mathrm{Ind}_{\mathfrak{p}^i}^{\mathfrak{g}}(\mathcal{IC}(\mathcal{O}, \mathcal{L})[\dim C - \dim \mathcal{O}]))_y \end{aligned} \right. \quad (2.3)$$

$$\begin{array}{ccccccc} & & & & Z^i = (p'_3)^{i-1}\{y\} & \longrightarrow & \{y\} \\ & & & & \downarrow & & \downarrow \\ \mathcal{O}^i & \xleftarrow{(p'_1)^i} & \mathcal{O}^i + \mathfrak{u}_n^i & \xrightarrow{(p'_2)^i} & G_0 \times^{(P^i)_0} (\mathcal{O}^i + \mathfrak{u}_n^i) & \xrightarrow{(p'_3)^i} & \mathfrak{g}_n \cap \mathcal{N}_G \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ C^i & \xrightarrow{i} & \mathfrak{l} & \xleftarrow{a'} & C^i + \mathfrak{u}_{P^i} & \xrightarrow{b^i} & G \times^{P^i} (C^i + \mathfrak{u}_{P^i}) & \xrightarrow{c^i} & \mathcal{N}_G \\ & \swarrow & & \searrow & & & & & \\ & & a^i & & & & & & \end{array}$$

we have,

$$\begin{aligned} \mathrm{H}_c^a((Y_y)^{\mathrm{C}^\times}, (b^* \mathrm{For}_P^G)^{-1} a^* \mathcal{E}[\dim C]|_{(Y_y)^{\mathrm{C}^\times}}) &= \bigoplus_i \mathrm{H}_c^a(Z^i, (b^* \mathrm{For}_P^G)^{-1} a^* \mathcal{E}[\dim C]|_{Z^i}) \\ &= \bigoplus_i \mathrm{H}^a(\mathrm{Ind}_{\mathfrak{p}^i}^{\mathfrak{g}}(\mathcal{IC}(\mathcal{O}, \mathcal{L})[\dim C - \dim \mathcal{O}]))_y. \end{aligned}$$

So we finally get,

$$\mathrm{H}_c^a((Y_y)^{\mathrm{C}^\times}, (b^* \mathrm{For}_P^G)^{-1} a^* \mathcal{E}|_{(Y_y)^{\mathrm{C}^\times}}[\dim C]) = \bigoplus_i \mathrm{H}^{a+\dim C - \dim \mathcal{O}}(\mathrm{Ind}_{\mathfrak{p}^i}^{\mathfrak{g}}(\mathcal{IC}(\mathcal{O}, \mathcal{L})))_y.$$

In the last sum one of these \mathfrak{p}^i is our original \mathfrak{p} . Hence from (2.2), we can conclude that

$$\mathrm{H}^{a+\dim C - \dim \mathcal{O}}(\mathrm{Ind}_{\mathfrak{p}}^{\mathfrak{g}}(\mathcal{IC}(\mathcal{O}, \mathcal{L})))_y = 0, \text{ for } a \text{ odd.} \quad (2.4)$$

If $\dim C - \dim \mathcal{O}$ is odd then $\mathrm{Ind}_{\mathfrak{p}}^{\mathfrak{g}} \mathcal{IC}(\mathcal{O}, \mathcal{L})$ is *-odd and if $\dim C - \dim \mathcal{O}$ is even then $\mathrm{Ind}_{\mathfrak{p}}^{\mathfrak{g}} \mathcal{IC}(\mathcal{O}, \mathcal{L})$ is *-even. As $\mathcal{IC}(\mathcal{O}, \mathcal{L}^\vee)$ is also cuspidal, so

$$\mathrm{H}^{a+\dim C - \dim \mathcal{O}}(\mathrm{Ind}_{\mathfrak{p}}^{\mathfrak{g}}(\mathcal{IC}(\mathcal{O}, \mathcal{L}^\vee)))_y = 0, \text{ for } a \text{ odd.} \quad (2.5)$$

$$\begin{aligned} \text{But, } \mathrm{Ind}_{\mathfrak{p}}^{\mathfrak{g}}(\mathcal{IC}(\mathcal{O}, \mathcal{L}^\vee)) &= \mathrm{Ind}_{\mathfrak{p}}^{\mathfrak{g}}(\mathbb{D}\mathcal{IC}(\mathcal{O}, \mathcal{L})) \\ &= \mathbb{D} \mathrm{Ind}_{\mathfrak{p}}^{\mathfrak{g}}(\mathcal{IC}(\mathcal{O}, \mathcal{L})) \text{ (by Theorem 5.1.1).} \end{aligned}$$

Therefore,

$$\begin{aligned}
\mathrm{H}^{a+\dim C-\dim \mathcal{O}}(j^! \mathrm{Ind}_{\mathfrak{p}}^{\mathfrak{g}}(\mathcal{IC}(\mathcal{O}, \mathcal{L}))) &= \mathrm{H}^{a+\dim C-\dim \mathcal{O}}(j^! \mathrm{Ind}_{\mathfrak{p}}^{\mathfrak{g}}(\mathbb{D}\mathcal{IC}(\mathcal{O}, \mathcal{L}^{\vee}))) \\
&= \mathrm{H}^{a+\dim C-\dim \mathcal{O}}(j^! \mathbb{D} \mathrm{Ind}_{\mathfrak{p}}^{\mathfrak{g}}(\mathcal{IC}(\mathcal{O}, \mathcal{L}^{\vee}))) \\
&= \mathrm{H}^{a+\dim C-\dim \mathcal{O}}(\mathrm{Ind}_{\mathfrak{p}}^{\mathfrak{g}}(\mathcal{IC}(\mathcal{O}, \mathcal{L}^{\vee})))_y,
\end{aligned}$$

where $j : \{y\} \hookrightarrow \mathfrak{g}_n \cap \mathcal{N}_G$. So by (2.5),

$$\mathrm{H}^{a+\dim C-\dim \mathcal{O}}(j^! \mathrm{Ind}_{\mathfrak{p}}^{\mathfrak{g}}(\mathcal{IC}(\mathcal{O}, \mathcal{L}^{\vee}))) = 0, \text{ for } a \text{ odd.}$$

Hence by the above fact, if $\dim C - \dim \mathcal{O}$ is odd then $\mathrm{Ind}_{\mathfrak{p}}^{\mathfrak{g}} \mathcal{IC}(\mathcal{O}, \mathcal{L})$ is !-odd and if $\dim C - \dim \mathcal{O}$ is even then $\mathrm{Ind}_{\mathfrak{p}}^{\mathfrak{g}} \mathcal{IC}(\mathcal{O}, \mathcal{L})$ is !-even, finally we can say $\mathrm{Ind}_{\mathfrak{p}}^{\mathfrak{g}} \mathcal{IC}(\mathcal{O}, \mathcal{L})$ satisfies the parity condition. □

Chapter 7. Existence Of Parity Sheaves

n -rigid

n -rigidity is introduced in [Lu]. Given a $x \in \mathfrak{g}_n$, we can always define a co-character χ and a parabolic subgroup P with a Levi subgroup L , so that (L, χ) is n -rigid, which immediately ensures that the L_0 -orbit containing x in \mathfrak{l}_n is open.

Existence

For any pair $(\mathcal{O}, \mathcal{L}) \in \mathcal{S}(\mathfrak{g}_n)$, existence of parity is still a question. Before going into that it is important to ensure the uniqueness as mentioned in [JMW]. Theorem 7.2.1 provides the essential condition needed to talk about parity sheaves in graded set-up, where n -rigidity plays an important role. Again using n -rigidity we prove Theorem 7.2.2 which is the key theorem used in proving the existence of parity for a general pair in $\mathcal{S}(\mathfrak{g}_n)$.

Modular reduction

In this section our aim is to prove an analogous Proposition like [Lu, Prop. 7.3]. But in positive characteristic the statement 2.6(c) from [Lu] does not hold. Also in characteristic 0, cuspidal pairs are clean. In positive characteristic, with the assumption on field characteristic (Assumption 3.2.4) we assume Matuner's cleanness conjecture (Conjecture 2.2.7) is true. Modular reduction provides us a triple $(\mathbb{K}, \mathbb{O}, \mathbb{k})$, with \mathbb{K} being of characteristic 0, that means whatever proved by Lustig in [Lu] holds if the sheaf coefficient is from \mathbb{K} . Modular reduction provides us a map from $K_G(\mathcal{N}_G, \mathbb{K})$ to $K_G(\mathcal{N}_G, \mathbb{k})$, where K_G denotes the Grothendieck group with objects from Perv_G . Finally at the end of this section we prove Proposition 7.3.4, which has the same version as [Lu, Prop. 7.3], but in positive

characteristic.

Normal complexes

Normal complexes were defined in [Lu] but in terms of semisimple complexes. Here we define it with parity sheaves. In Theorem 7.4.2 we prove that for any pair in $\mathcal{S}(\mathfrak{g}_n)$, the corresponding parity sheaf is a normal complex and therefore it exists. In Theorem 7.4.3, we prove that $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}}$ sends parity to parity.

7.1. n -rigidity

Let $n \in \mathbb{Z}$ be fixed. Recall the cocharacter map $\chi : \mathbb{C}^\times \rightarrow G$. Let $\phi : \mathfrak{sl}_2 \rightarrow \mathfrak{g}$ and $\tilde{\phi} : SL_2 \rightarrow G$ be such that $d\tilde{\phi} = \phi$. Define $\chi' : \mathbb{C}^\times \rightarrow G$ by,

$$\chi'(t) = \tilde{\phi} \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix}.$$

Now we define for $m \in \mathbb{Z}$,

$${}_m\mathfrak{g} = \{x \in \mathfrak{g} \mid \text{Ad}\chi'(t)x = t^m x\}.$$

Hence $\mathfrak{g} = \bigoplus_{m \in \mathbb{Z}} {}_m\mathfrak{g}$.

Definition 7.1.1. (G, χ) is said to be n -rigid if there exists ϕ such that

1. $\phi \in J_n$, which is defined in section 2.1.
2. ${}_m\mathfrak{g} = \mathfrak{g}_{nm/2}$ for $m \in \mathbb{Z}$ and $nm/2 \in \mathbb{Z}$,
3. ${}_m\mathfrak{g} = 0$ for $m \in \mathbb{Z}$ and $nm/2 \notin \mathbb{Z}$.

Proposition 7.1.2. If (G, χ) is n -rigid and $\phi(e) = x$, then

1. x is in the unique open G_0 -orbit in \mathfrak{g}_n ,
2. the map $G_0^x / (G_0^x)^\circ \rightarrow G^x / (G^x)^\circ$ is an isomorphism.

The proof of this proposition is given in [Lu, prop 4.2,5.8]. For the proof of Theorem 3.2.8 n -rigidity plays a role.

7.1.1. Construction of parabolic, nilpotent and Levi subgroups

In this section we first fix $x \in \mathfrak{g}_n$ and then we construct $\mathfrak{p}, \mathfrak{n}, \mathfrak{l}$ associated to x .

From Theorem 5.3.1, recall ϕ and the construction of χ' . Recall χ commutes with χ' and,

$${}_m\mathfrak{g} = \{g \in \mathfrak{g} | Ad(\chi'(t))g = t^m g\}.$$

Now we have the direct sum decomposition,

$$\mathfrak{g} = \bigoplus_{m, m' \in \mathbb{Z}_m} \mathfrak{g}_{m'}.$$

Here $m, m' \in \mathbb{Z}$ and ${}_m\mathfrak{g}_{m'} = {}_m\mathfrak{g} \cap \mathfrak{g}_{m'}$. We define,

$$\mathfrak{p} = \bigoplus_{m', m, 2m' / n \leq m} ({}_m\mathfrak{g}_{m'}), \mathfrak{n} = \bigoplus_{m', m, 2m' / n < m} ({}_m\mathfrak{g}_{m'}), \mathfrak{l} = \bigoplus_{m', m, 2m' / n = m} ({}_m\mathfrak{g}_{m'}).$$

Here, $\mathfrak{p}, \mathfrak{n}, \mathfrak{l}$ are parabolic, nilradical and Levi subalgebra of \mathfrak{g} [Lu, 5]. We give one example in 8.3, how to construct $\mathfrak{p}, \mathfrak{n}, \mathfrak{l}$ as defined here.

Theorem 7.1.3. *With the set-up above, $\phi(\mathfrak{sl}_2) \subset \mathfrak{l}$ and (L, χ) is n -rigid. Also x is in the open L_0 -orbit in \mathfrak{l}_n .*

7.2. Existence of parity sheaves

Let $(\mathcal{O}, \mathcal{L}) \in \mathcal{S}(\mathfrak{g}_n)$. Let $x \in \mathcal{O}$ and $\mathfrak{p}, \mathfrak{n}, \mathfrak{l}$ be Lie subalgebras of \mathfrak{g} constructed as above connected with x . Let P, U, L be the subgroups of G with Lie Algebras $\mathfrak{p}, \mathfrak{n}, \mathfrak{l}$ respectively. It follows that P contains the image $\chi(\mathbb{C}^\times)$. By Theorem 7.1.3, x is contained in an open L_0 -orbit in \mathfrak{l}_n , call it \mathcal{O}_L . As $x \in \mathcal{O}_L$, $\mathcal{O}_L \subset \mathcal{O}$. Now we can restrict \mathcal{L} to \mathcal{O}_L . By [Lu, prop 5.8], the inclusion induces isomorphisms on $G_0^x / (G_0^x)^\circ$ and $L_0^x / (L_0^x)^\circ$. Hence,

$$\text{Loc}_{f, G_0}(\mathcal{O}, \mathbb{k}) = \text{Loc}_{f, L_0}(\mathcal{O}_L, \mathbb{k}).$$

So $\mathcal{L}|_{\mathcal{O}_L}$ is a local system on \mathfrak{l}_n , let's call it \mathcal{L}' . Now we are ready to prove the parity vanishing theorem mentioned in section 3.1.

Theorem 7.2.1. *Let \mathcal{O} be a nilpotent orbit in \mathfrak{g}_n and $\mathcal{L} \in \text{Loc}_{f,G_0}(\mathcal{O}, \mathbb{k})$, then $H_{G_0}^*(\mathcal{L})$ vanishes in odd degrees.*

Proof. Define $\tilde{\mathcal{O}} := G_0/(G_0^x)^\circ$ and $\pi : \tilde{\mathcal{O}} \rightarrow \mathcal{O}$ by $g(G_0^x)^\circ \rightarrow g.x$. This is a Galois covering map with the Galois group $A_{G_0}(x)$. Also we know that

$$\text{Loc}_{f,G_0}(\mathcal{O}, \mathbb{k}) \cong \mathbb{k}[A_{G_0}(x)] - \text{mod}.$$

We can construct the Levi subgroup L as defined above. Then by Theorem 7.1.3, (L, χ) is n -rigid. Hence by Proposition 7.1.2, $L^x/(L^x)^\circ \cong L_0^x/(L_0^x)^\circ$. But in the above paragraph we mentioned $G_0^x/(G_0^x)^\circ \cong L_0^x/(L_0^x)^\circ$. By Remark 3.2.6, $|A_L(x)|$ is invertible in \mathbb{k} . Hence by the above isomorphisms, $|A_{G_0}(x)|$ is invertible in \mathbb{k} . Therefore any $\mathbb{k}[A_{G_0}(x)]$ -module is a summand of the direct sum of copies of the regular representation, which again corresponds to $\pi_*\mathbb{k}_{\tilde{\mathcal{O}}}$.

$$\begin{aligned} H_{G_0}^*(\mathcal{O}, \pi_*\mathbb{k}_{\tilde{\mathcal{O}}}) &\cong H_{G_0}^*(\tilde{\mathcal{O}}) \\ &\cong H_{(G_0^x)^\circ}^*(pt) \text{ (by quotient equivalence)} \\ &\cong H_{(G_0^x)^\circ - red}^*(pt). \end{aligned}$$

Here $(G_0^x)^\circ - red$ is the reductive quotient of $(G_0^x)^\circ$. By Lemma 3.2.3 and assumption 3.2.4, l is not a torsion prime for $(G^x)^\circ - red$. By [JMW, Theorem 2.44], $H_{(G_0^x)^\circ - red}^*(pt)$ will vanish in odd degrees if we can show that l is not a torsion prime for $(G_0^x)^\circ - red$. This will follow from showing $(G_0^x)^\circ - red$ is a regular subgroup of $(G^x)^\circ - red$. Define a map $\psi : \mathbb{C}^\times \mapsto \mathbb{C}^\times \times G$ by,

$$t \mapsto (t^n, \chi(t)).$$

Then $\mathbb{C}^\times \times G_0$ is the centralizer of $\psi(\mathbb{C}^\times)$. So $(\mathbb{C}^\times \times G_0)^x = (\mathbb{C}^\times \times G)^x \cap C_{\psi(\mathbb{C}^\times)}, C_{\psi(\mathbb{C}^\times)}$

is the centralizer of $\psi(\mathbb{C}^\times)$. Therefore any maximal torus in $(\mathbb{C}^\times \times G)^x$ containing $\psi(\mathbb{C}^\times)$ commutes with $\psi(\mathbb{C}^\times)$, hence in $(\mathbb{C}^\times \times G_0)^x$. So $(\mathbb{C}^\times \times G_0)^x$ is regular subgroup of $(\mathbb{C}^\times \times G)^x$. Let us define $\mathbb{C}^\times \ltimes G^x$ by the action of \mathbb{C}^\times on G^x as $(t, g) \rightarrow \chi(t)g\chi(t^{-1})$. Now we define a map $(\mathbb{C}^\times \times G)^x \rightarrow \mathbb{C}^\times \ltimes G^x$ by,

$$(t, g) \rightarrow [(t, \chi(t)g\chi(t^{-1}))].$$

It is easy to check this is an isomorphism and image of $\psi(\mathbb{C}^\times)$ under this isomorphism is contained in $1 \times G^x \cong G^x$. Similarly we have another isomorphism $(\mathbb{C}^\times \times G_0)^x \cong \mathbb{C}^\times \ltimes G_0^x$. Therefore from the previous deduction we can say any maximal torus in $G^x \subset 1 \times G^x \subset \mathbb{C}^\times \ltimes G^x$ containing $\psi(\mathbb{C}^\times)$ that commute with $\psi(\mathbb{C}^\times)$ will be contained in $\mathbb{C}^\times \ltimes G_0^x$, so in $1 \times G_0^x \cong G_0^x$. Now we can conclude from the previous deduction that G_0^x is regular subgroup of G^x . Hence $(G_0^x)^\circ - red$ is regular subgroup of $(G^x)^\circ - red$ and we are done. \square

Theorem 7.2.2. *Let $(\mathcal{O}, \mathcal{L}) \in \mathcal{I}(\mathfrak{g}_n)$ and \mathfrak{l}_n and \mathcal{L}' constructed above. Assume that $\mathcal{E}(\mathcal{O}_L, \mathcal{L}')$ exists, then*

- (a) *The support of $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} \mathcal{E}(\mathcal{O}_L, \mathcal{L}')$ is $\bar{\mathcal{O}}$, and*
- (b) *$\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} \mathcal{E}(\mathcal{O}_L, \mathcal{L}')|_{\mathcal{O}} = \mathcal{L}[\dim \mathcal{O}_L]$.*

Proof.

(a) Let $y \in \mathfrak{g}_n$ be in the support of $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} \mathcal{E}(\mathcal{O}_L, \mathcal{L}')$. We need to show that $y \in \bar{\mathcal{O}}$. From the definition of induction, there exists $\eta \in \mathfrak{p}_n$ and $g \in G_0$, such that, $Ad(g)\eta = y$. Now both the support of $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} \mathcal{E}(\mathcal{O}_L, \mathcal{L}')$ and $\bar{\mathcal{O}}$ are G_0 -invariant. Hence we can replace y by $\eta \in \mathfrak{p}_n$. By [Lu, 5.9], \mathfrak{p}_n coincides with the closure of the P_0 -orbit of x in \mathfrak{p}_n which is again contained in $\bar{\mathcal{O}}$. Hence $y \in \bar{\mathcal{O}}$ and part (a) is proved.

(b) Recall the induction diagram

$$\mathfrak{l}_n \xleftarrow{\pi} \mathfrak{p}_n \xrightarrow{e} G_0 \times^{P_0} \mathfrak{p}_n \xrightarrow{\mu} \mathfrak{g}_n.$$

Let $E_{\mathcal{O}} = \mu^{-1}(\mathcal{O})$. We first show that μ is an isomorphism when restricted to $E_{\mathcal{O}}$. Actions of G_0 on $E_{\mathcal{O}}$ and \mathcal{O} are compatible with the map μ . Also action of G_0 on \mathcal{O} is transitive. So to prove that μ is a bijection, it is enough to show that $\mu^{-1}(x)$ is a single point. Let $(g, \gamma) \in G_0 \times \mathfrak{p}_n$ be in the inverse image. So $Ad(g)\gamma = x$. Therefore $x \in Ad(g)\mathfrak{p}$ and by [Lu, 5.7], $Ad(g)\mathfrak{p} = \mathfrak{p}$. Hence $g \in P_0$. Hence $(g, \gamma) = (1, Ad(g)\gamma) = (1, x)$. Hence $\mu^{-1}(x)$ is a singleton and μ is a bijection of smooth varieties, thus isomorphism on $E_{\mathcal{O}}$.

Let

$$\begin{aligned} \mathcal{G} &= \text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} \mathcal{E}(\mathcal{O}_L, \mathcal{L}')|_{\mathcal{O}} = (\mu)_!(e^* \text{For}_{P_0}^{G_0})^{-1} \pi^*(\mathcal{E}(\mathcal{O}_L, \mathcal{L}'))|_{\mathcal{O}} \\ &= (\mu|_{E_{\mathcal{O}}})_!(e^* \text{For}_{P_0}^{G_0})^{-1} \pi^*(\mathcal{E}(\mathcal{O}_L, \mathcal{L}'))|_{E_{\mathcal{O}}}. \end{aligned}$$

As $\mu|_{E_{\mathcal{O}}}$ is an isomorphism, hence $(\mu|_{E_{\mathcal{O}}})_!$ is an equivalence of categories. So in other words, \mathcal{G} satisfies,

$$(\mu|_{E_{\mathcal{O}}})^*(\mathcal{G}) = (e^* \text{For}_{P_0}^{G_0})^{-1} \pi^*(\mathcal{E}(\mathcal{O}_L, \mathcal{L}'))|_{E_{\mathcal{O}}}$$

In fact $(\mu|_{E_{\mathcal{O}}})^*(\mathcal{G})$ is uniquely determined by,

$$((e|_{E'_{\mathcal{O}}})^* \text{For}_{P_0}^{G_0})(\mu|_{E_{\mathcal{O}}})^*(\mathcal{G}) = (\pi|_{E'_{\mathcal{O}}})^*(\mathcal{E}(\mathcal{O}_L, \mathcal{L}')|_{\mathcal{O}_L}) = (\pi|_{E'_{\mathcal{O}}})^* \mathcal{L}'[\dim \mathcal{O}_L], \quad (2.1)$$

where $E'_{\mathcal{O}} = e^{-1}(E_{\mathcal{O}})$.

$$\begin{array}{ccccc} \bar{\mathcal{O}}_L & \xleftarrow{\pi} & \bar{\mathcal{O}}_L + \mathfrak{u}_n & \xrightarrow{e} & G_0 \times^{P_0} (\bar{\mathcal{O}}_L + \mathfrak{u}_n) & \xrightarrow{\mu} & \bar{\mathcal{O}} \\ \uparrow & & \uparrow & & \uparrow & & \\ \mathcal{O}_L & \xleftarrow{\pi} & \mathcal{O}_L + \mathfrak{u}_n & \xrightarrow{e} & G_0 \times^{P_0} (\mathcal{O}_L + \mathfrak{u}_n) & & \end{array}$$

As we already proved $E_{\mathcal{O}} \cong^{\mu} \mathcal{O}$ and μ is G_0 equivariant, $E_{\mathcal{O}}$ is a single orbit. $G_0 \times^{P_0} (\mathcal{O}_L + \mathfrak{u}_n)$ is stable under G_0 action, therefore $E_{\mathcal{O}} \subset G_0 \times^{P_0} (\mathcal{O}_L + \mathfrak{u}_n)$ and it follows that $E'_{\mathcal{O}} \subset \mathcal{O}_L + \mathfrak{u}_n$. Now we have the diagram below,

$$\begin{array}{ccccc}
 \mathcal{O}_L & \longleftarrow & E'_{\mathcal{O}} & \xrightarrow{e|_{E'_{\mathcal{O}}}} & E_{\mathcal{O}} & \xrightarrow{\mu|_{E_{\mathcal{O}}}} & \mathcal{O} \\
 \uparrow id & & \nearrow j & & & & \nearrow i \\
 \mathcal{O}_L & & & & & &
 \end{array}$$

This diagram is commutative.

$$\begin{aligned}
 ((e|_{E'_{\mathcal{O}}})^* \text{For}_{P_0}^{G_0})(\mu|_{E_{\mathcal{O}}})^* \mathcal{L}[\dim \mathcal{O}_L] &= j_* i^* \mathcal{L}[\dim \mathcal{O}_L] \\
 &= (\pi|_{E'_{\mathcal{O}}})^* (\mathcal{L}[\dim \mathcal{O}_L]|_{\mathcal{O}_L}) \\
 &= (\pi|_{E'_{\mathcal{O}}})^* (\mathcal{L}'[\dim \mathcal{O}_L]) \\
 &= \pi^* \mathcal{E}(\mathcal{O}_L, \mathcal{L}')|_{E'_{\mathcal{O}}}.
 \end{aligned}$$

We already know $e|_{E'_{\mathcal{O}}}$ is closed embedding and $\mu|_{E_{\mathcal{O}}}$ is smooth, so the pull back of these maps induce faithful functor on the local systems. Hence from the above equation and (2.1), we have $\mathcal{G} = \mathcal{L}[\dim \mathcal{O}_L]$.

□

7.3. Modular reduction

Recall from the background, we assumed that there exists a finite extension \mathbb{K} of \mathbb{Q}_l with ring of integers \mathbb{O} and residue field \mathbb{k} . Also assume for each $x \in \mathcal{N}_G$, all the irreducible representations of $A_G(x)$ are defined over \mathbb{K} . Let $K_G(\mathcal{N}_G, \mathbb{k})$ denote the Grothendieck group generated by the isomorphism classes of simple objects in

$\text{Perv}_G(\mathcal{N}_G, \mathbb{k})$. Similarly define $K_G(\mathcal{N}_G, \mathbb{K})$. By [J2, 2.9], there exists a \mathbb{Z} -linear map,

$$d : K_G(\mathcal{N}_G, \mathbb{K}) \rightarrow K_G(\mathcal{N}_G, \mathbb{k}).$$

This map is called modular reduction map and is defined in the following way: If $\mathcal{F} \in \text{Perv}_G(\mathcal{N}_G, \mathbb{K})$ and $\mathcal{F}_\mathbb{O}$, a torsion free object in $\text{Perv}_G(\mathcal{N}_G, \mathbb{O})$ such that $\mathcal{F} \cong \mathbb{K} \otimes_\mathbb{O} \mathcal{F}_\mathbb{O}$, then

$$d([\mathcal{F}]) = [\mathbb{k} \otimes_\mathbb{O}^L \mathcal{F}_\mathbb{O}].$$

If $\mathcal{F} = \mathcal{IC}(C, \mathcal{E})$, then there exists a G -equivariant local system $\mathcal{E}_\mathbb{O}$ on C such that $\mathcal{E} = \mathbb{K} \otimes_\mathbb{O} \mathcal{E}_\mathbb{O}$, and

$$d[\mathcal{IC}(C, \mathcal{E})] = [\mathbb{k} \otimes_\mathbb{O}^L \mathcal{IC}(C, \mathcal{E}_\mathbb{O})].$$

By [JMW, prop 2.39], $\mathbb{k} \otimes_\mathbb{O}^L \mathcal{E}(C, \mathcal{E}_\mathbb{O}) \cong \mathcal{E}(C, \mathbb{k} \otimes_\mathbb{O}^L \mathcal{E}_\mathbb{O})$. We can call $\mathbb{k} \otimes_\mathbb{O}^L \mathcal{E}_\mathbb{O}$ to be $\mathcal{E}_\mathbb{k}$.

Theorem 7.3.1. *The modular reduction above gives a well-defined map with the following properties:*

1. $\text{Irr}(\mathbb{K}[G^x/(G^x)^\circ] - \text{mod}) \xrightarrow{\cong} \text{Irr}(\mathbb{k}[G^x/(G^x)^\circ] - \text{mod})$.
2. *If M is a torsion free module in $\mathbb{O}[G^x/(G^x)^\circ]$, then the direct summands of $\mathbb{K} \otimes_\mathbb{O} M$ are in bijection with the direct summands of $\mathbb{k} \otimes_\mathbb{O} M$.*

The proof follows from [CR, Theorem 82.1].

Let P be a parabolic subgroup with L as the Levi factor.

Theorem 7.3.2. *If $(C, \mathcal{F}) \in \mathcal{S}(L)^{0-\text{cusp}}$ and $(C, \mathcal{G}) \in \mathcal{S}(L, \mathbb{K})^{0-\text{cusp}}$, whose modular reduction is \mathcal{F} , then,*

1. $\mathcal{IC}(C, \mathcal{G}_\mathbb{O})$ is clean, and
2. $\text{Ind}_P^G \mathcal{IC}(C, \mathcal{G}_\mathbb{O})$ is parity.

Proof. 1. If there exists $y \in \bar{C} - C$ such that $\mathcal{IC}(C, \mathcal{G}_\mathbb{O})_y \neq 0$, then it must have torsion part only; otherwise, $\mathbb{K} \otimes^L \mathcal{IC}(C, \mathcal{G}_\mathbb{O})_y \neq 0$ but is same as $\mathcal{IC}(C, \mathcal{G})_y$ which is zero by Conjecture 2.2.7. Also by [J2, 2.6], $\mathbb{k} \otimes_\mathbb{O}^L \mathcal{IC}(C, \mathcal{G}_\mathbb{O})$ is a perverse sheaf with $\mathbb{k} \otimes_\mathbb{O}^L \mathcal{IC}(C, \mathcal{G}_\mathbb{O})|_C = \mathbb{k} \otimes_\mathbb{O}^L \mathcal{G}_\mathbb{O}[\dim C]$. We have the open and closed embeddings below,

$$C \xrightarrow{j} \bar{C} \xleftarrow{i} \bar{C} - C ,$$

which gives rise to the distinguished triangle,

$$j_! \mathbb{k} \otimes_{\mathbb{O}}^L \mathcal{G}_{\mathbb{O}}[\dim C] \rightarrow \mathbb{k} \otimes_{\mathbb{O}}^L \mathcal{IC}(C, \mathcal{G}_{\mathbb{O}}) \rightarrow i_* i^*(\mathbb{k} \otimes_{\mathbb{O}}^L \mathcal{IC}(C, \mathcal{G}_{\mathbb{O}})) \rightarrow .$$

By conjecture 2.2.7, $j_!(\mathbb{k} \otimes_{\mathbb{O}}^L \mathcal{G}_{\mathbb{O}}[\dim C]) = \mathcal{IC}(C, \mathbb{k} \otimes_{\mathbb{O}}^L \mathcal{G}_{\mathbb{O}})$. The third morphism in the above distinguished triangle is 0, as $i^!(\mathcal{IC}(C, \mathbb{k} \otimes_{\mathbb{O}}^L \mathcal{G}_{\mathbb{O}})) = 0$ by conjecture 2.2.7. Therefore this distinguished triangle splits and we have,

$$\mathbb{k} \otimes_{\mathbb{O}}^L \mathcal{IC}(C, \mathcal{G}_{\mathbb{O}}) = \mathcal{IC}(C, \mathbb{k} \otimes_{\mathbb{O}}^L \mathcal{G}_{\mathbb{O}}) \oplus i_* i^*(\mathbb{k} \otimes_{\mathbb{O}}^L \mathcal{IC}(C, \mathcal{G}_{\mathbb{O}})).$$

As $\mathbb{k} \otimes_{\mathbb{O}}^L \mathcal{IC}(C, \mathcal{G}_{\mathbb{O}})$ and $\mathcal{IC}(C, \mathbb{k} \otimes_{\mathbb{O}}^L \mathcal{G}_{\mathbb{O}})$ are perverse, so $i_* i^*(\mathbb{k} \otimes_{\mathbb{O}}^L \mathcal{IC}(C, \mathcal{G}_{\mathbb{O}}))$ must be perverse. Now it will not be hard to check that for a torsion \mathbb{O} -module M , $H^i(\mathbb{k} \otimes_{\mathbb{O}}^L M)$ is nonzero for $i = 0, -1$. So if we choose an open orbit \mathcal{O}' in the support of $i_* i^*(\mathbb{k} \otimes_{\mathbb{O}}^L \mathcal{IC}(C, \mathcal{G}_{\mathbb{O}}))$ such that y is in that orbit then by the above statement we will have $H^i(\mathbb{k} \otimes_{\mathbb{O}}^L \mathcal{IC}(C, \mathcal{G}_{\mathbb{O}})_y) \neq 0$ for $i = -\dim \mathcal{O}', -\dim \mathcal{O}' - 1$, which contradicts the perversity of $i_* i^*(\mathbb{k} \otimes_{\mathbb{O}}^L \mathcal{IC}(C, \mathcal{G}_{\mathbb{O}}))$.

2. Note that $\mathbb{K} \otimes_{\mathbb{O}}^L \text{Ind}_P^G \mathcal{IC}(C, \mathcal{G}_{\mathbb{O}}) = \text{Ind}_P^G \mathcal{IC}(C, \mathcal{G})$ which is parity because it is in characteristic 0. Also, $\mathbb{k} \otimes_{\mathbb{O}}^L \text{Ind}_P^G \mathcal{IC}(C, \mathcal{G}_{\mathbb{O}}) \cong \text{Ind}_P^G \mathcal{E}(C, \mathcal{F})$ is parity by Conjecture 3.2.9. Combining these two facts and using [JMW, Prop. 2.37], $\text{Ind}_P^G \mathcal{IC}(C, \mathcal{G}_{\mathbb{O}})$ is parity. □

Theorem 7.3.3. *For $(C, \mathcal{F}) \in \mathcal{S}(G)$, there exists a Levi subgroup L and a pair $(C', \mathcal{F}') \in \mathcal{S}(L)^{0\text{-cusp}}$ such that, $H^{-\dim C}(\text{Ind}_P^G \mathcal{IC}(C', \mathcal{F}'))|_C$ contains \mathcal{F} as a direct summand.*

Proof. By [AJHR], if l is rather good, which is our assumption here, we have a bijection,

$$K_G(\mathcal{N}_G, \mathbb{K}) \xrightarrow{\cong} K_G(\mathcal{N}_G, \mathbb{k}).$$

Let $(C, \mathcal{G}) \in \mathcal{S}(G, \mathbb{K})$ be the pair whose modular reduction is \mathcal{F} . By [Lu3], there exists a parabolic subgroup P with Levi L and $(C', \mathcal{G}') \in \mathcal{S}(L, \mathbb{K})^{0\text{-cusp}}$ be such that $\mathcal{IC}(C, \mathcal{G})$ is a direct summand of $\text{Ind}_P^G \mathcal{IC}(C', \mathcal{G}')$.

$$\mathbb{K} \otimes^L H^{-\dim C}(\text{Ind}_P^G \mathcal{IC}(C', \mathcal{G}'_{\mathbb{O}}))|_C \cong H^{-\dim C}(\text{Ind}_P^G \mathcal{IC}(C', \mathcal{G}'))|_C$$

which contains \mathcal{G} . Now, $H^{-\dim C}(\text{Ind}_P^G \mathcal{IC}(C', \mathcal{G}'_{\mathbb{O}}))|_C$ is torsion-free. If not, then $\mathbb{k} \otimes_{\mathbb{O}}^L$

$H^{-\dim C}(\text{Ind}_P^G \mathcal{IC}(C', \mathcal{G}'_{\mathbb{O}}))|_C$ has cohomology concentrated in two consecutive de-

grees, which contradicts Theorem 7.3.2(2). By Theorem 7.3.1, direct summands

appearing in $H^{-\dim C}(\text{Ind}_P^G \mathcal{IC}(C', \mathcal{G}'))|_C$ are in bijection with direct summands appearing in $H^{-\dim C}(\text{Ind}_P^G \mathcal{IC}(C', \mathcal{G}'_k))|_C$. Set \mathcal{F}' to be \mathcal{G}'_k . Hence $\mathcal{F}' \in \mathcal{S}(L)^{0\text{-cusp}}$ and $H^{-\dim C}(\text{Ind}_P^G \mathcal{IC}(C', \mathcal{F}'))|_C$ contains \mathcal{F} as a direct summand.

□

Proposition 7.3.4. *Let $(\mathcal{O}, \mathcal{L}) \in \mathcal{S}(\mathfrak{g}_n)$. There exists an integer b and $(\mathcal{O}', \mathcal{L}') \in \mathcal{S}(\mathfrak{g}_n)^{\text{cusp}}$ for some parabolic subgroup P with the Levi subgroup L , such that \mathcal{L} is a direct summand of $H^{b-\dim \mathcal{O}'}(\text{Ind}_P^{\mathfrak{g}} \mathcal{IC}(\mathcal{O}', \mathcal{L}'))|_{\mathcal{O}}$.*

Proof. Let $(\mathcal{O}, \mathcal{L}) \in \mathcal{S}(\mathfrak{g}_n)$ and $(C, \mathcal{E}) \in \mathcal{S}(G)$ such that $C \cap \mathfrak{g}_n = \mathcal{O}$ and $\mathcal{E}|_{\mathcal{O}} = \mathcal{L}$. By Theorem 7.3.3, there exists Q , a parabolic subgroup containing $\chi(\mathbb{C}^\times)$ with M , the Levi subgroup such that $(C', \mathcal{E}') \in \mathcal{S}(M)^{0\text{-cusp}}$ and $H^{-\dim C}(\text{Ind}_Q^G \mathcal{IC}(C', \mathcal{E}'))|_C$ contains \mathcal{E} as direct summand.

For this we will imitate the proof of Theorem 6.2.1. Let $y \in \mathcal{O}$, we construct the parabolic induction diagram corresponding to (Q, M, C') . That is,

$$C' \xleftarrow{a} C' + \mathfrak{u}_Q \xrightarrow{b} G \times^Q (C' + \mathfrak{u}_Q) \xrightarrow{c} \mathcal{N}_G$$

Let $Y_y = c^{-1}(y)$ and recall that,

$$H_c^a(Y_y, (b^* \text{For}_Q^G)^{-1} a^* \mathcal{E}'[\dim C']|_{Y_y}) = 0 \text{ for } a \text{ odd.}$$

Also recall the action of \mathbb{C}^\times on Y_y defined in the proof of Theorem 6.2.1. Let again $Y_y^{\mathbb{C}^\times}$ be the fixed point set of this action. It is not hard to see that the stabilizer of each point in the complement of $(Y_y)^{\mathbb{C}^\times}$ is trivial. Hence by The Lemma 4.1.3, $\dim H_{\mathbb{C}^\times}^*(Y_y - (Y_y)^{\mathbb{C}^\times}) < \infty$. Now using Lemma 4.1.8, the Euler characteristic of $H_c^*(Y_y - (Y_y)^{\mathbb{C}^\times})$ is 0, so we have,

$$\begin{aligned} & \sum_a (-1)^a H_c^a(Y_y, (b^* \text{For}_Q^G)^{-1} a^* \mathcal{E}'[\dim C']|_{Y_y}) \\ &= \sum_a (-1)^a H_c^a((Y_y)^{\mathbb{C}^\times}, (b^* \text{For}_Q^G)^{-1} a^* \mathcal{E}'[\dim C']|_{(Y_y)^{\mathbb{C}^\times}}) \end{aligned} \quad (3.2)$$

As the stabilizer of each point in $Y_y - (Y_y)^{\mathbb{C}^\times}$ is trivial, so from Theorem 4.1.7,

$$H_c^a(Y_y^{\mathbb{C}^\times}, (b^* \text{For}_Q^G)^{-1} a^* \mathcal{E}'[\dim C']|_{(Y_y)^{\mathbb{C}^\times}}) = 0 \text{ for } a \text{ odd.}$$

Combining both the result we have,

$$\sum_{a \text{ even}} H_c^a(Y_y, (b^* \text{For}_Q^G)^{-1} a^* \mathcal{E}'[\dim C']|_{Y_y}) = \sum_{a \text{ even}} H_c^a(Y_y)^{\mathbb{C}^\times}, (b^* \text{For}_Q^G)^{-1} a^* \mathcal{E}'[\dim C']|_{(Y_y)^{\mathbb{C}^\times}} \quad (3.3)$$

Now let Q^i 's denote the G_0 -orbits of Q in the set of all parabolic subgroups containing $\chi(\mathbb{C}^\times)$, for $i = 1, \dots, b$. Define Z^i as before but in terms of Q^i and C'^i , where an element of G conjugates Q to Q^i , conjugating C' by the same element gives C'^i contained in \mathfrak{m}^i .

$$Z^i = \{g(Q^i)_0 \in G_0/(Q^i)_0 \mid \text{Ad}(g^{-1})y \in (\pi^i)^{-1}(C'^i)^i\}.$$

Then as before we get,

$$Y_y^{\mathbb{C}^\times} = \sqcup_i Z^i,$$

and,

$$H_c^a(Z^i, (b^* \text{For}_Q^G)^{-1} a^* \mathcal{E}'[\dim C']|_{Z^i}) = H^a(\text{Ind}_{\mathfrak{q}^i}^{\mathfrak{g}}(\mathcal{IC}(\mathcal{O}^i, \mathcal{L}^i)[\dim C' - \dim \mathcal{O}^i])_y), \quad (3.4)$$

where $(\mathcal{O}^i, \mathcal{L}^i) \in \mathcal{S}(M^i)$. These come from conjugating and restricting (C', \mathcal{E}') . Now combining equation 3.3 and the above result we have,

$$\sum_{a \text{ even}} H_c^a(Y_y, (b^* \text{For}_Q^G)^{-1} a^* \mathcal{E}'[\dim C']|_{Y_y}) = \sum_{i, a \text{ even}} H^a(\text{Ind}_{\mathfrak{q}^i}^{\mathfrak{g}}(\mathcal{IC}(\mathcal{O}^i, \mathcal{L}^i)[\dim C' - \dim \mathcal{O}^i])_y).$$

Now $H_c^a(Y_y, (b^* \text{For}_Q^G)^{-1} a^* \mathcal{E}'[\dim C']|_{Y_y})$ is the same as $H^a(\text{Ind}_Q^G \mathcal{IC}(C', \mathcal{E}'))_y$. By assumption \mathcal{E} occurs as direct summand of $H^{-\dim C}(\text{Ind}_Q^G \mathcal{IC}(C', \mathcal{E}'))|_C$. As $\mathcal{L} = \mathcal{E}|_{\mathcal{O}}$, therefore for some i and a even, \mathcal{L} should appear as a direct summand of $H^a(\text{Ind}_{\mathfrak{q}^i}^{\mathfrak{g}}(\mathcal{IC}(\mathcal{O}^i, \mathcal{L}^i)[\dim C' - \dim \mathcal{O}^i])_y$. We call this Q^i to be P , M^i to be L and $(\mathcal{O}^i, \mathcal{L}^i)$ to be $(\mathcal{O}', \mathcal{L}')$. Hence we get the desired result that \mathcal{L} is a direct summand of $H^{a+\dim C' - \dim \mathcal{O}'}(\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} \mathcal{IC}(\mathcal{O}', \mathcal{L}'))$. we can call $a + \dim C'$ to be b . \square

7.4. Normal complexes

Definition 7.4.1. An $A \in D_{G_0}^b(\mathfrak{g}_n)$ is called normal if there exists $(\mathcal{O}, \mathcal{L}) \in \mathcal{S}(\mathfrak{l}_n)^{\text{cusp}}$ such that some shift of A is a direct summand of $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}}(\mathcal{E}(\mathcal{O}, \mathcal{L}))$.

Theorem 7.4.2. For $(\mathcal{O}, \mathcal{L}) \in \mathcal{S}(\mathfrak{g}_n)$, $\mathcal{E}(\mathcal{O}, \mathcal{L})$ exists and is a normal complex.

Proof. For $(\mathcal{O}, \mathcal{L}) \in \mathcal{S}(\mathfrak{g}_n)$, we can construct the Levi subgroup L as in 7.1.1 and $(\mathcal{O}_L, \mathcal{L}') \in \mathcal{S}(\mathfrak{l}_n)$ as in 7.2 and by Theorem 7.2.2, $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} \mathcal{E}(\mathcal{O}_L, \mathcal{L}')|_{\mathcal{O}} = \mathcal{L}[\dim \mathcal{O}_L]$. Now by Proposition 7.3.4, there exists a parabolic subgroup $Q \subset L$ and a Levi subgroup $M \subset Q$ with $(\mathcal{O}', \mathcal{L}'') \in \mathcal{S}(\mathfrak{m}_n)^{0\text{-cusp}}$ such that $H^{b-\dim \mathcal{O}'}(\text{Ind}_{\mathfrak{q}}^{\mathfrak{l}} \mathcal{IC}(\mathcal{O}', \mathcal{L}'')|_{\mathcal{O}_L})$ contains \mathcal{L}' as a direct summand. By Theorem 6.2.1, $\text{Ind}_{\mathfrak{q}}^{\mathfrak{l}} \mathcal{IC}(\mathcal{O}', \mathcal{L}'')$ is parity. By proposition 7.1.2, \mathcal{O}_L is open in \mathfrak{l}_n . Combining these two above facts, we can see that $\mathcal{E}(\mathcal{O}_L, \mathcal{L}')$ exists and is direct summand of $\text{Ind}_{\mathfrak{q}}^{\mathfrak{l}} \mathcal{IC}(\mathcal{O}', \mathcal{L}'')$. Using the fact that induction is transitive it follows, $\mathcal{L}[\dim \mathcal{O}_L]$ is direct summand of $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} \mathcal{IC}(\mathcal{O}', \mathcal{L}'')|_{\mathcal{O}}$. By Theorem 7.2.2, support of $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} \mathcal{E}(\mathcal{O}_L, \mathcal{L}')$ is $\bar{\mathcal{O}}$. Therefore $\mathcal{E}(\mathcal{O}, \mathcal{L})$ exists and is direct summand of $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} \mathcal{E}(\mathcal{O}', \mathcal{L}'')$. \square

7.4.1. Induction preserves parity

Theorem 7.4.3. Let P be a parabolic subgroup of G with a Levi factor L . For any pair $(\mathcal{O}, \mathcal{L}) \in \mathcal{S}(\mathfrak{l}_n)$, the induction functor sends parity complexes to parity complexes.

Proof. By Theorem 7.4.2, there exist a cuspidal pair $(\mathcal{C}, \mathcal{F}) \in \mathcal{I}(\mathfrak{m}_n)^{0\text{-cusp}}$, where M is the Levi subgroup of L such that

$$\text{Ind}_{\Gamma \cap \mathfrak{p}}^l(\mathcal{E}(\mathcal{C}, \mathcal{F})) = \mathcal{E}(\mathcal{O}, \mathcal{L})[k] \oplus \dots, \text{ for some } k \in \mathbb{Z}.$$

If we apply $\text{Ind}_{\mathfrak{p}}^g$ on both sides and use the transitivity of induction, then we get,

$$\text{Ind}_{\mathfrak{p}}^g(\mathcal{E}(\mathcal{C}, \mathcal{F})) = \text{Ind}_{\mathfrak{p}}^g(\mathcal{E}(\mathcal{O}, \mathcal{L})) [k] \oplus \dots$$

Now the left-hand side is parity by Theorem 6.2.1. Hence so is the right-hand side. So induction preserves the parity of $\mathcal{E}(\mathcal{O}, \mathcal{L})$. □

Chapter 8. Examples

Several cases for conjecture 3.2.9

Recall Conjecture 3.2.9, which plays an analogous role as the statement 2.6(c) from [Lu]. The result is still unknown in positive characteristic. Our next aim will be to prove this conjecture in general. Here we give some specific examples where the conjecture is true. More specifically, we calculate Ind_P^G for some classical algebraic groups (in 8.1 for \mathfrak{sp}_4 and in 8.2 for \mathfrak{sl}_4) for cuspidal pairs on the Levis and show that Ind_P^G applied to the cuspidal pair has vanishing stalks in odd degrees.

For $x \in \mathfrak{g}_n$, construction of parabolic with Levi as described in 7.1.1

Recall from subsection 7.1.1, that for $x \in \mathfrak{g}_n$, we first defined χ and then constructed $\mathfrak{p}, \mathfrak{l}$. The Levi L with χ was n -rigid, which played a key role in the proofs of section 7.1. In section 8.3, we construct $\mathfrak{p}, \mathfrak{l}$ for x being the representatives of various G_0 orbits in \mathfrak{g}_{-1} .

8.1. \mathfrak{sp}_4 -case:

Let $G = Sp_4$. The symplectic form is defined by the matrix

$$B = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}$$

and the inner product is $Q(v, w) = v^t B w$, where $v, w \in \mathbb{C}^4$. Sp_4 is defined as the group of automorphisms A from \mathbb{C}^4 to \mathbb{C}^4 , such that $Q(Av, Aw) = Q(v, w)$. Here a torus is of the form $\text{diag}(t_1, t_2, t_1^{-1}, t_2^{-2})$. Let $\{e_1, e_2, e_3, e_4\}$ be the standard basis of \mathbb{C}^4 . Hence the root system is $\Phi = \{\pm e_1 \pm e_2, \pm 2e_1, \pm 2e_2\}$. The orthogonal complement of a vector space V ,

denoted by V^\perp , is the set of all vectors having inner product 0 with all the vectors in V .

Clearly the orthogonal complement of $\langle e_1 \rangle$ is $\langle e_1, e_2, e_4 \rangle$.

Levi subgroups

The Levi subgroups are $T, GL(2), GL(1) \times Sp(2)$, up-to conjugacy. According to our assumption for the characteristic $l, l \neq 2$. So from [Lu3], we can see that T and $GL(1) \times Sp(2)$ have cuspidal pairs. For T the parity condition has been checked in [JMW, 4.3].

In the case of $GL(1) \times Sp(2)$, the cuspidal pair is of the form $(\mathcal{O}_{prin}, \mathcal{L})$, where \mathcal{O}_{prin} is the Sp_2 -principal orbit in $\mathfrak{sp}(2) = \mathfrak{sl}_2$ and \mathcal{L} is the nontrivial $SL(2)$ -equivariant local system on \mathcal{O}_{prin} . The Levi $GL(1) \times Sp(2)$ comes from the root $\alpha = 2e_2$. The Levi and nilpotent subalgebras related to the root $\alpha = 2e_2$ are of the form

$$\mathfrak{l} = \left\{ \begin{pmatrix} a & 0 & 0 & 0 \\ 0 & b & 0 & c \\ 0 & 0 & -a & 0 \\ 0 & d & 0 & -b \end{pmatrix} \mid a, b, c, d \in \mathbb{C} \right\} \cong \mathfrak{sl}_2 \times \mathbb{C},$$

and

$$\mathfrak{u}_P = \left\{ \begin{pmatrix} 0 & x & z & y \\ 0 & 0 & y & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -x & 0 \end{pmatrix} \mid x, y, z \in \mathbb{C} \right\}.$$

Hence, $\mathfrak{p} = \left\{ \begin{pmatrix} a & x & z & y \\ 0 & b & y & c \\ 0 & 0 & -a & 0 \\ 0 & d & -x & -b \end{pmatrix} \mid a, b, c, d, x, y, z \in \mathbb{C} \right\}$. Now we want to calculate Ind with

respect to these parabolic and Levi. Recall the parabolic induction diagram,

$$\mathcal{N}_L \xleftarrow{\pi} \mathcal{N}_L + \mathfrak{u}_P \xrightarrow{e} G \times^P (\mathcal{N}_L + \mathfrak{u}_P) \xrightarrow{\mu} \mathcal{N}_G.$$

The crucial step is to calculate the push forward of the map μ . We can interpret the space $G \times^P (\mathcal{N}_L + \mathfrak{u}_P)$ in a different way,

$$\begin{aligned} G \times^P (\mathcal{N}_L + \mathfrak{u}_P) &\cong \{(gP, x) \in G/P \times \mathcal{N}_G \mid \text{Ad}(g^{-1})x \in \mathcal{N}_L + \mathfrak{u}_P\}, \text{ by } (g, x) \rightarrow (gP, \text{Ad}(g)x). \\ &\cong \{(gP, x) \in G/P \times \mathcal{N}_G \mid \text{Ad}(g^{-1})x \in \text{Lie}(P)\}, \text{ as } x \in \mathcal{N}_G, \text{Ad}(g^{-1})x \in \mathcal{N}_G. \end{aligned}$$

$\text{Ad}(g^{-1})x \in \mathfrak{p}$ means it preserves the partial flag $\langle e_1 \rangle \subset \langle e_1, e_2, e_4 \rangle$, call it E . This implies that x preserves gE . Hence the definition becomes

$$G \times^P (\mathcal{N}_L + \mathfrak{u}_P) = \left\{ (V_1 \subset V_3, x) \left| \begin{array}{l} V_1 \subset V_3 \text{ is a partial flag of dimension 1 and 3,} \\ V_1^\perp = V_3, x \in \mathcal{N}_G \text{ preserves } V_3 \text{ and } V_1 \end{array} \right. \right\}. \quad (1.1)$$

Now the map μ becomes projection on the second coordinate. By [CM, 5.2], we can find the orbits in \mathfrak{sp}_4 , which are $\mathcal{O}[4], \mathcal{O}[2, 1^2], \mathcal{O}[2^2], \mathcal{O}[1^4]$. The representatives from these orbits are,

$$\begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \{0\},$$

respectively. Now we are interested in the fibers of the representatives of each orbits. For each orbit as above, we call x to be the representative.

Lemma 8.1.1. *For x defined above, $\mu^{-1}(x)$ has the following descriptions,*

1. for $x \in \mathcal{O}[2, 1^2]$, $\mu^{-1}(x) \cong \mathbb{P}^2$,
2. for $x \in \mathcal{O}[2^2]$, $\mu^{-1}(x) \cong \mathbb{P}^1$,
3. for $x \in \mathcal{O}[4]$, $\mu^{-1}(x) \cong \{pt\}$,
4. for $x \in \mathcal{O}[1^4]$, $\mu^{-1}(x) \cong G/P$.

Proof. 1. If $x \in \mathcal{O}[2, 1^2]$, then $\ker x = \langle e_1, e_2, e_4 \rangle$. For a flag $V_1 \subset V_3$ in $G \times^P (\mathcal{N}_L + \mathfrak{u}_P)$, x stabilizes both V_3 and V_1 . So if $\langle v \rangle = V_1$, then v should either go to 0 or to some scalar multiplication of v under the map x . Let $v = ae_1 + be_2 + ce_3 + de_4$, then $x.v = ce_1$. If $c = 0$, then $x.v = 0$ which means $v \in \ker x$. If $c \neq 0$ then $x.v \in V_1$, so $b = c = d = 0$, a contradiction. Hence $V_1 \subset \ker x$ which is 3 dimensional. Once we choose V_1 , V_3 is automatically determined by the condition $V_1^\perp = V_3$. Hence $\mu^{-1}(x) \cong \mathbb{P}^2$.

2. Now for $\mathcal{O}[2^2]$, to find the fiber we will proceed as before. For $\mathcal{O}[2^2]$, $\ker x = \langle e_1, e_2 \rangle$. If v is the generator of V_1 then $x.v$ is either 0 or in $\langle v \rangle$. Now if $v = ae_1 + be_2 + ce_3 + de_4$, then $x.v = ce_1 + de_2$. Hence either $c = d = 0$ implying $v \in \ker x$, otherwise, $ce_1 + de_2 = \lambda v$ implying $c = d = 0$, a contradiction. Therefore, $V_1 \subset \ker x$ which is two-dimensional. Hence in this case $\mu^{-1}(x) \cong \mathbb{P}^1$.

3. For $\mathcal{O}[4]$, we can check that $\langle e_1 \rangle \subset \langle e_2, e_1, e_4 \rangle$ is the only flag which satisfies all the conditions to be in the inverse image, so the fiber is just a point.

4. For $\mathcal{O}[1^4]$ the fiber is the whole space G/P .

□

We can find the dimension and the fundamental groups of the orbits from [CM].

Now as $L \cong SL_2 \times \mathbb{C}^\times$, hence the orbits in L are \mathcal{O}_{prin} and $\{0\}$. We want to calculate

$\mu^{-1}(x) \cap G \times^P (\mathcal{O}_0 + \mathfrak{u}_P)$ and $\mu^{-1}(x) \cap G \times^P (\mathcal{O}_{prin} + \mathfrak{u}_P)$ for each representative x .

Lemma 8.1.2. *Let $x \in \mathcal{N}_L + \mathfrak{u}_P$. Then,*

1. $x \in \mathcal{O}_{prin} + \mathfrak{u}_P$ if and only if the action of x on $\langle e_1, e_2, e_4 \rangle / \langle e_1 \rangle$ is nonzero,
2. $x \in \mathcal{O}_0 + \mathfrak{u}_P$ if and only if the action of x on $\langle e_1, e_2, e_4 \rangle / \langle e_1 \rangle$ is zero.

The proof follows from some simple matrix calculations.

Thus from the above lemma and the definition that we gave in the beginning,

$$G \times^P (\mathcal{O}_0 + \mathfrak{u}_P) = \left\{ (gP, x) \in G/P \times \mathcal{N}_G \left| \begin{array}{l} Ad(g^{-1})x \text{ preserves the flag } \langle e_1 \rangle \subset \langle e_1, e_2, e_3 \rangle \\ \text{and } Ad(g^{-1})x \text{ is zero on } \langle e_1, e_2, e_3 \rangle / \langle e_1 \rangle \end{array} \right. \right\}.$$

Which is same as,

$$G \times^P (\mathcal{O}_0 + \mathfrak{u}_P) = \left\{ (gP, x) \in G/P \times \mathcal{N}_G \left| \begin{array}{l} x \text{ preserves the flag } g.\langle e_1 \rangle \subset g.\langle e_1, e_2, e_3 \rangle \\ \text{and } x \text{ is zero on } g.\langle e_1, e_2, e_3 \rangle / g.\langle e_1 \rangle \end{array} \right. \right\}.$$

Which is again same as,

$$\left\{ (V_1 \subset V_3, x) \left| \begin{array}{l} V_1 \subset V_3 \text{ is a partial flag of dimension 1 and 3,} \\ V_1^\perp = V_3, x \in \mathcal{N}_G \text{ preserves } V_3 \text{ and } V_1 \text{ with } x \text{ is 0 on } V_3/V_1 \end{array} \right. \right\}.$$

Therefore for each representative x ,

$$\mu^{-1}(x) \cap G \times^P (\mathcal{O}_0 + \mathfrak{u}_P) = \left\{ V_1 \subset V_3 \left| \begin{array}{l} V_1 \subset V_3 \text{ is a partial flag of dimension 1 and 3,} \\ V_1^\perp = V_3, x \text{ preserves } V_3 \text{ and } V_1 \text{ with } x \text{ is 0 on } V_3/V_1 \end{array} \right. \right\}.$$

Lemma 8.1.3. *For x being the representative of each orbit, $\mu^{-1}(x) \cap G \times^P (\mathcal{O}_0 + \mathfrak{u}_P)$*

satisfies the fifth column in the table given below.

Proof. 1. For $\mathcal{O}[1^4]$, it is not hard to see, $\mu^{-1}(x) \cap G \times^P (\mathcal{O}_0 + \mathfrak{u}_P) = G/P$.

2. For $\mathcal{O}[2, 1^2]$, we have already seen that if $\langle v_1 \rangle \subset \langle v_1, v_2, v_3 \rangle$ is in $\mu^{-1}(x)$, then $v_1 \in \ker x = \langle e_1, e_2, e_4 \rangle$. As x is zero on $\langle v_1, v_2, v_3 \rangle / \langle v_1 \rangle$ that means v_2 and v_3 should either go to 0 or to $\langle v_1 \rangle$. If both go to 0, that is both are in the kernel, then $\langle v_1, v_2, v_3 \rangle = \langle e_1, e_2, e_4 \rangle$. Using the condition $V_1^\perp = V_3$, definitely $V_1 = \langle e_1 \rangle$. If one of them goes to v_1 , let's say v_2 . But then $x.v_2$ is some scalar multiplication of e_1 as $Im(x) = \langle e_1 \rangle$. This implies $\langle v_1 \rangle = \langle e_1 \rangle$. Again using the fact that $\langle v_1 \rangle^\perp = \langle v_1, v_2, v_3 \rangle$, we can see, $\langle v_1, v_2, v_3 \rangle = \langle e_1, e_2, e_4 \rangle$. Hence $\mu^{-1}(x) \cap G \times^P (\mathcal{O}_0 + \mathfrak{u}_P)$ is a single point $\{\langle e_1 \rangle \subset \langle e_1, e_2, e_4 \rangle\}$.

3. Now for $\mathcal{O}[2^2]$, again $\langle v_1 \rangle \subset \ker x$. But here $\ker x = \{e_1, e_2\}$. This means v_1 is of the form $ae_1 + be_2$. If $a = 0$ that means $\langle v_1 \rangle = \langle e_2 \rangle$ and $v_1^\perp = \langle e_2, e_1, e_3 \rangle$. But e_3 goes to e_1 under x , so the map does not induce a zero map on the quotient $\langle v_1, v_2, v_3 \rangle / \langle v_1 \rangle$. Similarly if $b = 0$ then $\langle v_1 \rangle = \langle e_1 \rangle$, therefore $v_1^\perp = \langle e_2, e_1, e_4 \rangle$. But e_4 goes to e_2 under x , so the map does not induce a 0 map on the quotient $\langle v_1, v_2, v_3 \rangle / \langle v_1 \rangle$. Now we consider the case where both a and b are non-zero. We have $\langle v_1 \rangle^\perp = \langle e_1, e_2, be_3 - ae_4 \rangle$. Now under the map x , e_1, e_2 both goes to 0 but $be_3 - ae_4$ goes to $be_1 - ae_2$. The action of x should be zero on the quotient, that means $be_1 - ae_2$ must be a scalar multiplication of v_1 . This implies $b^2 + a^2 = 0$ or $a = \pm ib$. Therefore the flags that satisfy all the conditions to be in $\mu^{-1}(x) \cap G \times^P (\mathcal{O}_0 + \mathfrak{u}_P)$ are $\langle ie_1 + e_2 \rangle \subset \langle e_1, e_2, e_3 - ie_4 \rangle$ and $\langle -ie_1 + e_2 \rangle \subset \langle e_1, e_2, e_3 + ie_4 \rangle$.
4. For $\mathcal{O}[4]$, $\langle e_1 \rangle \subset \langle e_2, e_1, e_4 \rangle$ is the only flag in the inverse image but it does not satisfy this condition hence $\mu^{-1}(x) \cap G \times^P (\mathcal{O}_0 + \mathfrak{u}_P) = \emptyset$.

□

Table 8.1. Orbits in \mathfrak{sp}_4

orbits:	$\mathcal{O}[4]$	$\mathcal{O}[2^2]$	$\mathcal{O}[2, 1^2]$	$\mathcal{O}[1^4]$
dim :	8	6	4	0
$\pi_1 :$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	0
$\mu^{-1}(x) :$	$\{pt\}$	\mathbb{P}^1	\mathbb{P}^2	G/P
$\mu^{-1}(x) \cap G \times^P (\mathcal{O}_0 + \mathfrak{u}_P) :$	\emptyset	$\{pt\} \sqcup \{pt\}$	$\{pt\}$	G/P
$\mu^{-1}(x) \cap G \times^P (\mathcal{O}_{prin} + \mathfrak{u}_P) :$	$\{pt\}$	$\mathbb{A}^1 - \{pt\}$	$\mathbb{P}^2 - \{pt\}$	\emptyset

We are now ready to calculate Ind_P^G for cuspidal pairs. The only cuspidal pair on SL_2 is $(\mathcal{O}_{prin}, \mathcal{L})$, where \mathcal{L} is the nontrivial local system on \mathcal{O}_{prin} . Now recall the parabolic induction diagram for cuspidal pair defined in ???. As $(\mathcal{O}_{prin}, \mathcal{L})$ is cuspidal, so $\text{Ind}_P^G \mathcal{IC}(\mathcal{O}_{prin}, \mathcal{L}) = c_!(b^* \text{For}_P^G)^{-1} a^* \mathcal{L}[\dim \mathcal{O}_{prin}] = c_!(b^* \text{For}_P^G)^{-1} a^* \mathcal{L}[2]$. As we know pull-back of some local system is again a local system, thus $(b^* \text{For}_P^G)^{-1} a^* \mathcal{L}[2]$ is a local

system on $G \times^P (\mathcal{O}_{prin} + \mathfrak{u}_P)$.

$$\begin{array}{ccc} G \times^P (\mathcal{O}_{prin} + \mathfrak{u}_P) & \xrightarrow{c} & \mathcal{N}_G \\ \uparrow & & \uparrow \\ G \times^P (\mathcal{O}_{prin} + \mathfrak{u}_P) \cap \mu^{-1}(x) & \xrightarrow{c} & x \end{array}$$

Using the above diagram, $\text{Ind}_P^G \mathcal{IC}(\mathcal{O}_{prin}, \mathcal{L})_x$ becomes the $!$ -pushforward of a local system on $G \times^P (\mathcal{O}_{prin} + \mathfrak{u}_P) \cap \mu^{-1}(x)$ by a constant map. From the table above, $G \times^P (\mathcal{O}_{prin} + \mathfrak{u}_P) \cap \mu^{-1}(x)$ is simply connected for $\mathcal{O}[4]$, $\mathcal{O}[2, 1^2]$ and $\mathcal{O}[1^4]$. A local system on a simply connected space is constant sheaf. Therefore for these orbits, each stalk of $\text{Ind}_P^G \mathcal{IC}(\mathcal{O}_{prin}, \mathcal{L})$ is the cohomology of $G \times^P (\mathcal{O}_{prin} + \mathfrak{u}_P) \cap \mu^{-1}(x)$. But for $\mathcal{O}[2^2]$, $G \times^P (\mathcal{O}_{prin} + \mathfrak{u}_P) \cap \mu^{-1}(x)$ is $\mathbb{A}^1 - \{pt\}$, which is not simply connected. Here we will abuse the notation little bit, both the representative of $\mathcal{O}[2^2]$ and its image under the projection $\mathcal{N}_P \rightarrow \mathcal{N}_L$ will be called x . Recall we started with a nontrivial L -equivariant local system on \mathcal{O}_{prin} . The projection $\pi : \mathcal{O}_{prin} + \mathfrak{u}_P \rightarrow \mathcal{O}_{prin}$ is a trivial vector bundle, hence induces isomorphism of the equivariant fundamental groups. The inclusion $\mathcal{O}_{prin} + \mathfrak{u}_P \hookrightarrow G \times^P (\mathcal{O}_{prin} + \mathfrak{u}_P)$ induces isomorphism on the equivariant fundamental groups via induction equivalence. So the pullback of the local system we started with is still a nontrivial local system on $G \times^P (\mathcal{O}_{prin} + \mathfrak{u}_P)$.

Let

$$S = \left\{ \begin{pmatrix} A & 0 \\ 0 & A \end{pmatrix} \mid A = \begin{pmatrix} a & b \\ -b & a \end{pmatrix}, a, b \in \mathbb{C}, a^2 + b^2 = 1 \right\}.$$

It is not hard to check $S \subset G^x$. If we choose a flag in $\mu^{-1}(x) \cap G \times^P (\mathcal{O}_{prin} + \mathfrak{u}_P)$,

say $F = \langle e_1 \rangle \subset \langle e_1, e_2, e_4 \rangle$ then we can see the elements of S that fix the flag F are $\left\{ \begin{pmatrix} Id & 0 \\ 0 & Id \end{pmatrix}, \begin{pmatrix} -Id & 0 \\ 0 & -Id \end{pmatrix} \right\}$. Therefore,

$$S^F/(S^F)^\circ = \left\{ \begin{pmatrix} Id & 0 \\ 0 & Id \end{pmatrix}, \begin{pmatrix} -Id & 0 \\ 0 & -Id \end{pmatrix} \right\} \cong \mathbb{Z}/2\mathbb{Z}.$$

Now we aim to show $S^F/(S^F)^\circ \cong L^x/(L^x)^\circ$. It is not hard to see

$$L^x \cong \left\{ \begin{pmatrix} a & & & \\ & b & & \\ & & a^{-1} & \\ & & & b \end{pmatrix} \mid a \in \mathbb{C}, b = \pm 1 \right\} \cong G_m \times \{\pm 1\}.$$

Hence $L^x/(L^x)^\circ \cong \mathbb{Z}/2\mathbb{Z}$ and the map $S^F/(S^F)^\circ \rightarrow L^x/(L^x)^\circ$ is an isomorphism. Therefore

$(b^* \text{For}_P^G)^{-1} a^* \mathcal{L}[2]|_{\mu^{-1}(x) \cap G \times^P (\mathcal{O}_{\text{prin}} + \mathfrak{u}_P)}$ is a nontrivial local system.

For a connected, locally contractible space X if the universal cover is contractible, then the inclusion functor $\text{Loc}(X, \mathbb{k}) \rightarrow \text{Sh}(X, \mathbb{k})$ induces an equivalence of categories,

$$D^b \text{Loc}(X, \mathbb{k}) \rightarrow D_{\text{loc}}^b(X, \mathbb{k}).$$

It can be proved by a minor variation on the proof that the (co)homology of an Eilenberg-MacLane space is isomorphic to group (co)homology [Bro, Prop II.4.1]. For $X = \mathbb{A}^1 - \{pt\}$, $\text{Loc}(X, \mathbb{k}) \cong \mathbb{k}[\pi_1(\mathbb{A}^1 - \{pt\})] - \text{mod}$, which is same as $\mathbb{k}[\mathbb{Z}] - \text{mod} \cong \mathbb{k}[T, T^{-1}] - \text{mod}$. Therefore, for the local system $(b^* \text{For}_P^G)^{-1} a^* \mathcal{L}[2]|_{\mu^{-1}(x) \cap G \times^P (\mathcal{O}_{\text{prin}} + \mathfrak{u}_P)}$, there exists a $\mathbb{k}[T, T^{-1}]$ module M on which T acts by (-1) . To calculate the cohomology of this local system is same as calculating $R\text{Hom}(\mathbb{k}, M)$. Now,

$$\rightarrow \mathbb{k}[T, T^{-1}] \xrightarrow{\times(T-1)} \mathbb{k}[T, T^{-1}] \xrightarrow{T-1} \mathbb{k}$$

is a projective resolution of \mathbb{k} . Applying $\text{Hom}(\cdot, M)$ we get, M in degree 0 and 1.

$$\rightarrow 0 \rightarrow M \xrightarrow{\times(-2)} M \rightarrow 0 \rightarrow .$$

Multiplying by -2 induces isomorphism. Hence $R\text{Hom}(\mathbb{k}, M)$ is 0 in every degree.

Table 8.2. Stalks of $\text{Ind}_P^G \mathcal{IC}(\mathcal{O}_{prin}, \mathcal{L})$

dim	$\mathcal{O}[4]$	$\mathcal{O}[2^2]$	$\mathcal{O}[2, 1^2]$	$\mathcal{O}[1^4]$
0				
-1				
-2			rank 1	
-3				
-4			rank 1	
-5				
-6				
-7				
-8				
-9				
-10	rank 1			

Hence the parity condition of Conjecture 3.2.9 is satisfied.

8.2. \mathfrak{sl}_4 -case:

Let $G = SL_4$. First we talk about the Levi subgroups of G and find out which of them have cuspidal pairs. The conjugacy classes of proper Levis are of the form

$$S(GL_3 \times GL_1), S(GL_2 \times GL_2),$$

$S(GL_2 \times GL_1 \times GL_1), T$. Here $S(GL_m \times GL_n) = \left\{ \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \mid A \in GL_m, B \in GL_n, \det(A) \det(B) = 1 \right\}$. According to the discussion in 6.2 and Theorem 6.3 in

[AJHR3], we can see cuspidal pair only appears for $S(GL_2 \times GL_2)$ and is of the form

$(\mathcal{O}_{prin} \times \mathcal{O}_{prin}, \mathcal{L} \boxtimes \mathcal{L})$. Here each \mathcal{L} is a rank one $SL(2)$ -equivariant local system on \mathcal{O}_{prin}

and \mathcal{O}_{prin} is the SL_2 -principle nilpotent orbit in \mathfrak{sl}_2 .

For \mathfrak{sl}_4 , the root system is $\Phi = \{e_i - e_j \mid i \neq j, 1 \leq i, j \leq 4\}$. The parabolic subgroup

associated to $\{e_1 - e_2, e_3 - e_4\}$ is of the form,

$$\begin{pmatrix} * & * & * & * \\ * & * & * & * \\ & & * & * \\ & & * & * \end{pmatrix}. \text{ The Levi subgroup is then}$$

$S(GL_2 \times GL_2)$ and the unipotent radical is of the form,

$$\begin{pmatrix} & & * & * \\ & & * & * \\ & & & \\ & & & \end{pmatrix}. \text{ Now the genera-}$$

tors of the nilpotent orbits come from the Jordan block of size depending on the partition.

Hence the representatives of $\mathcal{O}[4], \mathcal{O}[3, 1], \mathcal{O}[2^2], \mathcal{O}[2, 1^2], \mathcal{O}[1^4]$ are respectively,

$$\begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \{0\}.$$

Now we calculate $\mu^{-1}(x)$ for each x as we did for \mathfrak{sp}_4 . Here again,

$$G \times^P (\mathcal{N}_L + \mathfrak{u}_P) = \{(gP, x) \in G/P \times \mathcal{N}_G \mid Ad(g^{-1})x \in \text{Lie}(P)\}.$$

But $Ad(g^{-1})x \in \mathfrak{p}$ means it preserves the two dimensional subspace $\langle e_1, e_2 \rangle$. Hence,

$$G \times^P (\mathcal{N}_L + \mathfrak{u}_P) = \{(H, x) \mid x \in \mathcal{N}_G, H \text{ is a two dimensional subspace preserved by } x\},$$

Lemma 8.2.1. *Let $x \in \mathcal{N}_L + \mathfrak{u}_P$. Then, $x \in \mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P$ if and only if $x|_{\langle e_1, e_2 \rangle} \neq 0$ and $x|_{\mathbb{C}^4 / \langle e_1, e_2 \rangle} \neq 0$.*

The proof follows from some easy matrix calculations. Therefore following the same

process as for \mathfrak{sp}_4 ,

$$\mu^{-1}(x) \cap (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P) = \{(H, x) \mid x \text{ preserves the subspace } H, x|_H \neq 0, x|_{\mathbb{C}^4/H} \neq 0\}.$$

Lemma 8.2.2. *Let x be the representative of each orbits in \mathfrak{sl}_4 .*

1. For $\mathcal{O}[4]$, $\mu^{-1}(x)$ is $\{\langle e_1, e_2 \rangle\}$ and $\mu^{-1}(x) \cap (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P) = \{\langle e_1, e_2 \rangle\}$.
2. For $\mathcal{O}[3, 1]$, $\mu^{-1}(x) \cong \mathbb{P}^1$ and $\mu^{-1}(x) \cap (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P) \cong \mathbb{P}^1 - \{[0, 1], [1, 0]\}$.
3. For $\mathcal{O}[2, 1^2]$, if x preserves H , then either $\langle e_1 \rangle \subset H$ or $H \subset \ker(x)$. Also $\mu^{-1}(x) \cong \mathbb{P}^2 \sqcup_{\mathbb{P}^1} \mathbb{P}^2$ and $\mu^{-1}(x) \cap (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P) \cong \emptyset$.
4. For $x \in \mathcal{O}[2^2]$, $\mu^{-1}(x) \cap (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P) \cong \mathbb{A}^1 \sqcup \mathbb{A}^2$.

Proof. 1. For $\mathcal{O}[4]$, the only choice for $\mu^{-1}(x)$ is $\langle e_1, e_2 \rangle$. Now $x|_{\langle e_1, e_2 \rangle} \neq 0$ and $x|_{\mathbb{C}^4/\langle e_1, e_2 \rangle} \neq 0$, hence $\mu^{-1}(x) \cap (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P) = \{\langle e_1, e_2 \rangle\}$.

2. The first claim is that if $H \in \mu^{-1}(x)$, then H contains e_1 . Let H does not contains e_1 . if H contains $v = ae_1 + be_2 + ce_3 + de_4$, then as $x.v$ is in H , so $be_1 + ce_2 \in H$. If both $b = c = 0$, then $v \in \ker(x) = \langle e_1, e_4 \rangle$. If $v \neq e_1$ then it is a linear combination of e_1 and e_4 . In this case the other basis element of H must be a linear combination of e_2 and e_3 , which contradicts the fact that H x -invariant. So b and c both can not be 0. If $c = 0$ we are done. If $c \neq 0$ then $x.(x.v) = ce_1 \in H$, therefore $e_1 \in H$.

Now as e_1 is fixed, we have one choice left for the second generator. Now let the second generator $v = ae_2 + be_3 + de_4$, then $x.v = ae_1 + ce_2$. As H is stable under x , so $x.v$ must be a scalar multiple of e_1 or v . In both cases $c = 0$, therefore v can be linear combination of e_2 and e_4 . So we have $\langle e_1 \rangle \subset H \subset \langle e_1, e_2, e_4 \rangle$ and $\mu^{-1}(x) \cong \mathbb{P}^1$.

For $\mu^{-1}(x) \cap (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P)$, $x|_H \neq 0$. So again if we take the other generator v in H , which is of the form $ae_2 + be_4$, then using the condition $x|_H \neq 0$ we can say $a \neq 0$. Now if $b = 0$, then $H = \langle e_1, e_2 \rangle$, which implies $x|_{\mathbb{C}^4/H} = 0$. Therefore for H to be in $\mu^{-1}(x) \cap (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P)$, a and b both must be nonzero. So $\mu^{-1}(x) \cap (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P) \cong \mathbb{P}^1 - \{[0, 1], [1, 0]\}$.

3. Let H is not contained in $\ker(x) = \langle e_1, e_3, e_4 \rangle$. Let $v = ae_1 + be_2 + ce_3 + de_4 \in H$ with $b \neq 0$, then $x.v = be_1 \in H$. Hence H contains e_1 . Now if $\langle e_1 \rangle \subset H \subset \mathbb{C}^4$, then the choice for H is \mathbb{P}^2 . If $H \subset \ker(x) = \langle e_1, e_3, e_4 \rangle$, then again choice is \mathbb{P}^2 . If $\langle e_1 \rangle \subset H \subset \ker(x)$, then the choice is \mathbb{P}^1 . Hence $\mu^{-1}(x) \cong \mathbb{P}^2 \sqcup_{\mathbb{P}^1} \mathbb{P}^2$. For $\mu^{-1}(x) \cap (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P)$, $x|_H \neq 0$, so H can not be contained in $\ker(x)$. But

still whatever be the choice of the other generator we can see $x|_{\mathbb{C}^4/H}$ is always 0. Therefore $\mu^{-1}(x) \cap (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P) = \emptyset$.

4. If $ae_1 + be_2 + ce_3 + de_4 \in H$, then $be_1 + de_3 \in H$. If $x|_H \neq 0$, then $H \cap \ker(x)$ is one dimensional. Call the subspace $\ker(x) \cap H$ to be L which is definitely generated by elements of the form $be_1 + de_3$. Now clearly $L \subset H \subset x^{-1}L = \langle e_1, e_3, be_2 + de_4 \rangle$. If both b and d are 0, then $H = \langle e_1, e_3 \rangle$. This implies $x|_H = 0$. The converse is also true. For H to be in $\mu^{-1}(x) \cap (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P)$ we need $x|_H \neq 0$ and $x|_{\mathbb{C}^4/H} \neq 0$, which is not true for the above case. So one of them must be non-zero. Now if we consider one of them is zero, say $d = 0$, then $L = \langle e_1 \rangle$. In this case we can consider $b = 1$, hence H is generated by e_1 and $e_2 + ce_3$, and in this case $x|_{\mathbb{C}^4/H} \neq 0$, so the choice is \mathbb{A}^1 . The remaining case is $d \neq 0$. Here we can consider $d = 1$ and then $L = \langle be_1 + e_3 \rangle$. In this case, H is generated by $be_1 + e_3$ and $ae_1 + be_2 + ce_3 + e_4$, which is the same as $\langle be_1 + e_3, a'e_1 + be_2 + e_4 \rangle$, so the choice is \mathbb{A}^2 . In this case also x is nonzero on both H and the quotient.

□

Table 8.3. Orbits in \mathfrak{sl}_4

orbits:	$\mathcal{O}[4]$	$\mathcal{O}[3, 1]$	$\mathcal{O}[2^2]$	$\mathcal{O}[2, 1^2]$	$\mathcal{O}[1^4]$
dim :	12	10	8	6	0
$\pi_1 :$	$\mathbb{Z}/4$	$\{1\}$	$\mathbb{Z}/2$	$\{1\}$	$\{1\}$
$\mu^{-1}(x) :$	$\{\langle e_1, e_2 \rangle\}$	\mathbb{P}^1	-	$\mathbb{P}^2 \sqcup_{\mathbb{P}^1} \mathbb{P}^2$	G/P
$\mu^{-1}(x) \cap G \times^P (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P) :$	$\{\langle e_1, e_2 \rangle\}$	$\mathbb{P}^1 - \{[0, 1], [1, 0]\}$	$\mathbb{A}^2 \sqcup \mathbb{A}^1$	\emptyset	\emptyset

Now we are ready to find the Ind_P^G . The only cuspidal pair in L is $(\mathcal{O}_{prin} \times \mathcal{O}_{prin}, \mathcal{L} \boxtimes \mathcal{L})$ where \mathcal{L} is the nontrivial local system on \mathcal{O}_{prin} . We know $\mathcal{IC}(\mathcal{O}_{prin} \times \mathcal{O}_{prin}, \mathcal{L} \boxtimes \mathcal{L}) \cong \mathcal{IC}(\mathcal{O}_{prin}, \mathcal{L}) \boxtimes \mathcal{IC}(\mathcal{O}_{prin}, \mathcal{L})$. We can use the parabolic induction diagram introduced in ??, therefore $\text{Ind}_P^G \mathcal{IC}(\mathcal{O}_{prin} \times \mathcal{O}_{prin}, \mathcal{L} \boxtimes \mathcal{L}) = c_!(b^* \text{For}_P^G)^{-1} a^*(\mathcal{L}[2] \boxtimes \mathcal{L}[2])$. Now we will follow the same steps as we did for \mathfrak{sp}_4 and using the same diagram, $\text{Ind}_P^G \mathcal{IC}(\mathcal{O}_{prin} \times \mathcal{O}_{prin}, \mathcal{L} \boxtimes \mathcal{L})_x$ becomes the !-pushforward of a local system on $G \times^P (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P) \cap \mu^{-1}(x)$ by a constant map. From the table above, $G \times^P (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P) \cap \mu^{-1}(x)$ is simply connected for $\mathcal{O}[4]$, $\mathcal{O}[2, 1^2]$, $\mathcal{O}[2^2]$ and $\mathcal{O}[1^4]$. A local system on a simply connected space is constant sheaf. Therefore

for these orbits, the stalks of $\text{Ind}_P^G \mathcal{IC}(\mathcal{O}_{prin} \times \mathcal{O}_{prin}, \mathcal{L} \boxtimes \mathcal{L})$ are the cohomologies of $G \times^P (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P) \cap \mu^{-1}(x)$. But for $\mathcal{O}[3, 1]$, $G \times^P (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P) \cap \mu^{-1}(x)$ is $\mathbb{P}^1 - \{[1, 0], [0, 1]\}$, which is not simply connected. Here we will use the same abuse of notation, both the representative of $\mathcal{O}[3, 1]$ and its image under the projection $\mathcal{N}_P \rightarrow \mathcal{N}_L$ will be called x . Recall we started with a nontrivial L -equivariant local system on $\mathcal{O}_{prin} \times \mathcal{O}_{prin}$. The projection $\pi : \mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P \rightarrow \mathcal{O}_{prin} \times \mathcal{O}_{prin}$ is a trivial vector bundle, hence induces isomorphism of the equivariant fundamental groups. The inclusion $\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P \hookrightarrow G \times^P (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P)$ induces isomorphism on the equivariant fundamental groups via induction equivalence. So the pullback of the local system we started with is still a nontrivial local system on $G \times^P (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P)$. Let

$$S = \left\{ \begin{pmatrix} A & 0 \\ 0 & A \end{pmatrix} \mid A = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}, a \in \mathbb{C}^\times \right\}.$$

and surely $S \subset G^x$. If we choose a subspace in $\mu^{-1}(x) \cap G \times^P (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P)$, say $H = \langle e_1, e_2 + e_4 \rangle$ then we can see S stabilizes H , so $S^H = S$. Therefore, $S^H / (S^H)^\circ$ is trivial. Now we aim to show $S^H / (S^H)^\circ \cong L^x / (L^x)^\circ$. It is not hard to see

$$L^x \cong \left\{ \begin{pmatrix} a & & & \\ & b & & \\ & & a^{-1} & \\ & & & c & b^{-1} \end{pmatrix} \mid a, b \in \mathbb{C}^\times \text{ and } c \in \mathbb{C} \right\} \cong \mathbb{C}^\times \times \mathbb{C}^\times \times \mathbb{C}.$$

Hence $L^x / (L^x)^\circ$ is also trivial and the map $S^H / (S^H)^\circ \rightarrow L^x / (L^x)^\circ$ is an isomorphism.

Therefore $(b^* \text{For}_P^G)^{-1} a^* (\mathcal{L}[2] \boxtimes \mathcal{L}[2])|_{\mu^{-1}(x) \cap G \times^P (\mathcal{O}_{prin} \times \mathcal{O}_{prin} + \mathfrak{u}_P)}$ is a nontrivial local system.

Now again using the same argument from [Bro, Prop. II.4.1],

$$D^b \text{Loc}(X, \mathbb{k}) \rightarrow D_{loc}^b(X, \mathbb{k}).$$

Here $X = \mathbb{P}^1 - \{[1, 0], [0, 1]\}$, $\text{Loc}(X, \mathbb{k}) \cong \mathbb{k}[\pi_1(\mathbb{P}^1 - \{[0, 1], [1, 0]\})] - \text{mod}$, which is same as $\mathbb{k}[\mathbb{Z}] - \text{mod} \cong \mathbb{k}[T, T^{-1}] - \text{mod}$. Therefore, for the local system $(b^* \text{For}_P^G)^{-1} a^* \mathcal{L}[2] \boxtimes \mathcal{L}[2]|_{\mu^{-1}(x) \cap G \times^P (\mathcal{O}_{\text{prin}} \times \mathcal{O}_{\text{prin}+u_P})}$, there exists a $\mathbb{k}[T, T^{-1}]$ module M on which T acts by (-1) . Now we use the same calculation as we did for \mathfrak{sp}_4 to conclude $R\text{Hom}(\mathbb{k}, M)$ is 0 in every degree.

Table 8.4. Stalks of $\text{Ind}_P^G \mathcal{IC}(\mathcal{O}_{\text{prin}} \times \mathcal{O}_{\text{prin}}, \mathcal{L} \boxtimes \mathcal{L})$

dim	$\mathcal{O}[4]$	$\mathcal{O}[3, 1]$	$\mathcal{O}[2^2]$	$\mathcal{O}[2, 1^2]$	$\mathcal{O}[1^4]$
-6					
-7					
-8				rank 1	
-9					
-10			rank 1		
-11					
-12					
-13					
-14	rank 1				
-15					
-16					

Hence the parity condition is again satisfied.

8.3. Construction of \mathfrak{p} , \mathfrak{n} , and \mathfrak{l} described in 7.1.1

Let $G = SL_4$ and $\chi : \mathbb{C}^\times \rightarrow G$ be defined as $t \rightarrow (t, 1, 1, t^{-1})$. Then the matrix that gives $\mathfrak{g}_{m'}$ for all m' is

$$\begin{pmatrix} 0 & 1 & 1 & 2 \\ -1 & 0 & 0 & 1 \\ -1 & 0 & 0 & 1 \\ -2 & -1 & -1 & 0 \end{pmatrix}. \quad (3.2)$$

Now $\mathfrak{g}_{m'}$ comes from the above matrix by putting nonzero entries wherever we have m' in (3.2) and 0 elsewhere. For example,

$$\mathfrak{g}_2 = \begin{pmatrix} * \\ \end{pmatrix}, \mathfrak{g}_1 = \begin{pmatrix} * & * \\ & * \\ & * \\ \end{pmatrix}, \text{ and}$$

$$\mathfrak{g}_0 = \begin{pmatrix} * \\ & * & * \\ & * & * \\ & & * \\ \end{pmatrix}.$$

Similarly we can find $\mathfrak{g}_{-1}, \mathfrak{g}_{-2}$.

Now choose a point $x \in \mathfrak{g}_{-1}$. We can think of \mathfrak{g}_{-1} as $\text{Hom}(\mathbb{C}, \mathbb{C}^2) \times \text{Hom}(\mathbb{C}^2, \mathbb{C})$, which is a space of representations of quivers of finite type of dimension $(1, 2, 1)$. By [DW, Theorem 4.3.9], isomorphism classes of G_0 -orbits in \mathfrak{g}_{-1} are in bijection with the isomorphism classes of finite type quiver representations of dimension $(1, 2, 1)$. This is again a linear combination of roots of A_3 that add up to $\alpha_1 + 2\alpha_2 + \alpha_3$, where $\alpha_1, \alpha_2, \alpha_3$ are all the simple roots in A_3 and $\alpha_4 = \alpha_1 + \alpha_2, \alpha_5 = \alpha_1 + \alpha_2 + \alpha_3, \alpha_6 = \alpha_2 + \alpha_3$. Let us pick one such linear combination which gives a representative of that orbit and call it x . Note that $\alpha_4 + \alpha_6$ adds up to the desired sum. This gives the representative

$$x = \begin{pmatrix} 1 \\ 0 \\ 0 & 1 \end{pmatrix}$$

Note that the Jordan canonical form of x is

$$\begin{pmatrix} 0 & 1 & & \\ & 0 & & \\ & & 0 & 1 \\ & & & 0 \end{pmatrix}$$

This matrix is associated to the partition $[2, 2]$, hence we can use [CM, Lemma 3.2.6] to

find a map $\mathfrak{sl}_2 \rightarrow \mathfrak{g}$ which takes e to $\begin{pmatrix} 0 & 1 & & \\ & 0 & & \\ & & 0 & 1 \\ & & & 0 \end{pmatrix}$ and h to $\begin{pmatrix} 1 & & & \\ & -1 & & \\ & & 1 & \\ & & & -1 \end{pmatrix} \in \mathfrak{g}_0$. We

can see the matrix $\begin{pmatrix} & & & 1 \\ & & 1 & \\ & & & 1 \\ 1 & & & \end{pmatrix}$ conjugates x to $\begin{pmatrix} 0 & 1 & & \\ & 0 & & \\ & & 0 & 1 \\ & & & 0 \end{pmatrix}$, therefore conjugating

the above map by the same matrix we get a map $\phi : \mathfrak{sl}_2 \mapsto \mathfrak{g}$ which sends e to x and h to

$$\begin{pmatrix} -1 & & & \\ & 1 & & \\ & & -1 & \\ & & & 1 \end{pmatrix} \in \mathfrak{g}_0$$

Clearly, $\tilde{\phi} \begin{pmatrix} t \\ t^{-1} \end{pmatrix} = \begin{pmatrix} t^{-1} \\ t \\ t^{-1} \\ t \end{pmatrix}$. Hence we get the required $\chi' : \mathbb{C}^\times \rightarrow G$, as

$\chi'(t) = \tilde{\phi} \begin{pmatrix} t \\ t^{-1} \end{pmatrix} = \begin{pmatrix} t^{-1} \\ t \\ t^{-1} \\ t \end{pmatrix}$. So the matrix ${}_m\mathfrak{g}$ for all m comes from the

matrix below by the same procedure as above.

$$\begin{pmatrix} 0 & -2 & 0 & -2 \\ 2 & 0 & 2 & 0 \\ 0 & -2 & 0 & -2 \\ 2 & 0 & 2 & 0 \end{pmatrix}. \quad (3.3)$$

We can see what ${}_1\mathfrak{g}, {}_2\mathfrak{g}, {}_3\mathfrak{g}$ are as before. In this example $n = -1$. So we need conditions

on $m + 2m'$ to find $\mathfrak{p}, \mathfrak{n}, \mathfrak{l}$. The matrix ${}_m\mathfrak{g}_{m'}$ for $m + 2m'$ is given below.

$$\begin{pmatrix} 0 & 0 & 2 & 2 \\ 0 & 0 & 2 & 2 \\ -2 & -2 & 0 & 0 \\ -2 & -2 & 0 & 0 \end{pmatrix}.$$

Hence, with the conditions on $m + 2m'$ we can say,

$$\mathfrak{l} = \begin{pmatrix} * & * & & & & \\ * & * & & & & \\ & & * & * & & \\ & & * & * & & \\ & & & & & \end{pmatrix}.$$

Table 8.5. Table for the Levi

x	Representative	dim	Associated Levi
$\alpha_2 + \alpha_3 + \alpha_4$	$\begin{pmatrix} 0 & & & & & \\ & 0 & 0 & & & \\ & & & & & \end{pmatrix}$	2	$\begin{pmatrix} * & * & * & & & \\ * & * & * & & & \\ * & * & * & & & \\ & & & & & * \\ & & & & & \end{pmatrix}$
$\alpha_6 + \alpha_4$	$\begin{pmatrix} 1 & & & & & \\ 0 & & & & & \\ & 0 & 1 & & & \\ & & & & & \end{pmatrix}$	3	$\begin{pmatrix} * & * & & & & \\ * & * & & & & \\ & & & & * & * \\ & & & & * & * \\ & & & & & \end{pmatrix}$
$\alpha_2 + \alpha_5$	$\begin{pmatrix} 0 & & & & & \\ 1 & & & & & \\ & 0 & 1 & & & \\ & & & & & \end{pmatrix}$	4	$\begin{pmatrix} * & * & & & & \\ * & * & & & & \\ & & & & * & \\ & & & & & * \\ & & & & & \end{pmatrix}$
$\alpha_1 + \alpha_2 + \alpha_6$	$\begin{pmatrix} 0 & & & & & \\ 0 & & & & & \\ & 0 & 1 & & & \\ & & & & & \end{pmatrix}$	2	$\begin{pmatrix} * & & & & & \\ & * & & & & \\ & & & & * & * \\ & & & & * & * \\ & & & & & \end{pmatrix}$
$\alpha_1 + 2\alpha_2 + \alpha_3$	$\{0\}$	0	G

Bibliography

- [AJHR2] P. Achar, A. Henderson, D. Juteau, S. Riche. *Modular Generalized Springer Correspondence I: the general linear group*. J. Eur. Math. Soc. (JEMS) 18 (2016), no. 7, 14051436.
- [AJHR3] P. Achar, A. Henderson, D. Juteau, S. Riche. *Modular Generalized Springer Correspondence II: classical groups*. J. Eur. Math. Soc. (JEMS) 19 (2017), no. 4, 10131070.
- [AJHR] P. Achar, A. Henderson, D. Juteau, S. Riche. *Constructible sheaves on nilpotent cones in rather good characteristic*. Selecta Math. (N.S.) 23 (2017), no. 1, 203243.
- [BBD] A. Beilinson, J. Bernstein, and P. Deligne, *Faisceaux pervers, Analyse et topologie sur les espaces singuliers, I* (Luminy, 1981), Asterisque, vol. 100, Soc. Math. France, Paris, 1982, pp. 5171.
- [BO] J. Brundan and V. Ostrik. *Cohomology of Spaltenstein varieties*. Transform. Groups, 16(3):619648, 2011.
- [BL] J. Bernstein, V. Lunts. *Equivariant sheaves and functors*, Lecture Notes in Math., vol. 1578, Springer-Verlag, Berlin, 1994
- [BR] Tom. Braden. *Hyperbolic Localization of Intersection Cohomology* Transform. Groups 8 (2003), no. 3, 209216.
- [Bro] Brown, Kenneth S. *Cohomology of groups*. Corrected reprint of the 1982 original. Graduate Texts in Mathematics, 87. Springer-Verlag, New York, 1994. ISBN: 0-387-90688-6 20J05 (20-02)
- [Car] R. W. Carter. *Finite groups of Lie type. Conjugacy classes and complex characters*. Reprint of the 1985 original. Wiley Classics Library. A Wiley-Interscience Publication. John Wiley and Sons, Ltd., Chichester, 1993. xii+544 pp. ISBN: 0-471-94109-3 20C33 (20-02 20G40)
- [CM] D. H. Collingwood and W. M. McGovern. *Nilpotent orbits in semisimple Lie Algebras*. Van Nostrand Reinhold Mathematics Series. Van Nostrand Reinhold Co., New York, 1993.
- [CR] C. W. Curtis, I. Reiner. *Representation theory of finite groups and associative algebras*. ISBN 0-8218-4066-5.
- [DCLP] C. De Concini, G. Lusztig, and C. Procesi. *Homology of the zero-set of a nilpotent vector field on a flag manifold*. J. Amer. Math. Soc., 1(1):1534, 1988.

- [DW] Harm Derksen and Jerzy Weyman *An Introduction to Quiver Representations*(2017). ISBN-10: 1-4704-2556-4 ISBN-13: 978-1-4704-2556-2.
- [Fr] L. Fresse. *Partial flag varieties and nilpotent elements*. arXiv:1305.3355, 2013.
- [GIN] V. Ginzburg. *Perverse sheaves on a Loop group and Langlands' duality*. <https://arxiv.org/abs/alg-geom/9511007>.
- [Hum] James. E. Humphreys *Linear Algebraic Group*. ISBN 978-1-4612-6398-2.
- [Her] S. Herpel. *On the smoothness of centralizers in reductive groups*. Trans. Amer. Math. Soc., 365(7):37533774, 2013.
- [J2] D. Juteau. *Decomposition numbers for perverse sheaves*. Ann. Inst. Fourier 59(2009), 1177-1229.
- [JMW] D. Juteau, C. Mautner, and G. Williamson. *Parity Sheaves*. Journal of the American Mathematical Society. Vol 27, Number 4, Oct 2014, 1169-1212.
- [KL] D. Kazhdan and G. Lusztig, *Proof of the Deligne-Langlands conjecture for Hecke Algebras*. Invent. Math. 87(1987). 153-215.
- [Lu] George Lusztig. *Study of perverse sheaves on Graded Lie Algebras*. Advances in Mathematics 112, 147-217(1995).
- [Lu1] Lusztig, G. *Vanishing properties of cuspidal local systems*. Proc. Nat. Acad. Sci. U.S.A. 91 (1994), no. 4, 14381439. 20G05 (22E46).
- [Lu2] George Lusztig. *Character sheaves V*. Advances in Mathematics 61.(1986), 103-155.
- [Lu3] George Lusztig. *Intersection cohomology complexes on a reductive group*. Invent. Math. 75(1984), no. 2, 205272.
- [Ma] Ruslan Maksimau. *Flag versions of quiver Grassmannians for Dynkin quivers have no odd cohomology over \mathbb{Z}* . arXiv:1909.04907v1 [math.RT]. 11 Sep 2019.
- [Sw] P. Slodowy. *Simple singularities and simple algebraic groups*. Lecture Notes in Mathematics(Springer, New York),Vol.815(1980).
- [Su] H. Sumihiro *Equivariant completion, I-II*, J.Math.Kyoto Univ.14 (1974), 1-28 and 15 (1975), 573-605.

Vita

Tamanna Chatterjee was born in a small town named Suri, Located in the eastern part of India . She earned her high school degree in 2010. In May 2013, she earned her Bachelor of Science degree in Mathematics from Vidyasagar college, which is affiliated under the University of Burdwan. She earned her Masters in Mathematics from Ramkrishna Mission Vivekananda Educational Research Institute(RKMVERI) in 2016. Chatterjee began her graduate studies at Louisiana State University in August 2016 and earned her Master of Science in Mathematics in May 2018. Currently she is a candidate for the degree of Doctor of Philosophy in Mathematics. In August 2021, she will begin a three year postdoctoral position at University of Georgia in mathematics.