

1. Benedetto, E., Sacerdote, L. and Zucca, C.: A first passage problem for a bivariate diffusion process: Numerical solution with an application to neuroscience when the process is Gauss-Markov, *J. Comput. Appl. Math.* 242 (2013), no. 1, 41-52.

<https://doi.org/10.1016/j.cam.2012.10.014>

2. Caravelli, F., Mansour, T., Sindoni, L. and Severini, S.: On moments of the integrated exponential Brownian motion, *Eur. Phys. J. Plus* 131 (2016), Article number 245.

<https://doi.org/10.1140/epjp/i2016-16245-9>

3. Cox, D. R. and Miller, H. D.: *The Theory of Stochastic Processes*, Methuen, London, 1965.

4. Goldman, M.: On the first passage of the integrated Wiener process, *Ann. Math. Statist.* 42 (1971), no. 6, 2150-2155.

<https://doi.org/10.1214/aoms/1177693084>

5. Gor'kov, Yu. P.: A formula for the solution of a certain boundary value problem for the stationary equation of Brownian motion, *Soviet. Math. Dokl.* 16 (1975), 904-908.

6. Hesse, C. H.: The one-sided barrier problem for an integrated Ornstein-Uhlenbeck process, *Stoch. Models* 7 (1991), no. 3, 447-480.

<https://doi.org/10.1080/15326349108807200>

7. Karatzas, I. and Shreve, S. E.: *Brownian Motion and Stochastic Calculus*, 2nd ed., Springer-Verlag, New York, 1991.

8. Lachal, A.: Sur le premier instant de passage de l'integrale du mouvement brownien, *Ann. Inst. H. Poincar'e Probab. Statist.* 27 (1991), no. 3, 385-405.

9. Lachal, A.: L'integrale du mouvement brownien, *J. Appl. Probab.* 30 (1993), no. 1, 17-27.

<https://doi.org/10.2307/3214618>

10. Lefebvre, M.: Moment generating function of a first hitting place for the integrated Ornstein-Uhlenbeck process, *Stochastic Process. Appl.* 32 (1989), no. 2, 281-287.

[https://doi.org/10.1016/0304-4149\(89\)90080-X](https://doi.org/10.1016/0304-4149(89)90080-X)

11. Lefebvre, M.: Integrated diffusion processes inside rectangles, *Bol. Soc. Mat. Mex.* 8 (2002), no. 3, 171-183.

12. Lefebvre, M.: First hitting time and place for the integrated geometric Brownian motion, *Int. J. Differ. Equ. Appl.* 9 (2004), no. 4, 365-374.

13. Lefebvre, M.: Moments of first-passage places and related results for the integrated Brownian motion, *ROMAI J.* 2 (2006), no. 2, 101-108.

14. Lefebvre, M. and Leonard, 'E.: On the first hitting place of the integrated Wiener process, *Adv. in Appl. Probab.* 21 (1989), no. 4, 945-948.

<https://doi.org/10.2307/1427780>

15. Levy, E.: On the moments of the integrated geometric Brownian motion, *J. Comput. Appl. Math.* 342 (2018), 263-273.

<https://doi.org/10.1016/j.cam.2018.04.005>

16. Makasu, C.: Exit probability for an integrated geometric Brownian motion, *Statist. Probab. Lett.* 79 (2009), no. 11, 1363-1365.

<https://doi.org/10.1016/j.spl.2009.02.009>

17. McKean, H. P.: A winding problem for a resonator driven by a white noise, *J. Math. Kyoto Univ.* 2 (1963), 227-235.

<https://doi.org/10.1215/kjm/1250524936>

18. Metzler, A.: The Laplace transform of hitting times of integrated geometric Brownian motion, *J. Appl. Probab.* 50 (2013), no. 1, 295-299.

<https://doi.org/10.1239/jap/1363784440>